

Scotland's Rural College

The effect of four different feeding regimes on rabbit behaviour

Prebble, JL; Langford, FM; Shaw, DJ; Meredith, AL

Published in:

Applied Animal Behaviour Science

DOI:

[10.1016/j.applanim.2015.05.003](https://doi.org/10.1016/j.applanim.2015.05.003)

Print publication: 01/01/2015

Document Version

Peer reviewed version

[Link to publication](#)

Citation for published version (APA):

Prebble, JL., Langford, FM., Shaw, DJ., & Meredith, AL. (2015). The effect of four different feeding regimes on rabbit behaviour. *Applied Animal Behaviour Science*, 169, 86 - 92.
<https://doi.org/10.1016/j.applanim.2015.05.003>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

1 The effect of four different feeding regimes on rabbit behaviour

2

3 **Jennifer L Prebble^{1,2,3}, Fritha M Langford⁴, Darren J Shaw¹ & Anna L Meredith¹**

4 ¹Royal (Dick) School of Veterinary Studies and the Roslin Institute, University of Edinburgh, Easter

5 Bush Campus, Midlothian, EH25 9RG¹

6 ²JP was employed on a KTP partnership between the Royal (Dick) School of Veterinary Studies and

7 Burgess Pet Care, Victory Mill, Priestman's Lane, Thornton-Le-Dale, Pickering, North Yorkshire, YO18

8 7RU

9 ³Current address; Askham Bryan College, Askham Bryan, York, YO23 3FR

10 ⁴ Animal and Veterinary Sciences, SRUC, West Mains Road, Edinburgh, EH9 3JG.

11

12 **Corresponding author**

13 **Professor Anna Meredith:** Anna.Meredith@ed.ac.uk

14

15 **Keywords**

16 **Rabbit; Feeding; Behaviour; Abnormal behaviour; Hay.**

17 Abstract

18 Dietary composition and presentation impacts on the behaviour of animals, and failure to provide a
19 suitable diet can lead to reduced welfare through the development of poor health, the inability to
20 express normal behaviours and the development of abnormal behaviours. This study assessed the
21 effects of two commonly fed pet rabbit diets (extruded nuggets with hay (EH) and muesli with hay
22 (MH)) alongside hay only (HO) and muesli only (MO) on the behaviour of 32 Dutch rabbits observed
23 over 17 months. Increased time spent feeding was observed in the groups fed ad libitum hay (HO,
24 EH, MH) compared to the MO group ($P < 0.05$). A corresponding high level of inactivity was observed
25 in the MO group compared to rabbits receiving hay ($P < 0.05$). In the groups provided with hay a
26 preference to consume hay in a natural grazing posture was observed. The higher activity levels and
27 absence of abnormal behaviours when hay was fed support recommendations that forage should
28 form a significant portion of the diet for domestic rabbits.

29 Introduction

30 As herbivores, wild rabbits consume relatively large amounts of a high fibre diet of low nutritional
31 quality (Williams and Wells, 1974). This requires them to apportion a large amount of their time
32 budget to grazing. Rabbits spend 30-70% of time outside the burrow grazing, pausing occasionally to
33 groom (Mykytowycz, 1958; Myers and Poole, 1961; Myers and Mykytowycz, 1958; Lockley, 1961).
34 Time spent eating varies with age, sex and social status within the group and has also been shown
35 to increase when food availability falls during drought (Myers and Mykytowycz, 1958; Mykytowycz,
36 1958). Grazing occurs mainly during late afternoon and throughout the night and daylight hours are
37 spent underground in warrens (Myers and Mykytowycz, 1958; Mykytowycz, 1958; Lockley, 1961;
38 Lockley, 1962). Caecotrophy is performed while underground (Southern, 1942). Domestic rabbits
39 kept in free range conditions exhibit a similar feeding pattern to their wild counterparts (Vastrade,
40 1987; Lehmann, 1991). In contrast, many pet rabbits are housed in small hutches with limited
41 exercise opportunities (Mullan and Main, 2006; PDSA, 2011) and a diet consisting largely of
42 concentrates (mono-component nugget or muesli mixes) (PDSA 2011) which can be consumed
43 rapidly (Lidfors, 1997), with limited or no access to hay or grass (Mullan and Main, 2006; PDSA,
44 2011).

45 Stereotypic behaviours are described as behaviours that are relatively invariant, regularly repeated
46 and without an obvious function (Mason, 1991). Stereotypic behaviours reported to occur in
47 laboratory rabbits include excessive grooming, sham chewing (chewing with nothing in mouth), bar
48 biting, licking parts of cage, digging against cage, biting water nipple, sliding nose against bars, head
49 pressing and running repeatedly in a defined pattern (Gunn and Morton, 1995; Lidfors, 1997). An
50 apathetic state of inactivity and boredom has also been reported by Gunn and Morton (1995).
51 Stereotypic behaviours occur most frequently during the night (Gunn and Morton, 1995) when
52 rabbits are naturally at their most active (Mykytowycz 1958).

53 Whilst not studied in pet rabbits, the beneficial impact of providing hay to laboratory rabbits has
54 been demonstrated (Lidfors, 1997; Berthelsen and Hansen, 1999) . The provision of hay to
55 individually housed laboratory rabbits has proved effective at reducing the expression of abnormal
56 behaviours (Lidfors, 1997; Berthelsen and Hansen, 1999).

57 Rabbits can consume pelleted feeds rapidly (Lidfors, 1997) and, whilst they may provide adequate
58 nutrition for the maintenance of the rabbit, foraging behaviour is limited. If fed in limited amounts
59 the rapid consumption of the daily ration may leave the rabbit in a state of hunger for a considerable
60 portion of the day (Lidfors, 1997). It has been suggested that stereotypies in pigs and broiler
61 breeder chickens develop through hunger and frustration at an inability to forage (Lawrence and
62 Terlouw, 1993; de Jong et al., 2003; de Jong et al., 2005) when a restricted diet is fed. Rabbits
63 provided with hay spend considerably more time interacting with it than with other forms of
64 environmental enrichment, suggesting its importance to this species (Lidfors, 1997).

65 Despite recommendations that the ideal diet for pet rabbits is one of grasses, herbs, and leaves
66 mimicking that of their wild counterparts (Clauss, 2012), the benefits of hay for meeting behavioural
67 needs of rabbits , and that rabbits are able to maintain weight gain on forage only diets ([Lebas,
68 2004](#); [Leiber et al., 2008](#)), studies suggest that at least 15-17% of pet rabbits do not have access to
69 hay (Mullan and Main 2006; Schepers et al. 2009) and 36-42% are not fed recommended amounts
70 (PDSA 2011; 2012). Veterinary surgeons frequently recommend feeding rabbits concentrates in
71 limited amounts, as it is recognised that owners may often feed what may be considered excessive
72 amounts of concentrate diets ([Harcourt-Brown, 2002a](#); [Meredith, 2006](#); [PDSA, 2011](#)) but it is unclear
73 as to whether ad libitum (*ad lib*) access to hay is routinely recommended.

74 This study aimed to assess the effect of two commonly fed/recommended diet regimes (extruded
75 nugget with *ad lib* hay and a mixed muesli type diet with *ad lib* hay), alongside a forage based diet
76 and a muesli only diet, on the feeding behaviour and time budget of pet rabbits.

77 Materials and methods

78 Study animals

79 This study was conducted as part of a long term study to assess the effect of diet on the health and
80 welfare of pet rabbits, as previously described by Prebble and Meredith (2014). Thirty two Dutch
81 rabbits from five mixed litters (20 male and 12 female) were purchased at 8-9 weeks old from a
82 single breeder. They were weighed on arrival (mean weight 0.84 kg±0.084). The rabbits were housed
83 in 12 male-female neutered pairs and four male-male neutered pairs in wooden floor pens (0.96m²)
84 with 12mm thick rubber matting, with shavings provided as bedding. The pens were split between
85 two rooms with a 12hr light (0600-1800): 12 hour dark cycle, a temperature of 18°C (±2°C) and
86 relative humidity between 40 and 70%. Rabbits were randomly allocated in pairs to four diet
87 treatment groups and day 0 was designated as the day when the rabbits had been transitioned
88 completely onto the treatment diet after an acclimatisation period. Detailed consideration of the
89 experimental setup of the study with regard to the design and housing was undertaken and
90 approved by the Ethical Review Committees of the Royal (Dick) School of Veterinary Studies and the
91 Food and Environment Research Agency (FERA). The rabbits were housed in a facility licensed by the
92 Home Office, however a project licence under the Animals (Scientific Procedures) Act 1986 (ASPA)
93 was not required for this study. The study was continually monitored by the FERA Ethics Committee
94 and Home Office inspector throughout its duration.

95 In the week following arrival six rabbits displayed signs of digestive disease, subsequently diagnosed
96 as an outbreak of coccidiosis and clostridial enterotoxaemia. Supportive treatment was provided.
97 Despite this, three rabbits died, but three recovered over the following 2 weeks. All remaining
98 rabbits then received prophylactic treatment with a two day course of toltrazuril (Baycox 50 mg/ml
99 Oral Suspension for Piglets, Calves and Lambs, Bayer plc, Newbury, Berkshire, UK) at a dose rate of
100 2.5mg/kg ([Redrobe et al., 2010](#)) on day -43 and -42 and repeated five days later on day -37 and -36
101 and metronidazole (Flagyl S 200mg/5ml Oral Suspension, Winthrop Pharmaceuticals UK Limited,

102 Guildford, Surrey, UK) at a dose rate of 20mg/kg twice daily for five days. Three 8 week old Dutch
103 rabbits arrived on day -23 to replace those that died.

104

105 Diets

106 On arrival, rabbits were acclimatised over a period of 40 days (days -54 to -14) by maintaining their
107 weaning diet and were fed 50g per rabbit of an extruded diet (Burgess® Excel-Junior and Dwarf
108 Rabbit; Burgess Pet Care, Thornton Le Dale, North Yorkshire, UK) once a day plus *ad lib* Timothy Hay.
109 Hay was provided in wall mounted hayracks to enable intake to be monitored and to prevent faecal
110 and urinary contamination which may reduce intake of hay. Water was provided *ad lib* in 700ml
111 bottles. At day-14 the paired rabbits were allocated to one of four diet treatment groups:

112 1 - Hay Only (HO) - *ad lib* supply of Timothy Hay (n=8);

113 2 – Extruded diet and Hay (EH) - 50g per rabbit Burgess Excel- Adult Rabbit (Burgess Pet Care,
114 Thornton Le Dale, North Yorkshire, UK) with *ad lib* supply of hay (n=8);

115 3 - Muesli and Hay (MH) - 60g per rabbit Russell Rabbit Complete Muesli (Supreme Petfoods Limited,
116 Ipswich, Suffolk, UK) with *ad lib* hay (n=8);

117 4 - Muesli Only (MO) - *ad lib* supply (125g per rabbit) of Russell Rabbit Complete Muesli (Supreme
118 Petfoods Limited, Ipswich, Suffolk, UK) (n=8).

119 Rabbits were gradually transitioned on to their respective new diets over a two week period (day -14
120 to day 0) to prevent digestive conditions associated with sudden dietary changes (Tzika et al. 2004).

121 From day 0 to the end of the study, (day 510; 17 months), rabbits were only fed the diet of that
122 group.

123 The nutritional compositions of diets are shown in Table 1. The EH and MH diets represent two
124 commonly fed diets fed according to the manufacturer's guidelines. The muesli consisted of 11

125 components: extrudates (4 types), pellets (2 types), grains (3 types), rolled peas and alfalfa stalks.
126 Quantities of concentrates offered in the EH and MH groups were based on the lower end of the
127 range of the manufacturer's stated guidelines to replicate dietary advice given by veterinary
128 surgeons (Harcourt-Brown 2002a; Meredith 2006). The inclusion of the HO group was to provide a
129 forage only diet similar to that of wild rabbits. The MO group was included because many
130 commercially available muesli based diets are labelled as complete or 'nutritionally complete',
131 leading owners to feed them alone. Muesli was provided to the MO group in sufficient quantities to
132 ensure an ad libitum supply of food. All concentrates were weighed out and replaced daily to ensure
133 accurate and consistent weights were offered. No measures were taken to prevent selective sorting
134 or feeding of the different components of the muesli.

135 In the month following transition all rabbits were neutered and vaccinated against myxomatosis
136 (Nobivac Myxo, MSD Animal Health, Milton Keynes, UK) and Viral Haemorrhagic Disease (Cylap,
137 Pfizer Limited, Sandwich, UK) as recommended for pet rabbits at that age.

138 The rabbits continued on the four trial diets for 17 months (72 weeks). One pair of rabbits in the MO
139 group had to be separated in week 3 following the development of aggressive behaviour and are not
140 included in the analysis.

141 Data collection

142 Behavioural observations were performed over two 24 hour periods in week 8 (timepoint 1: T1) and
143 week 18 (timepoint 2: T2) of the trial diet period. A video recording using closed circuit television
144 (CCTV was made for 24 hours from 0900h. Daily food rations (concentrates and/or hay) and water
145 were provided prior to the start of recording, following which no one entered the room for the 24
146 hour period. Red light was used during the 12h dark period which was present for 28 days prior to
147 data collection to allow habituation. Recordings were carried out with WebCCTV NVR (Quadrox,
148 Herent, Belgium).

149

150 Instantaneous scans of the CCTV recordings were performed every 15 minutes and the behaviour
151 performed by each of the rabbits in the pen recorded. Data was collected by pen and
152 measurements relating to bout duration were not recorded as rabbits were housed in pairs and
153 could not be distinguished from each other on recordings. At both timepoints (T1 and T2) the first
154 observations were in sequence from the lowest to the highest pen numbers. Individual behaviours
155 were categorised into five groups (Feeding, Maintenance, Active, Inactive and Investigative) for
156 analysis.

157 An ethogram based on Gunn and Morton (1993) was developed following prior observation of the
158 rabbits and is detailed in Table 2. In addition, assessment of proximity was made during each
159 instantaneous scan. Proximity was described as rabbits either separated by more than a third of the
160 pen (apart), in the same third of the pen without touching (near to each other) , or in direct physical
161 contact (together).

162

163 Statistical Analysis

164 Analysis of the data by behaviour group (Table 2) was performed on the total data set and also
165 separately on data from light and dark periods. Behaviours in the feeding category were also
166 analysed individually. The effects of sex on behavioural expression could not be assessed as it was
167 not possible to distinguish between individual rabbits. Statistical analysis was performed using R
168 software (v3.1.2 © 2014 The R Foundation for Statistical Computing and the R package *lme4* v 1.1-7).
169 Overall percentages of time spent performing each behaviour when both time points were
170 considered together were analysed using generalised linear mixed-effect models with binomial
171 errors (*glmeb*) with Pen as the random effect. *Glmeb* models were also used to compare how hay
172 was consumed, and whether rabbits differed in how much they stayed physically closer together
173 between the 2 timepoints. When considering differences in behaviour between diet treatment

174 groups at specific time points, general linear models with binomial errors (*glmb*) were utilised.
175 Tukey's post-hoc tests were carried out for both sets of models to assess pair-wise differences
176 between groups where overall differences were found. Pearson's correlation was used to examine
177 relationships between behaviours. $P < 0.05$ was taken to indicate statistical significance and mean \pm
178 standard errors are quoted throughout.

179

180 **Results**

181 Figure 1 shows the percentage of time spent performing different behaviours at the two timepoints
182 by diet treatment group. Overall, *feeding* was negatively correlated with *inactivity* at both time
183 points (T1: $\rho = -0.892$, T2: $\rho = -0.918$, $P < 0.001$) and *maintenance* behaviours at timepoint 2 ($\rho = -0.547$,
184 $P = 0.035$).

185 Effect of dark/ light period on behaviour

186 Differences in percentage of time spent performing different behaviours were affected by the light
187 dark cycle (Figure 2). *Feeding* (Dark: mean = $32.44\% \pm 2.84$, Light: $28.54\% \pm 4.68$), *maintenance*
188 (Dark: $9.76\% \pm 1.34$, Light: $5.93\% \pm 1.06$) and *investigative* (Dark: $3.15\% \pm 0.69$, Light: $1.5\% \pm 0.29$)
189 behaviours occurred more frequently in the dark period (Overall: $P < 0.001$, T1: $P < 0.019$, T2: $P < 0.013$)
190 whilst *inactive* (Dark: $52.31\% \pm 2.29$, Light: $63.20\% \pm 3.90$) behaviours occurred more frequently in the
191 light period ($P < 0.001$ for overall, T1 and T2).

192 In the light period, the HO group spent more time *feeding* (overall $49.7\% \pm 2.4$) in comparison to
193 9.3% (± 3.5) of time by the MO group ($P < 0.001$ for overall, T1 and T2), and the HO group spent less
194 time *inactive* (overall $45.8\% \pm 3.3$) than the MO group (overall $83\% \pm 6.2$, $P < 0.001$ for overall, T1 and
195 T2). No differences between the 4 groups in percentage of time spent performing *maintenance* and
196 *investigative* behaviours were present in the light period ($P > 0.085$ for overall, T1 and T2). In
197 contrast, *maintenance* behaviours were performed more frequently in the MO group than all other

198 groups in the dark period overall ($P<0.029$) and T1 ($P<0.037$), but only the HO group spent less in
199 *maintenance* than MO at timepoint 2 ($P<0.001$, $P>0.120$ other groups) (Figure 2).

200

201 Effect of diet group on behaviour

202 Over the total twenty four hour period (combining light and dark periods) at each timepoint, the MO
203 group spent significantly less percentage time *feeding* (T1:11.33% \pm 4.33, T2:9.33% \pm 4.68) and more
204 time *inactive* (T1:71% \pm 4.02, T2:74.67% \pm 3.35) than all other groups ($P<0.006$, Figure 1). The HO
205 group also spent significantly more percentage time *feeding* (T1:44.25% \pm 1.60, T2:40.75% \pm 1.31)
206 than the MH (T1:26% \pm 1.68, T2:23.5% \pm 1.44) group at both timepoints ($P<0.001$). The EH group
207 spent significantly less percentage time *feeding* (T1:26% \pm 1.68, T2:23.5% \pm 1.44) than the HO group
208 ($P<0.001$), however there was no significant difference in levels of *inactivity* between these two
209 groups (53% \pm 2.42) at time point 1 ($P=0.846$). At time point 2 the EH group spent significantly more
210 percentage time *inactive* (59.25% \pm 1.70, than the HO group ($P<0.002$). Compared to the EH group,
211 the MH group spent significantly less percentage time *feeding* and more time *inactive* at timepoint 1
212 ($P<0.001$), but there was no significant difference spent *inactive* at timepoint 2 ($P>0.09$).

213 Consumption of hay from between the bars, with the front paws placed on the hayrack (T1:16.38%
214 \pm 5.75, T2:26.45% \pm 14.9) occurred significantly less than consumption of hay whilst in a natural
215 grazing posture from either the floor or whilst sat in the hay rack (T1:83.62% \pm 5.75, T2:73.55%
216 \pm 14.9, $P<0.001$, Figure 3).

217 Observations of *maintenance* behaviours were significantly higher in the MO group (T1:11.67%
218 \pm 1.47, T2:12.67% \pm 1.89) than in the HO group (T1:5% \pm 0.41, T2:6% \pm 1.35) at both timepoints
219 ($P<0.001$) and the EH group (T1:7% \pm 1.37, T2:9.75% \pm 2.14) at timepoint 1 ($P=0.027$). *Maintenance*
220 behaviours occurred significantly more frequently in the MH group (T1:9% \pm 2.86, T2:8.25% \pm 1.7)

221 than the HO group at timepoint 1 ($P<0.02$) but not timepoint 2 ($P=0.391$). While at timepoint 2
222 *maintenance* behaviours were significantly greater in the MO group than the MH group ($P=0.038$).
223 *Active* behaviours were observed significantly more frequently in the MH group (T1:2.5% \pm 0.65,
224 T2:2% \pm 0.58) than the HO group (T1:0.5% \pm 0.29, T2:1.75% \pm 0.85) only at timepoint 1 ($P=0.018$).
225 *Investigative* behaviour occurred more frequently in the MO group (T1:4.67% \pm 0.63, T2:2% \pm 0.48)
226 than the HO group again only at timepoint 1 (T1:1.25% \pm 0.25, T2:2% \pm 0.41, $P=0.001$).

227

228 In all groups, time spent in direct physical contact (together) was greater at T2 (38.73% \pm 2.37) than
229 at T1 (10.13% \pm 1.58, Figure 4, $P<0.001$). The HO group spent less time together (T1:4% \pm 1.08, T2:
230 27.75% \pm 3.09) than the MH (T1:14.25% \pm 8.76, T2:46.5% \pm 7.44) and MO (T1:15.33% \pm 6.22, T2:58%
231 \pm 6.08) groups at both timepoints ($P<0.015$). The MH and MO groups spent a greater proportion of
232 time together than the EH (T1:8.25% \pm 2.78, T2:27.5% \pm 4.17) group only at the second timepoint
233 ($P<0.001$). In addition the MO group spent more time together than the MH group only at timepoint
234 2 ($P=0.019$)

235

236 Other observations

237 Throughout the study, rabbits from MH, HO and EH groups were observed pulling hay from the hay
238 rack using a digging motion with their front paws and subsequently consuming hay from the floor.
239 In addition, rabbits in the majority of pens also jumped into the hay rack to consume hay.

240 Over the whole study period (17 months), hair chewing was recorded in one rabbit within the MO
241 group and occurred in the period following the removal of its pen mate from the trial. Chewing of
242 objects in the pen was seen infrequently on the videos (0.14% of observations), however damage to
243 wooden fixtures within the pen as a result of chewing occurred in all groups over the whole study

244 period. Chewing of the rubber matting used as flooring was observed only in the MO group. In
245 addition to chewing the rubber matting, rabbits in the MO group were also often observed eating
246 the shavings provided as bedding over the duration of the entire 17 month trial.

247 **Discussion**

248 This study has provided preliminary information on the effect of diet on the time budgets of pet
249 rabbits. Diet affects time budgets of many herbivorous mammals including horses, sheep and pigs
250 (Ruckebusch and Gaujoux, 1976; Robert et al., 1993; Thorne et al., 2005). Previous studies in rabbits
251 have been limited to laboratory rabbits which focus on environmental enrichment including the use
252 of hay and food based enrichments rather than their use as a diet (Lidfors, 1997; Berthelsen and
253 Hansen, 1999; Harris et al., 2001), whereas this study assessed the effect of different feeding
254 regimes on the time budgets of pet rabbits.

255 Time spent feeding was highest in the HO group which had the diet of lowest nutritional value. Wild
256 rabbits spend between 30-70% of their time above ground eating (Myers and Mykytowycz, 1958;
257 Mykytowycz, 1958) but this can rise to 90% when diet quality is poor (Myers and Poole, 1961). As
258 wild rabbits are above ground between 11 and 13 hours per day (Mykytowycz, 1958) this equates to
259 a range of approximately 14 - 38% of total time spent eating. The time spent feeding in the groups
260 provided with hay (HO, EH and MH) groups falls within this range at both timepoints, however the
261 average time spent feeding by the HO group was at the top end of this range and is similar to figures
262 reported by Myers and Poole (1961) in rabbits during summer when diet quality is of low nutritional
263 value. Conversely the MO group spent less time feeding (9-11%) than wild rabbits. Reduced
264 opportunity for foraging in herbivores, including rabbits, has been associated with the development
265 of abnormal and stereotypical behaviours (Robert et al., 1993; Lidfors, 1997; Berthelsen and Hansen,
266 1999; Thorne et al., 2005). These may occur as a result of boredom, hunger or frustrated attempts
267 to display foraging behaviours which they are motivated to perform (Newberry, 1995; D'Eath et al.,
268 2009).

269 The reduction in time spent feeding was matched with corresponding increases in time spent
270 inactive; a similar pattern of behaviour is also reported in sheep (Ruckebusch and Gaujoux, 1976).
271 The increase in inactivity and reduction in active feeding behaviours may have contributed to both
272 the development of abnormal behaviours (see below) and high weight gain and increased body
273 condition observed in the MO group (Prebble and Meredith, 2014).

274 Stereotypy has been reported to occur at a mean frequency of 11% over a 24 hr period in laboratory
275 rabbits (Gunn and Morton, 1995). Stereotypical behaviours include hair chewing, chewing objects,
276 nose sliding and licking objects (Gunn and Morton, 1995). Hair chewing along with other abnormal
277 behaviours are suggested as indicators of stress (Podberscek et al., 1991; Gunn and Morton, 1995)
278 and therefore in the one rabbit in which hair chewing was observed, this could have been triggered
279 by the loss of the companion, as rabbits are highly motivated to have social contact (Seaman et al.,
280 2008). The lack of access to hay may also have contributed to the hair chewing in this rabbit, as
281 provision of hay has been demonstrated to prevent hair chewing in laboratory rabbits (Mulder et al.,
282 1992). Although chewing of wood occurred in all diet groups in this study, the chewing of rubber
283 matting and ingesting of shavings in the MO group could be indicative of abnormal behaviour. The
284 chewing of rubber matting may be misdirected chewing as a consequence of reduced time feeding.
285 Increased levels of abnormal behaviours have been found in pigs that do not have a suitable
286 substrate for foraging (Spooler et al. 1995). This would indicate that the rabbits needed or craved a
287 higher fibre food source and is a behaviour also reported in horses fed on a low forage diet (Houpt et
288 al. 1988).

289

290 The level of potential stereotypy and abnormal behaviour is lower in this study than that reported by
291 Gunn and Morton (1995) and this may reflect differences in the rabbit's husbandry. Rabbits in this
292 study were housed in pairs whereas rabbits studied previously were housed in social isolation which
293 increases expression of abnormal behaviours (Chu et al. 2004). The observations of Gunn and

294 Morton (1995) were also performed in laboratory cages which limited behaviours such as rearing,
295 hoping and lying outstretched. Increasing the amount of space available reduces inactivity and
296 allows expression of a fuller behavioural repertoire in rabbits (Dixon et al. 2010) and reduces
297 abnormal behaviour (Chu et al. 2004). Therefore the social contact and increased space (allowing
298 expression of most behaviours) provided in this study may have contributed to the low overall levels
299 of abnormal behaviours seen. In addition, the use of instantaneous scans rather than continuous
300 monitoring may have reduced detection of misdirected chewing behaviour as it occurs as an event
301 rather than a state.

302

303 The higher proportion of time spent consuming hay from the floor or whilst sitting in the hayrack
304 suggests that rabbits prefer to consume hay in a natural grazing posture with the head down rather
305 than from a hay rack which requires elevation of the head. Similar findings have been observed in
306 horses which have been observed actively pull hay onto the floor and spending a higher proportion
307 of time consuming it from the floor than from troughs (Sweeting et al., 1985).

308

309 The findings of this study highlight the effect of forage on behaviour in pet rabbits. The increased
310 inactivity and reduction in time spent eating in rabbits on the muesli only diet indicates an impact on
311 welfare through the inability to express normal foraging behaviour (evidenced by misdirected
312 chewing of rubber matting and consumption of bedding), in addition to the development of obesity,
313 also reported elsewhere in the MO rabbits (Prebble et al., 2014) through reduced activity levels.

314 Provision of forage is essential to promote welfare of rabbits and should form a significant portion of
315 the rabbit's diet.

316

317

318 Acknowledgements

319 The authors acknowledge financial support from the Technology Strategy board provided through a
320 Knowledge Transfer Partnership between the Royal (Dick) School of Veterinary Studies and Burgess
321 Pet Care and also thank Burgess Pet Care for additional financial support. We also thank FERA for
322 allowing us the use of their facilities, ASIST for providing routine rabbit care and Station House Vets
323 for neutering and providing treatment to the rabbits.

324 **References**

- 325 Berthelsen, H., Hansen, L.T., 1999: The Effect of Hay on the Behaviour of Caged Rabbits (*Oryctolagus*
326 *Cuniculus*). *Animal Welfare* **8**, 9.
- 327 Chu, L., Garner J.P., Mench, J.A., 2004: A behavioural comparison of New Zealand White rabbits
328 (*Oryctolagus cuniculus*) housed individually or in pairs in conventional laboratory cages. *Applied*
329 *Animal Behaviour Science* **85**, 121-139.
- 330 Clauss, M., 2012: Clinical technique: feeding hay to rabbits and rodents. *Journal of Exotic Pet*
331 *Medicine* **21**, 80-86.
- 332 D'Eath, R.B. Tolkamp, B.J. Kyriazakis, I. Lawrence, A.B., 2009: 'Freedom from hunger' and preventing
333 obesity: the animal welfare implications of reducing food quantity or quality. *Animal Behaviour* **77**,
334 275-288.
- 335 de Jong, I. C., Sander van Voorst, A., Blokhuis, H.J., 2003: Parameters for quantification of hunger in
336 broiler breeders. *Physiology and Behaviour* **78**, 773-783.
- 337 de Jong, I.C., Enting, H., van Voorst, A., Blokhuis, H.J., 2005: Do Low-Density Diets Improve Broiler
338 Breeder Welfare During Rearing and Laying? *Poultry Science* **84**, 194-203.
- 339 Dixon, L.M., Hardiman, J.R., Cooper, J.J., 2010: The effects of spatial restriction on the behavior of
340 rabbits (*Oryctolagus cuniculus*). *Journal of Veterinary Behavior* **5**, 302-308.
- 341 Gunn, D., Morton, D.B., 1995: Inventory of the behaviour of New Zealand White rabbits in laboratory
342 cages. *Applied Animal Behaviour Science* **45**, 277-292.
- 343 Harcourt-Brown, F., 2002: Diet and husbandry, In: Harcourt-Brown, F. (ed.), Textbook of rabbit
344 medicine. Butterworth-Heinemann, Oxford.
- 345 Harris, L.D., Custer, L.B., Soranaka, E.T., Burge, J.R., Ruble, G.R., 2001: Evaluation of objects and food
346 for environmental enrichment of NZW rabbits. *Contemp Top Lab Anim Sci* **40**, 27-30.
- 347 Houpt, K.A., Perry, P.J., Hintz, H.F., Houpt, T.R., 1988: Effect of meal frequency on fluid balance and
348 behavior of ponies. *Physiology and Behaviour* **42**, 401-407.

349 Lawrence, A.B., Terlouw, E.M., 1993: A review of behavioural factors involved in the development
350 and continued performance of stereotypic behaviours in pigs. *Journal of Animal Science* **71**, 2815-
351 2825.

352 Lebas, F., 2004: Reflections on rabbit nutrition with a special emphasis on feed ingredients
353 utilization, 8th World Rabbit Congress, Puebla, Mexico.

354 Lehmann, M., 1991: Social behaviour in young domestic rabbits under semi natural conditions.
355 *Applied Animal Behaviour Science* **32**, 269-292.

356 Leiber, F., Meier, J.S., Burger, B., Wettstein, H.-R., Kreuzer, M., Hatt, J.-M., Clauss, M., 2008:
357 Significance of coprophagy for the fatty acid profile of rabbits fed different diets. *Lipids* **43**, 853-865.

358 Lidfors, L., 1997: Behavioural effects of environmental enrichment for individually caged rabbits.
359 *Applied Animal Behaviour Science* **52**, 157-169.

360 Lockley, R.M., 1961: Social Structure and stress in the rabbit warren. *Journal of Animal Ecology* **30**,
361 385-425.

362 Lockley, R.M., 1962: Production of faecal Pellets in the wild rabbit. *Nature* **194**, 988-989.

363 Mason, G.J., 1991: Stereotypies: a critical review. *Animal Behaviour* **41**, 1015-1037.

364 Meredith, A., 2006: General Biology and Husbandry, In: Meredith, A. (ed.), BSAVA Manual of Rabbit
365 Medicine and Surgery. British Small Animal Veterinary Association. 1-17.

366 Mulder, A., Nieuwenkamp, A.E., van der Palen, J.G., van Rooijen, G.H., Beynen, A.C., 1992:
367 Supplementary hay reduces fur chewing in rabbits (abstract only). *Tijdschrift voor diergeneeskunde*
368 **12**, 655-658.

369 Mullan, S.M., Main, D.C., 2006: Survey of the husbandry, health and welfare of 102 pet rabbits.
370 *Veterinary Record* **159**, 103-109.

371 Myers, K., Mykytowycz, R., 1958: Social Behaviour in the Wild Rabbit. *Nature* **181**, 1515-1516.

372 Myers, K., Poole, W.E., 1961: A study of the biology of the wild rabbit, *Oryctolagus cuniculus* (L.), in
373 confined populations II. The effects of season and population increase on behaviour. *CSIRO Wildlife*
374 *Research* **6**, 1-41.

375 Mykytowycz, R., 1958: Continuous observations of the activity of the wild rabbit, *Oryctolagus*
376 *cuniculus* (L.), during 24-hour periods. *C.S.I.R.O. Wildl. Res.*

377 Newberry, R.C., 1995: Environmental enrichment: Increasing the biological relevance of captive
378 environments. *Applied Animal Behaviour Science* **44**, 229-243.

379 PDSA, 2011: PDSA Animal Wellbeing Report 2011.

380 PDSA, 2012: PDSA Animal Wellbeing Report 2012: Issues in Focus.

381 Prebble, J.L., Meredith A.L., 2014. Food and water intake and selective feeding in rabbits on four
382 feeding regimes. *Journal of Animal Physiology and Animal Nutrition* **98**, 991-1000.

383 Prebble, J.L., Shaw, D.J., Meredith, A.L. 2014a: Bodyweight and body condition score in rabbits on
384 four different feeding regimes. [Journal of Small Animal Practice](#). doi: 10.1111/jsap.12301. [Epub
385 ahead of print]

386 Podberscek, A.L., Blackshaw, J.K., Beattie, A.W., 1991: The behaviour of group penned and
387 individually caged laboratory rabbits. *Applied Animal Behaviour Science* **28**, 353-363.

388 Redrobe, S.P., Gakos, G., Elliot, S.C., Saunders, R., Martin, S., Morgan E.R., 2010: Comparison of
389 toltrazuril and sulphadimethoxine in the treatment of intestinal coccidiosis in pet rabbits. *Veterinary*
390 *Record* **167**, 283-290.

391 Robert, S., Matte, J.J., Farmer, C., Girard, C.L., Martineau, G.P., 1993: High-fibre diets for sows:
392 effects on stereotypies and adjunctive drinking. *Applied Animal Behaviour Science* **37**, 297-309.

393 Ruckebusch, Y., Gaujoux, M., 1976: Sleep-inducing effect of a high protein diet in sheep. *Physiology*
394 *and Behaviour* **17**, 9-12.

395 Schepers, F., Koene, P., Beerda, B., 2009: Welfare assessment in pet rabbits. *Animal Welfare* **18**, 477-
396 485.

397 Seaman, S.C., Waran, N.K., Mason, G., D'eath, R.B., 2008: Animal Economics: assessing the motivation
398 of female laboratory rabbits to reach a platform, social contact and food. *Animal Behaviour* **75**, 31-
399 42.

400 Southern, H.N., 1942: Periodicity of refectation in the wild rabbit. *Nature* **149**, 553-554.

401 Spoolder, H.A.M., Burbridge, J.A., Edwards, S.A., Simmins, P.H., Lawrence, A.B., 1995: Provision of
402 straw as a foraging substrate reduces the development of excessive chain and bar manipulation in
403 food restricted sows. *Applied Animal Behaviour Science* **43**, 249-262.

404 Sweeting, M.P., Houpt, C.E., Houpt, K.A., 1985: Social Facilitation of feeding and time budgets in
405 stabled ponies. *Journal of Animal Science* **60**, 369-374.

406 Thorne, J.B., Goodwin, D., Kennedy, M.J., Davidson, H.P.B., 2005: Foraging enrichment for
407 individually housed horses: Practicallity and effects on behaviour. *Applied Animal Behaviour Science*
408 **94**, 149-164.

409 Tzika, E.D., Papatsiros, V., Tassis, P.T., Saoulidis, K., 2004: Prevalance of Enteritis in Greek Rabbitries,
410 International Society for Animal Hygiène Saint-Malo.

411 Vastrade, F.M., 1987: Spacing behaviour of free-ranging domestic rabbits, *Oryctolagus cuniculus* L.
412 *Applied Animal Behaviour Science* **18**, 185-195.

413 Williams, O.B., Wells, T.C.E., 1974: Grazing management of Woodwalton Fen:Seasonal changes in
414 the diet of cattle and rabbits. *Journal of Applied Ecology* **11**, 499-516.

415

416

417

418 **Table 1.** Nutritional composition of diets offered, with values expressed as % Dry Matter (DM)

419

420 **Table 2.** Rabbit Ethogram used in this study (based on Gunn and Morton (1993))

421

422 **Figure 1.** Time budgets of Dutch rabbits fed on four diets, hay only (HO), Extruded nugget and hay
423 (EH), Muesli and hay (MH) and Muesli only (MO). Percentage of observations consisting of feeding,
424 inactive, maintenance and other (made up of active and investigative behaviours) behaviours for a
425 24 hour period at timepoint 1 (week 8) and timepoint 2 (week 18) after the start of the dietary
426 treatments.

427

428 **Figure 2.** Time budgets of Dutch rabbits fed on four diets, hay only (HO), Extruded nugget and hay
429 (EH), Muesli and hay (MH) and Muesli only (MO). Percentage of observations consisting of feeding,
430 inactive, maintenance and other (made up of active and investigative behaviours) behaviours for the
431 light and dark periods (12 hour light dark cycle) at timepoint 2 (week 18) after the start of the dietary
432 treatments.

433

434 **Figure 3.** Feeding behaviour of Dutch rabbits fed on four diets, hay only (HO), Extruded nugget and
435 hay (EH), Muesli and hay (MH) and Muesli only (MO). Breakdown of the feeding behaviours
436 observed at timepoint 1 (week 8) and timepoint 2 (week 18) after the start of the dietary
437 treatments.

438

439 **Figure 4.** Proximity of paired Dutch rabbits to each other. Rabbits fed on one of four diets, hay only
440 (HO), Extruded nugget and hay (EH), Muesli and hay (MH) and muesli only (MO). Percentage of time
441 spent separated by more than a third of the pen, in the same third of the pen without touching, or in
442 direct physical contact over a 24hour period at timepoint 1 (week 8) and timepoint 2 (week 18) after
443 the start of the dietary treatments.