

Scotland's Rural College

## Genetic profile of scrapie codons 146, 211 and 222 in the PRNP gene locus in three breeds of dairy goats

Vouraki, S; Gelasakis, AI; Alexandri, P; Boukouvala, E; Ekateriniadou, LV; Banos, G; Arsenos, G

Published in:  
PLoS ONE

DOI:  
[10.1371/journal.pone.0198819](https://doi.org/10.1371/journal.pone.0198819)

First published: 07/06/2018

Document Version  
Peer reviewed version

[Link to publication](#)

### Citation for published version (APA):

Vouraki, S., Gelasakis, AI., Alexandri, P., Boukouvala, E., Ekateriniadou, LV., Banos, G., & Arsenos, G. (2018). Genetic profile of scrapie codons 146, 211 and 222 in the PRNP gene locus in three breeds of dairy goats. *PLoS ONE*, 13(6), Article 0198819. Advance online publication. <https://doi.org/10.1371/journal.pone.0198819>

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

### Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

1 **Genetic profile of scrapie codons 146, 211 and 222 in the *PRNP* gene locus in**  
2 **three breeds of dairy goats**

3

4 Sotiria Vouraki<sup>1\*</sup>, Athanasios I. Gelasakis<sup>2</sup>, Panoraia Alexandri<sup>1</sup>, Evridiki  
5 Boukouvala<sup>2</sup>, Loukia V. Ekateriniadou<sup>2</sup>, Georgios Banos<sup>1,3</sup>, Georgios Arsenos<sup>1</sup>

6

7 <sup>1</sup>Laboratory of Animal Husbandry, School of Veterinary Medicine, Faculty of Health  
8 Sciences, Aristotle University of Thessaloniki, Thessaloniki, Greece

9 <sup>2</sup>Veterinary Research Institute of Thessaloniki, Hellenic Agricultural Organization  
10 Demeter, Thessaloniki, Greece

11 <sup>3</sup>Scotland's Rural College and The Roslin Institute, University of Edinburgh,  
12 Scotland, UK

13

14 \*Corresponding author

15 Email: [svouraki@vet.auth.gr](mailto:svouraki@vet.auth.gr) (SV)

16

## 17 **Abstract**

18 Polymorphisms at *PRNP* gene locus have been associated with resistance against  
19 classical scrapie in goats. Genetic selection on this gene within appropriate breeding  
20 programs may contribute to the control of the disease. The present study characterized  
21 the genetic profile of codons 146, 211 and 222 in three dairy goat breeds in Greece. A  
22 total of 766 dairy goats from seven farms were used. Animals belonged to two  
23 indigenous Greek, Eghoria (n=264) and Skopelos (n=287) and a foreign breed,  
24 Damascus (n=215). Genomic DNA was extracted from blood samples from individual  
25 animals. Polymorphisms were detected in these codons using Real-Time PCR  
26 analysis and four different Custom TaqMan<sup>®</sup> SNP Genotyping Assays. Genotypic,  
27 allelic and haplotypic frequencies were calculated based on individual animal  
28 genotypes. Chi-square tests were used to examine Hardy-Weinberg equilibrium state  
29 and compare genotypic distribution across breeds. Genetic distances among the three  
30 breeds, and between these and 30 breeds reared in other countries were estimated  
31 based on haplotypic frequencies using fixation index  $F_{ST}$  with Arlequin v3.1 software;  
32 a Neighbor-Joining tree was created using PHYLIP package v3.695. Level of  
33 statistical significance was set at  $P=0.01$ . All scrapie resistance-associated alleles  
34 (146S, 146D, 211Q and 222K) were detected in the studied population. Significant  
35 frequency differences were observed between the indigenous Greek and Damascus  
36 breeds. Alleles 222K and 146S had the highest frequency in the two indigenous and  
37 the Damascus breed, respectively (*ca.* 6.0%). The studied breeds shared similar  
38 haplotypic frequencies with most South Italian and Turkish breeds but differed  
39 significantly from North-Western European, Far East and some USA goat breeds.  
40 Results suggest there is adequate variation in the *PRNP* gene locus to support  
41 breeding programs for enhanced scrapie resistance in goats reared in Greece. Genetic

42 comparisons among goat breeds indicate that separate breeding programs should  
43 apply to the two indigenous and the imported Damascus breeds.

44

## 45 **Introduction**

46 Scrapie is an infectious, neurodegenerative and fatal disease of sheep and goats.  
47 Along with bovine spongiform encephalopathy, Creutzfeldt-Jacob disease in humans  
48 and chronic wasting disease in cervids, scrapie belongs to the group of transmissible  
49 spongiform encephalopathies (TSEs), also known as prion diseases. Scrapie is the  
50 result of the accumulation of PrP<sup>Sc</sup> in the central nervous system, which is an  
51 abnormal  $\beta$ -sheet rich isoform of the normal  $\alpha$ -helix rich PrP<sup>C</sup> protein [1].

52 In sheep, susceptibility to classical scrapie is modulated by the *PRNP* gene which  
53 encodes PrP<sup>C</sup> protein [2]. Polymorphisms at codons 136 (A/V), 154 (Q/R) and 171  
54 (Q/R/H) of this gene have been associated with susceptibility (ARQ and VRQ  
55 haplotypes) or resistance (ARR haplotypes) to classical scrapie [3,4,5,6]. Based on  
56 these findings a five-group risk classification system has been developed [7] and  
57 applied by different countries in selective breeding programs to control and eradicate  
58 the disease [8].

59 In goats, previous research has revealed more than 30 polymorphic codons in the  
60 *PRNP* gene. Amongst these, polymorphisms at codons 142 (I/M), 143 (H/R), 146  
61 (N/S/D), 154 (H/R), 211 (R/Q) and 222 (Q/K) have been associated with resistance or  
62 susceptibility to clinical manifestation of the disease in field studies [9-16].  
63 Experimental studies have shown that alleles 222K, 146S and 211Q confer the  
64 strongest degree of resistance after oral and/or intracerebral challenge and, therefore,  
65 are considered the most suitable candidates for selective breeding programs in goats  
66 [17-22]. This is also supported by the European Food Security Authority (EFSA) in

67 their scientific opinion on the genetic resistance to TSEs in goats [23]. However, there  
68 are currently no formal regulations to underpin selective breeding programs in goats,  
69 equivalent to those in sheep. The notion is that success of such programs is subject to  
70 in-depth knowledge of the frequency and distribution of *PRNP* polymorphisms  
71 conferring resistance to classical scrapie [14,23]. Different estimates of allelic  
72 frequency have been published for breeds in China [24], Japan [25], USA [26], Italy  
73 [27-29], France [13], UK [30], Cyprus [31] and Turkey [32]. Given the observed  
74 differences between countries and breeds, EFSA concluded that any selective  
75 breeding programs should be developed and managed at national level, and in each  
76 case, the frequency of the resistance-associated alleles and haplotypes should be  
77 assessed for each breed of interest [23].

78 Dairy goats constitute a significant sector of livestock farming in Greece. The Greek  
79 goat population is approximately 3.8 million [33], which makes it the largest in  
80 Europe and one of the largest worldwide. This population is composed mainly of the  
81 Eghoria and Skopelos indigenous breeds, as well as the imported Damascus breed  
82 along with various types of crossbreeds. Scrapie was first reported in Greece in 1986  
83 and has since been considered a major health problem in both sheep flocks and goat  
84 herds. Studies on the frequencies of classical scrapie resistance-associated alleles in  
85 goats of Greece are limited [15,34,35]. These studies either focused on a small  
86 number of goats from scrapie-affected herds [34] or animals of unspecified breeds  
87 [15,35]. A systematic large-scale population study of the *PRNP* gene polymorphisms  
88 in dairy goat breeds reared in Greece is missing.

89 The objective of the present study was to assess the genetic profile of codons 146, 211  
90 and 222 in the *PRNP* gene locus in the three key dairy goat breeds in Greece, namely

91 Eghoria, Skopelos and Damascus in order to determine the feasibility of breeding  
92 programs aiming at enhancing resistance to classical scrapie.

93

## 94 **Materials and methods**

### 95 **Ethics statement**

96 This study followed the European Directive 86/609/EEC and its national  
97 implementation in Greece Presidential Decree No 160/1991 (Governmental Gazette  
98 No A' 64). Blood sampling of dairy goats was performed in commercial farms within  
99 the EU SOLID project (FP7-KBBE-266367). This research was approved by the  
100 Research Committee of The Aristotle University of Thessaloniki (26362/03.05.2011).

101

### 102 **Animals**

103 A total of 766 dairy goats of two indigenous Greek breeds (Eghoria and Skopelos, n =  
104 264 and n = 287 goats, respectively) and one foreign breed (Damascus, n = 215 goats)  
105 were used. Animals of 1-4 years of age were randomly selected from seven farms  
106 (two with Eghoria, two with Skopelos and three with Damascus goats). Goat herds  
107 were selected as representatives of the prevailing farming systems in the country  
108 based on the *a posteriori* typology scheme described by Gelasakis *et al.* [36]. These  
109 were low-input pastoral farming systems characterized by grazing throughout the  
110 year, random mating and minor differences in management practices between herds  
111 [36]. Based on the latter, the selected number of seven herds was considered sufficient  
112 for the study. Relations of the selected animals were unknown since in all cases,  
113 random mating was performed. However, considering that the number of bucks and  
114 their replacement rate were high, over-representation of lineages is unlikely.

115

## 116 **Genotyping**

117 Blood samples were collected from the jugular vein in EDTA vacutainers and  
118 genomic DNA was extracted using GeneJET Whole Blood Genomic DNA  
119 Purification Mini Kit (Thermo Scientific, Waltham, Massachusetts, USA) according  
120 to manufacturer's instructions. Polymorphisms at codons 146 (N/S/D), 211 (R/Q) and  
121 222 (Q/K) were detected using a novel genotyping approach. Contrary to previous  
122 studies, which used PCR amplification of the open reading frame of caprine *PRNP*  
123 gene and sequencing of the PCR products, we focused on the most important codons  
124 (146, 211 and 222) and developed four separate Real-Time PCR reactions with four  
125 different Custom TaqMan<sup>®</sup> Single Nucleotide Polymorphism (SNP) Genotyping  
126 Assays (Applied Biosystems, Foster City, California, USA). Real-Time PCR is fast,  
127 sensitive and suitable for SNP detection. This method has been also used in sheep for  
128 the detection of polymorphisms at codons 136, 154 and 171 in the *PRNP* gene [37].  
129 Each SNP Genotyping Assay consisted of a mix of sequence-specific forward and  
130 reverse primers to amplify the polymorphic sequence of interest and two TaqMan<sup>®</sup>  
131 MGB (minor groove binder) probes, FAM (6-carboxyfluorescein) and VIC (2'-chloro-  
132 7-phenyl-1,4-dichloro-6-carboxy-fluorescein) dye-labeled to detect the amplified  
133 product (Table 1). Primers and probes were designed by Applied Biosystems using  
134 designated software and then individually tested by mass spectroscopy to verify the  
135 accuracy of the resulting synthesized oligonucleotide. Validation of the genotyping  
136 method was performed using sequenced goat genomic DNA samples with all possible  
137 genotypes at codons 146, 211 and 222.

138

139 **Table 1. Primer and probe sequences used in four Custom TaqMan® SNP**  
 140 **Genotyping Assays for the detection of polymorphisms 146 (N/S), 146 (N/D), 211**  
 141 **(R/Q) and 222 (Q/K).**

SNP	Primer	
	Forward	Reverse
146(N/S)	GCCATGAGCAGGCCTCTTATA	GGGTAACGGTACATGTTTTTCACGAT
146(N/D)	GCCATGAGCAGGCCTCTT	GGGTAACGGTACATGTTTTTCACGAT
211(R/Q)	GAACTTCACCGAAACTGACATCAAG	ACTGGGTGATGCACATTTGCT
222(Q/K)	TGGTGGAGCAAATGTGCATCA	GGGAAGAAAAGAGGATCACACTTG
	Probe <sup>†</sup>	
	FAM	VIC
146(N/S)	TTTTGGCAGTGA CTATG	CATTTTGGCAATGA CTATG
146(N/D)	CATTTTGGCAATGA CT	ATACATTTTGGCGATGA CT
211(R/Q)	AATGGAGCAAGTGGTG	ATAATGGAGCGAGTGGTG
222(Q/K)	CTGGGATTCTCTTGTACTG	TGGGATTCTCTTGGTACTG

142 <sup>†</sup>In all cases except for 146(N/D), FAM dye-labeled probes were used for the  
 143 detection of the mutated sequences and VIC dye-labeled probes for the detection of  
 144 the wildtype sequences.

145

146 PCR reactions were performed in 12.5 µl mixtures containing 6.25 µl of KAPA  
 147 PROBE FAST qPCR Master Mix (2X) Universal (Kapa Biosystems, Wilmington,  
 148 Massachusetts, USA), 0.3125 µl of each SNP Genotyping Assay Mix (40X) and 1 µl  
 149 (*ca.* 50 ng) of genomic DNA. Thermal cycling included: a) an initial denaturation step  
 150 at 95°C for 3 min and b) 45 cycles of denaturation at 95°C for 3 s and primer  
 151 annealing/extension at 62°C for 30 s. All Real-Time PCR analyses (n = 3,064) were



152 performed using Applied Biosystems StepOnePlus™ Real-Time PCR System and  
153 genotypes were determined through amplification plots with QPCR Applied  
154 Biosystems Step One Software v2.3 ([dx.doi.org/10.17504/protocols.io.qeidtce](https://doi.org/10.17504/protocols.io.qeidtce)).

155

## 156 **Genetic Analyses**

157 Genotypic, allelic and haplotypic frequencies regarding codons 146, 211 and 222  
158 were calculated within and across breed, based on counting of the respective  
159 genotypes of individual animals. Hardy-Weinberg equilibrium state was examined at  
160 each codon and breed using a chi-square test:

$$161 \chi^2 = \sum[(O - E)^2/E]$$

162 Where

163 O is the observed number of each genotype;

164 E is the expected number of each genotype assuming Hardy-Weinberg equilibrium;

165 Summation is over all possible genotypes.

166 Pairwise comparison of genotypic distribution between breeds at each codon was  
167 carried out using a chi-square test. In each case, number of observed genotypes in one  
168 breed was compared to expected number based on the allelic frequency in the other  
169 breed. Moreover, genetic distances among the three studied breeds were estimated on  
170 the basis of haplotypic frequencies using fixation index  $F_{ST}$  and Arlequin v3.1  
171 software [38]. To add substance to these comparisons, genetic distances were also  
172 calculated between the three studied breeds and 30 other goat breeds from different  
173 countries, for which haplotypic frequencies were available in the existing literature  
174 (S1 Table). In each case, the significance of difference from the null value was tested  
175 with 1,000 permutations. The matrix of genetic distances between breed pairs was  
176 used to create a Neighbor-Joining tree deploying the PHYLIP package v3.695 [39].

177 In all the above analyses, level of statistical significance was set at  $P=0.01$ .

178

## 179 **Results**

### 180 **Frequencies**

181 Genotypic, allelic and haplotypic frequencies in the *PRNP* gene locus are presented in  
182 Tables 2 and 3 and S2 Table, respectively. Across all breeds, four out of six possible  
183 genotypes at codon 146 (NN, NS, ND and SS), and two out of three at each of codons  
184 211 (RR and RQ) and 222 (QQ and QK) were detected. Genotypes carrying  
185 resistance-associated alleles, namely 222QK, 146NS, 211RQ, 146ND and 146SS  
186 were found at a frequency of 8.74%, 3.39%, 2.87%, 0.13% and 0.13%, respectively  
187 (Table 2). All resistance-associated alleles were observed in the three breeds  
188 collectively; the most frequent was 222K (4.37%), followed by 146S (1.83%), 211Q  
189 (1.44%) and 146D (0.06%, Table 3). Within breed, allele 222K had the highest  
190 frequency in the two indigenous Greek breeds (5.87% and 5.92% in Eghoria and  
191 Skopelos, respectively), whereas 146S was the most frequent in Damascus (6.05%,  
192 Table 3). Frequencies of the resistance-associated polymorphisms dictated the  
193 estimation of haplotypic frequencies in the *PRNP* gene locus; haplotypes carrying  
194 multiple resistance-associated polymorphisms were not detected (S2 Table).

195 All three codons were found to be in Hardy-Weinberg equilibrium ( $P>0.01$ ) in all  
196 breeds, suggesting absence of direct or indirect genetic selection pressure.

197

198 **Table 2. Genotypic frequencies (%) and corresponding number of animals (in**  
199 **parenthesis) at codons 146, 211 and 222 of the *PRNP* gene locus.**

		<b>Breed</b>
--	--	--------------

Codon	Genotype	Eghoria <sup>a</sup>	Skopelos <sup>a</sup>	Damascus <sup>b</sup>	Total
146	NN	99.62 (263)	99.65 (286)	87.91 (189)	96.35 (738)
	NS	0.00 (0)	0.35 (1)	11.63 (25)	3.39 (26)
	SS	0.00 (0)	0.00 (0)	0.46 (1)	0.13 (1)
	ND	0.38 (1)	0.00 (0)	0.00 (0)	0.13 (1)
	DD	0.00 (0)	0.00 (0)	0.00 (0)	0.00 (0)
	DS	0.00 (0)	0.00 (0)	0.00 (0)	0.00 (0)
211	RR	96.97 (256)	100.00 (287)	93.49 (201)	97.13 (744)
	RQ	3.03 (8)	0.00 (0)	6.51 (14)	2.87 (22)
	QQ	0.00 (0)	0.00 (0)	0.00 (0)	0.00 (0)
222	QQ	88.26 (233)	88.15 (253)	99.07 (213)	91.26 (699)
	QK	11.74 (31)	11.85 (34)	0.93 (2)	8.74 (67)
	KK	0.00 (0)	0.00 (0)	0.00 (0)	0.00 (0)

200 <sup>a,b</sup> Comparison of genotypic distributions between breeds; different superscripts  
201 denote significant differences (P<0.01); distributions were compared within codon  
202 and outcome was the same in all three codons.

203

204 **Table 3. Allelic frequencies (%) at codons 146, 211 and 222 of the *PRNP* gene**

205 **locus.**

Codon	Allele	Breed			
		Eghoria	Skopelos	Damascus	Total
146	N	99.81	99.83	93.95	98.11
	S*	0.00	0.17	6.05	1.83
	D*	0.19	0.00	0.00	0.06

211	R	98.48	100	96.74	98.56
	Q*	1.52	0.00	3.26	1.44
222	Q	94.13	94.08	99.53	95.63
	K*	5.87	5.92	0.47	4.37

206 \*Indicates alleles which are considered to confer resistance to scrapie.

207

## 208 **Genetic comparison among the three goat breeds in Greece**

209 Genotypic distribution in Damascus breed differed significantly ( $P < 0.01$ ) from that in  
 210 Eghoria and Skopelos breeds for each of the three codons (Table 2). No significant  
 211 differences ( $P > 0.01$ ) were observed between Eghoria and Skopelos breeds in any  
 212 codon.

213 Genetic distance analysis (Table 4) showed that Eghoria and Skopelos breeds shared  
 214 similar haplotypic frequencies implying genetic relatedness ( $F_{ST} = 0.000$ ,  $P > 0.01$ ),  
 215 whereas they differed significantly from Damascus breed ( $F_{ST} = 0.020$ ,  $P < 0.001$  and  
 216  $F_{ST} = 0.028$ ,  $P < 0.001$ , respectively), thereby corroborating the genotypic distribution  
 217 results.

218

219 **Table 4. Genetic ( $F_{ST}$ ) distances (above diagonal) and corresponding P-values**  
 220 **(below diagonal) among Eghoria, Skopelos and Damascus breeds.**

Breeds	Eghoria	Skopelos	Damascus
Eghoria		0.000	0.020
Skopelos	0.31543		0.028
Damascus	<0.00001*	<0.00001*	

221 \*Indicates significant P-values ( $P < 0.01$ ); five decimal points is the maximum number  
 222 provided by the Arlequin software.

223

224 **Genetic comparison between the three goat breeds in Greece**  
225 **and breeds in other countries**

226 Estimated genetic distances between goat breeds reared in Greece and in other  
227 countries are presented in Table 5. Small  $F_{ST}$  estimates and high (non-significant) P-  
228 values denote high levels of genetic relatedness between respective breeds. The three  
229 studied breeds shared similar haplotypic frequencies ( $P>0.01$ ) with most breeds reared  
230 in neighboring Mediterranean regions (Southern Italy, Turkey and Cyprus). Notable  
231 exceptions were the observed distances ( $P<0.01$ ) from four Southern Italian breeds  
232 (Garganica, Girgentana, Derivata di Siria and Pantellaria), and the Damascus breed of  
233 Cyprus and Turkey. Significant genetic distances ( $P<0.01$ ) were also observed from  
234 goat breeds reared in North-Western Europe (Northern Italy, France and UK) and Far  
235 East (Japan and China). Regarding breeds from the USA, Greek goats were  
236 genetically distant ( $P<0.01$ ) from LaMancha and Nubian, while a significant  
237 difference was also observed between Skopelos and Alpine and Saanen breeds.

238

239 **Table 5. Genetic ( $F_{ST}$ ) distances and corresponding P-values between goat breeds**  
240 **reared in Greece and other countries.**

Other breeds (Region/Country <sup>†</sup> )	Breeds in Greece					
	Eghoria		Skopelos		Damascus	
	$F_{ST}$	P-value	$F_{ST}$	P-value	$F_{ST}$	P-value
Garganica (Southern IT)	0.058	<0.00001*	0.076	<0.00001*	0.082	<0.00001*
Maltese (Southern IT)	0.009	0.19824	0.018	0.09961	0.037	0.01562
Ionica (Southern IT)	0.000	0.90723	0.000	0.78711	0.018	0.07812

Red Mediterranean (Southern IT)	0.000	0.77441	0.000	0.99902	0.016	0.06445
Girgentana (Southern IT)	0.064	<0.00001*	0.079	<0.00001*	0.087	<0.00001*
Derivata Di Siria (Southern IT)	0.034	<0.00001*	0.044	<0.00001*	0.060	<0.00001*
Pantellaria (Southern IT)	0.169	<0.00001*	0.216	<0.00001*	0.119	<0.00001*
Camosciata Delle Alpi (Northern IT)	0.059	<0.00001*	0.087	<0.00001*	0.040	<0.00001*
Saanen (Northern IT)	0.030	0.00098*	0.051	<0.00001*	0.021	0.00879*
Roccoverano (Northern IT)	0.063	<0.00001*	0.093	<0.00001*	0.046	<0.00001*
Valdostana (Northern IT)	0.027	<0.00001*	0.047	<0.00001*	0.017	0.00390*
Damascus (CY)	0.037	<0.00001*	0.046	<0.00001*	0.011	0.00293*
Damascus (TR)	0.263	<0.00001*	0.307	<0.00001*	0.159	<0.00001*
Akkeci (TR)	0.043	0.04980	0.074	0.01465	0.018	0.12793
Saanen (TR)	0.002	0.24805	0.013	0.06250	0.007	0.12109
Kilis (TR)	0.024	0.01171	0.031	0.01170	0.023	0.01855
Alpine (FR)	0.018	<0.00001*	0.032	<0.00001*	0.025	<0.00001*
Saanen (FR)	0.099	<0.00001*	0.127	<0.00001*	0.078	<0.00001*
Dairy breeds (UK)	0.027	<0.00001*	0.025	<0.00001*	0.042	<0.00001*
Boer (CN)	0.612	<0.00001*	0.656	<0.00001*	0.518	<0.00001*
Saanen (JP)	0.053	<0.00001*	0.077	<0.00001*	0.027	<0.00001*
Alpine (USA)	0.030	0.01465	0.052	0.00488*	0.010	0.12109
Oberhasli (USA)	0.012	0.09570	0.024	0.05371	0.007	0.17480
Toggenburg (USA)	0.000	0.86426	0.000	0.35742	0.010	0.08105
LaMancha (USA)	0.176	<0.00001*	0.212	<0.00001*	0.093	<0.00001*
Nubian (USA)	0.317	<0.00001*	0.360	<0.00001*	0.214	<0.00001*
Saanen (USA)	0.013	0.01367	0.019	0.00391*	0.007	0.07227

241 †IT= Italy, CY= Cyprus, TR= Turkey, FR= France, UK= United Kingdom, CN=  
242 China, JP= Japan, USA= United States of America

243 \*Indicates significant P-values ( $P < 0.01$ ); five decimal points is the maximum number  
244 provided by the Arlequin software.

245

246 Results in Table 5 are broadly consistent with the Neighbor-Joining tree presented in  
247 Fig 1. The latter revealed two major geographical clusters; one including the three  
248 studied breeds and most breeds from other Mediterranean countries, and another  
249 including most of the North-Western European and the Far East breeds. Goat breeds  
250 from the USA were distributed across both clusters.

251

252 **Fig 1. Unrooted Neighbor-Joining tree based on  $F_{ST}$  genetic distances.**

253 Colors indicate two major geographical clusters: Blue cluster includes: E\_GR,  
254 Eghoria (Greece); S\_GR, Skopelos (Greece); I\_S\_IT, Ionica (Southern Italy);  
255 RM\_S\_IT, Red Mediterranean (Southern Italy); D\_GR, Damascus (Greece); D\_CY,  
256 Damascus (Cyprus); K\_TR, Kilis (Turkey); O\_USA, Oberhasli (United States of  
257 America); S\_USA, Saanen (United States of America); DB\_UK, Dairy breeds (United  
258 Kingdom); T\_USA, Toggenburg (United States of America); A\_FR, Alpine (France);  
259 M\_S\_IT, Maltese (Southern Italy); DDS\_S\_IT, Derivata Di Siria (Southern Italy);  
260 G\_S\_IT, Garganica (Southern Italy); GI\_S\_IT, Girgentana (Southern Italy). Red  
261 cluster includes: S\_TR, Saanen (Turkey); V\_N\_IT, Valdostana (Northern Italy);  
262 A\_USA, Alpine (United States of America); S\_IT\_N, Saanen (Northern Italy); A\_TR,  
263 Akkeci (Turkey); S\_JP, Saanen (Japan); CDA\_N\_IT, Camosciata Delle Alpi  
264 (Northern Italy); R\_N\_IT, Roccaverano (Northern Italy); P\_S\_IT, Pantellaria  
265 (Southern Italy); S\_FR, Saanen (France); LM\_USA, La Mancha (United States of

266 America), D\_TR, Damascus (Turkey); N\_USA, Nubian (United States of America);  
267 B\_CN, Boer (China).

268

## 269 **Discussion**

270 In the present study, the genetic profile of three codons (146, 211 and 222) in the  
271 *PRNP* gene locus was characterized for the first time in the three most common dairy  
272 goat breeds in Greece (Eghoria, Skopelos and Damascus). Our results demonstrated  
273 the presence of polymorphism in this locus which renders it amenable to change via  
274 genetic selection. Moreover, differences in *PRNP* haplotypic frequencies were  
275 revealed among the studied goat breeds but also between them and breeds reared in  
276 other countries, suggesting the importance of breeding programs being custom-  
277 designed according to the genetic profile of specific populations. In this regard, our  
278 results indicate that separate breeding programs should apply to the two indigenous  
279 and the imported Damascus breeds aiming at enhancing scrapie resistance.

280 Although resistance-associated allele frequencies have been previously reported in  
281 Greek goats, this is the first large-scale population study involving animals from  
282 specific breeds, without clinical signs of scrapie and from farms representative of the  
283 major farming systems as described in relevant typologies [36]. Regarding the  
284 Eghoria and Skopelos breeds, a previous study reported allele frequencies for 222K  
285 (*ca.* 24.0% in Skopelos) and 211Q (*ca.* 7.0% in both breeds) using goats from one  
286 highly infected herd [34]. However, these results are not directly comparable with  
287 ours as they were based on a different study design and a smaller sample size. Kanata  
288 *et al.* [35] reported frequencies in Greek goats higher than our study (3.0%, 0.5%,  
289 6.0% and 5.6% for 146S, 146D, 211Q and 222K alleles, respectively) but for fewer



290 animals (436 individuals) of unspecified breeds. Contrary to the latter study, we used  
291 purebred goats of the three most common dairy goat breeds reared in Greece.

292 The present study confirms that the dairy goat population in Greece, collectively  
293 represented here by the three studied breeds, is one of the very few worldwide where  
294 all scrapie resistance-associated alleles in the three codons (146, 211 and 222) have  
295 been detected. Previous research in Turkey has also reported presence of all these  
296 alleles in the goat population; however, much lower frequencies have been reported  
297 for alleles 222K and 211Q [32]. Polymorphic variation in the *PRNP* gene locus  
298 reported in our study enables the design and implementation of breeding programs  
299 towards enhanced scrapie resistance in goats reared in Greece.

300 The genetic profiles of scrapie codons 146, 211 and 222, revealed certain differences  
301 between the studied breeds. The most common resistance-associated allele was 222K  
302 in Eghoria and Skopelos (indigenous Greek breeds), whereas 146S in Damascus (*ca.*  
303 6.0% in all cases). Such observations were further supported by comparison of  
304 genotypic distributions and genetic distance analyses, which showed significant  
305 differences between the two indigenous Greek breeds and Damascus. The latter is an  
306 imported breed of Middle East origin. Previous studies have suggested that  
307 geographical origin may explain a large part of genetic variability among goat breeds  
308 [39]. In a microsatellite analysis of European and Middle Eastern goats, Greek breeds  
309 were placed in the Central-Mediterranean cluster within which genetic differences  
310 among breeds were found to be relatively low [40]. In the same study, breeds of  
311 Middle Eastern origin were placed in a separate, East-Mediterranean genetic cluster  
312 [40].

313 To further investigate differences in the *PRNP* gene locus among different goat  
314 populations, we estimated the genetic distance between the studied breeds and 30

315 other breeds reared in various countries. To the best of our knowledge this is the first  
316 time that a large-scale comparison of the *PRNP* gene profile is attempted using  
317 collectively all the available data regarding polymorphism frequencies at codons 146,  
318 211 and 222 in different dairy goat populations worldwide. In this regard, we derived  
319 novel and interesting results about the level of relatedness of these populations. Of  
320 course a certain level of caution needs to be exercised when interpreting these results  
321 as the relatively small number of polymorphisms and occasionally limited  
322 representation in the studied breeds might have influenced the accuracy of the genetic  
323 distance estimates. Furthermore, some of the previous studies on breeds reared in  
324 Italy, Turkey and USA had been based on just a few animals, which might have  
325 affected the representativeness of the calculated haplotypic frequencies.

326 According to previous research, allelic frequencies vary across countries and breeds  
327 [13, 24-32]. Results from our comparisons suggest that geographical origin of the  
328 breeds combined with population structure, breeding schemes and scrapie incidence  
329 seem to be the key determinants of the *PRNP* gene profile. Thus, the studied breeds  
330 shared similar haplotypic frequencies with many goat breeds reared in neighboring  
331 Mediterranean countries, such as certain breeds in Southern Italy (Maltese, Ionica and  
332 Red Mediterranean) [27] and Turkey (Saanen, Akkeci and Kilis) [32]. However,  
333 significant genetic distances were also detected in respect to some other breeds reared  
334 in neighboring countries. For example, much higher allelic frequencies of 222K were  
335 observed in Girgentana (18.7%) and Derivata Di Siria (15.0%), but the allele was  
336 absent in Pantellaria [28,29]. Girgentana is an endangered breed of Sicily with a small  
337 population size in which significant inbreeding and genetic drift have been recorded  
338 [28] potentially leading to this high allelic frequency. Derivata Di Siria goats are  
339 reared in a high-scrapie endemic area of Sicily, whereas Pantellaria goats are found in

340 an area where scrapie has never been detected [29], quite possibly explaining the  
341 corresponding allelic frequencies in each case. Moreover, Pantellaria goats had allele  
342 211Q in high frequency (23.0%) [29]. This could be interpreted as a consequence of  
343 the extensive crossbreeding with Saanen and Alpine goats of Northern Italy [29] in  
344 which allele 211Q was also found to be very common [27].

345 Furthermore, significant genetic distances were also detected between Greek  
346 Damascus and goats of the same breed in Turkey and Cyprus. In both Greek and  
347 Cypriot Damascus goats, 146S was the most common allele (*ca.* 6.0% to 7.0%) [31].  
348 However, allele 146D, which was not detected in the present study, was also reported  
349 in Cyprus. In Damascus goats of Turkey, allele 146D was not detected, but a much  
350 higher frequency of 146S (*ca.* 28.0%) was reported [32]. Moreover, alleles 211Q and  
351 222K were detected for the first time in Damascus goats of Greece. These results  
352 suggest a diverse evolution pattern of the Damascus breed in the three neighboring  
353 countries, where different breeding schemes, possibly involving crossbreeding with  
354 indigenous breeds, have taken place. Within-breed differences in haplotypic  
355 frequencies were also detected between Saanen goats reared in different countries  
356 (France, Italy, Turkey and USA) further enhancing the notion of diverse selective  
357 breeding in different localities.

358 All goat breeds in the present study were significantly distant from North-Western  
359 European and Far East breeds in which the most common alleles were 211Q  
360 (Northern Italian, French Alpine, French Saanen and Japanese Saanen breeds) or  
361 146S (Chinese Boer), whereas 222K was either not detected or found in very low  
362 frequencies [13,24,25,27]. These findings could be explained by the different  
363 geographical origin of the breeds and are in agreement with the phylogeographical  
364 structure described by Canon *et al.* [40], which placed Northern Italian and French

365 breeds in a different genetic cluster than the Greek ones. Our findings are also in  
366 agreement with previous studies of Italian goat breeds [27], where Northern Italian  
367 breeds were clustered separately from Southern Italian; the latter shared similar allelic  
368 frequencies with the Greek goats.

369 The three breeds of the present study had similar haplotypic frequencies with some  
370 breeds from USA. Such results might have been unexpected given that the latter were  
371 imported into USA from North-Western Europe. However, the sample size in these  
372 breeds on which haplotypic frequencies had been based was relatively small [26],  
373 which might have compromised the significance of the comparisons. Furthermore, no  
374 genetic scrapie studies of native USA goat breeds were found in the literature. Based  
375 on the phylogenetic analysis of goat mitochondrial sequences reported by Amills *et*  
376 *al.* [41], significant differences between native American and European populations  
377 could be expected.

378 All genetic comparisons among goat breeds discussed above support the scientific  
379 opinion of EFSA, which suggests that selective breeding programs towards scrapie  
380 resistance should be developed and managed independently within each country and  
381 according to established frequencies of resistance-associated alleles for each breed  
382 [23]. Therefore, our results suggest that separate breeding programs would be  
383 advisable for the Eghoria and Skopelos breeds on the one hand, and Damascus on the  
384 other. In all cases, however, caution should be exercised as uni-dimensional selection  
385 towards enhancing scrapie resistance could potentially lead to reduction of genetic  
386 variability and increased inbreeding, especially given the relatively low population  
387 frequencies of resistance-associated alleles [42]. An evidence-based approach to  
388 determine the best strategy is necessary. In a previous study in sheep, the impact of  
389 various selection strategies on inbreeding was investigated by simulating different

390 frequencies of the resistant ARR haplotype and no negative effect was found when  
391 haplotypic frequency was at least 5.0% [43]. Moreover, in Chios Greek sheep, ARR  
392 haplotype was found at a frequency of 6.9% [44], which enabled the design of a  
393 selective breeding program towards enhancing scrapie resistance. This program is  
394 currently being implemented. Furthermore, Cyprus was able to start a breeding  
395 program for goat scrapie resistance focused on alleles 146S and 146D with  
396 frequencies of *ca.* 6.0% in the goat population [23]. Based on the above, a similar  
397 program for goat breeds reared in Greece is feasible. Such a program should aim at  
398 increasing the frequency of 222K allele in Eghoria and Skopelos goats, and the  
399 frequencies of 146S and, potentially, 211Q alleles in Damascus goats.

400 However, before such breeding programs are implemented, future research should  
401 address potentially adverse effects of genetic selection for enhanced scrapie resistance  
402 on other important animal traits. There might be a direct impact of *PRNP* gene in the  
403 phenotypic expression of such traits and/or genetic linkage between the *PRNP* gene  
404 and genes influencing animal performance [42]. Although relevant studies have been  
405 published for sheep [44-49], studies on goats are missing.

406 Finally, a critical step towards establishing effective breeding programs is to fully  
407 understand the allelic interactions in all codons of interest. A partial dominant effect  
408 of allele 222K over the wild-type was experimentally detected by intra-cerebrally  
409 inoculating mice with various scrapie isolates; all KK animals were resistant and QK  
410 were more resistant than QQ animals showing reduced incidence rates and/or longer  
411 incubation periods [19]. To the best of our knowledge, no relevant studies have been  
412 published for alleles in other codons. Therefore, experimental goat studies including  
413 both heterozygous and homozygous carriers of resistance-associated alleles are  
414 warranted.

415

## 416 **Conclusions**

417 The results of the present study indicate that there is adequate variation in the *PRNP*  
418 gene locus to support breeding programs for enhanced scrapie resistance in goats  
419 reared in Greece. According to the genetic profile in the three studied codons,  
420 separate breeding programs should apply to the indigenous breeds and the imported  
421 Damascus breed. Future breeding programs in other countries and breeds should first  
422 study the genetic profile of the goat population in question. However, before any  
423 measures are taken it is necessary to: i) determine possible adverse effects of selection  
424 towards scrapie resistance on other important animal traits and ii) fully understand the  
425 allelic interactions in the involved codons.

426

## 427 **Acknowledgements**

428 The authors thank Rebekka Giannakou, Sofia Termatzidou, Panagiota Kazana,  
429 Aikaterini Soufleri and Maria Karatzia for their assistance in field work and Mary  
430 Kalamaki for assistance in lab work.

431

## 432 **References**

- 433 1. Prusiner, SB. Prions. Proc Natl Acad Sci. 1998; 95:13363-13383.
- 434 2. Baylis M, Goldmann W. The genetics of scrapie in sheep and goats. Curr Mol  
435 Med. 2004; 4:385–396.
- 436 3. Goldmann W. PrP genetics in ruminant transmissible spongiform  
437 encephalopathies. Vet Res. 2008; 39:30.

- 438 4. Hunter N, Foster JD, Goldmann W, Stear MJ, Hope J, Bostock C. Natural  
439 scrapie in a closed flock of Cheviot sheep occurs only in specific PrP  
440 genotypes. *Arch Virol.* 1996; 141:809-824.
- 441 5. Hunter N, Cairns D, Foster JD, Smith G, Goldmann W, Donnelly K. Is scrapie  
442 solely a genetic disease? *Nature.* 1997; 386:137.
- 443 6. Bossers A, Schreuder BEC, Muileman IH, Belt PBGM, Smits MA. PrP  
444 genotype contributes to determining survival times of sheep with natural  
445 scrapie. *J Gen Virol.* 1996; 77:2669-2673.
- 446 7. DEFRA. National Scrapie Plan for Great Britain. Schemes Brochure  
447 [Internet]. Department for Environment, Food and Rural Affairs; 2001 [cited  
448 2018 February 10]. Available from:  
449 [http://adlib.everysite.co.uk/resources/000/054/063/NSP\\_english.pdf](http://adlib.everysite.co.uk/resources/000/054/063/NSP_english.pdf)
- 450 8. Commission Decision 2003/100/EC of 13 February 2003 on laying down  
451 minimum requirements for the establishment of breeding programmes for  
452 resistance to transmissible spongiform encephalopathies in sheep (Text with  
453 EEA relevance) (notified under document number C (2003) 498). *OJ.* 041:41-  
454 45.
- 455 9. Goldmann W, Martin T, Foster J, Hughes S, Smith G, Hughes K, et al. Novel  
456 polymorphisms in the caprine PrP Gene: a codon 142 mutation associated with  
457 scrapie incubation period. *J Gen Virol.* 1996; 77:2885-2891.
- 458 10. Billinis C, Panagiotidis CH, Psychas V, Argyroudis S, Nicolaou A, Leontides  
459 S, et al. Prion protein gene polymorphisms in natural goat scrapie. *J Gen*  
460 *Virol.* 2002; 83:713–721.

- 461 11. Acutis PL, Bossers A, Priem J, Riina MV, Peletto S, Mazza M, et al.  
462 Identification of prion protein gene polymorphisms in goats from Italian  
463 scrapie outbreaks. *J Gen Virol.* 2006; 87:1029–1033.
- 464 12. Papasavva-Stylianou P, Kleanthous M, Toumazos P, Mavrikiou P, Loucaides  
465 P. Novel polymorphisms at codons 146 and 151 in the prion protein gene of  
466 Cyprus goats, and their association with natural scrapie. *Vet J.* 2007; 173:459–  
467 462.
- 468 13. Barillet F, Mariat D, Amigues Y, Faugeras R, Caillat H, Moazami-Goudarzi  
469 K, et al. Identification of seven haplotypes of the caprine PrP gene at codons  
470 127, 142, 154, 211, 222 and 240 in French alpine and saanen breeds and their  
471 association with classical scrapie. *J Gen Virol.* 2009; 90:769–776.
- 472 14. Vaccari G, Panagiotidis CH, Acin C, Peletto S, Barillet F, Acutis P, et al.  
473 State-of-the-art review of goat TSE in the European Union, with special  
474 emphasis on *PRNP* genetics and epidemiology. *Vet Res.* 2009; 40:48.
- 475 15. Fragkiadaki EG, Vaccari G, Ekateriniadou LV, Agrimi U, Giadinis ND,  
476 Chiappini B, et al. *PRNP* genetic variability and molecular typing of natural  
477 goat scrapie isolates in a high number of infected flocks. *Vet Res.* 2011;  
478 42:104.
- 479 16. Boukouvala E, Katharopoulos E, Stewart P, Palaska V, Giadinis N, Arsenos  
480 G, et al. P.87: *PRNP* polymorphisms in Greek goats affected with natural  
481 scrapie. *Prion.* 2014; 8:65.
- 482 17. Acutis PL, Martucci F, D'Angelo A, Peletto S, Colussi S, Maurella C, et al.  
483 Resistance to classical scrapie in experimentally challenged goats carrying  
484 mutation K222 of the prion protein gene. *Vet Res.* 2012; 43:8.



- 485 18. White SN, Reynolds JO, Waldron DF, Schneider DA, O'Rourke KI. Extended  
486 scrapie incubation time in goats singly heterozygous for *PRNP* S146 or K222.  
487 *Gene*. 2012; 501:49–51.
- 488 19. Aguilar-Calvo P, Espinosa JC, Pintado B, Gutierrez-Adan A, Alamillo E,  
489 Miranda A, et al. Role of the Goat K<sub>222</sub>-PrP<sup>C</sup> Polymorphic Variant in Prion  
490 Infection Resistance. *J Virol*. 2014; 88:2670–2676.
- 491 20. Lacroux C, Perrin-Chauvineau C, Corbiere F, Aron N, Aguilar-Calvo P,  
492 Torres JM, et al. Genetic Resistance to Scrapie Infection in Experimentally  
493 Challenged Goats. *J Virol*. 2014; 88:2406–2413.
- 494 21. Mazza M, Guglielmetti C, Ingravalle F, Brusadore S, Langeveld JPM,  
495 Ekateriniadou LV, et al. Low fraction of the 222K PrP variant in the protease-  
496 resistant moiety of PrPres in heterozygous scrapie positive goats. *J Gen Virol*.  
497 2017; 98:1963-1967.
- 498 22. Fast C, Goldmann W, Berthon P, Tauscher K, Andréoletti O, Lantier I, et al.  
499 Protecting effect of PrP codons M142 and K222 in goats orally challenged  
500 with bovine spongiform encephalopathy prions. *Vet Res*. 2017; 48:52.
- 501 23. Ricci A, Allende A, Bolton D, Chemaly M, Davies R, Fernández Escámez PS,  
502 et al. Genetic resistance to transmissible spongiform encephalopathies (TSE)  
503 in goats [Scientific Opinion]. *EFSA Journal*. 2017; 15(8):4962.
- 504 24. Zhang L, Li N, Fan B, Fang M, Xu W. *PRNP* polymorphisms in Chinese  
505 ovine, caprine and bovine breeds. *Anim Genet*. 2004; 35:457–461.
- 506 25. Kurosaki Y, Ishiguro N, Horiuchi M, Shinagawa M. Polymorphisms of  
507 caprine PrP gene detected in Japan. *J Vet Med Sci*. 2005; 67:321–323.

- 508 26. White S, Herman-Hoesing L, O'Rourke K, Waldron D, Rowe J, Alverson J.  
509 Prion gene (*PRNP*) haplotype variation in United States goat breeds (*Open*  
510 *Access publication*). *Genet Select Evol.* 2008; 40:553–561.
- 511 27. Acutis PL, Colussi S, Santagada G, Laurenza C, Maniaci MG, Riina MV, et  
512 al. Genetic variability of the *PRNP* gene in goat breeds from Northern and  
513 Southern Italy. *J Appl Microbiol.* 2008; 104:1782–1789.
- 514 28. Migliore S, Agnello S, Chiappini B, Vaccari G, Mignacca SA, Di Marco Lo  
515 Presti V, et al. Biodiversity and selection for scrapie resistance in goats:  
516 Genetic polymorphism in “Girgentana” breed in Sicily, Italy. *Small Rumin*  
517 *Res.* 2015; 125:137–141.
- 518 29. Migliore S, Agnello S, D'Avola S, Goldmann W, Di Marco Lo Presti V,  
519 Vitale M. A cross-sectional study of *PRNP* gene in two native Sicilian goat  
520 populations in Italy: a relation between prion gene polymorphisms and scrapie  
521 incidence. *J Genet.* 2017; 96:319–325.
- 522 30. Goldmann W, Ryan K, Stewart P, Parnham D, Xicohtencatl R, Fernandez N,  
523 et al. Caprine prion gene polymorphisms are associated with decreased  
524 incidence of classical scrapie in goat herds in the United Kingdom. *Vet Res.*  
525 2011; 42:110.
- 526 31. Papasavva-Stylianou P, Windl O, Saunders G, Mavrikiou P, Toumazos P,  
527 Kakoyiannis C. PrP gene polymorphisms in Cyprus goats and their association  
528 with resistance or susceptibility to natural scrapie. *Vet J.* 2011; 187:245–250.
- 529 32. Meydan H, Pehlivan E, Özkan MM, Yildiz MA, Goldmann W. Prion protein  
530 gene polymorphisms in Turkish native goat breeds. *J Genet.* 2017; 96:299–  
531 305.

- 532 33. Eurostat  
533 ([http://appsso.eurostat.ec.europa.eu/nui/show.do?dataset=apro\\_mt\\_lsgoat&lan](http://appsso.eurostat.ec.europa.eu/nui/show.do?dataset=apro_mt_lsgoat&lang=en)  
534 [g=en](http://appsso.eurostat.ec.europa.eu/nui/show.do?dataset=apro_mt_lsgoat&lang=en)). Accessed 2018 February 18.
- 535 34. Bouzalas IG, Dovas CI, Banos G, Papanastasopoulou M, Kritas S, Oevermann  
536 A, et al. Caprine *PRNP* polymorphisms at codons 171, 211, 222 and 240 in a  
537 Greek herd and their association with classical scrapie. *J Gen Virol.* 2010;  
538 91:1629–1634.
- 539 35. Kanata E, Humphreys-Panagiotidis C, Giadinis ND, Papaioannou N,  
540 Arsenakis M, Sklaviadis T. Perspectives of a scrapie resistance breeding  
541 scheme targeting *Q211*, *S146* and *K222* caprine *PRNP* alleles in Greek goats.  
542 *Vet Res.* 2014; 45:43.
- 543 36. Gelasakis AI, Rose G, Giannakou R, Valergakis GE, Theodoridis A,  
544 Fortomaris P. Typology and characteristics of dairy goat production systems  
545 in Greece. *Livest Sci.* 2017; 197:22-29.
- 546 37. Boukouvala E, Gelasakis AI, Kanata E, Fragkiadaki E, Giadinis ND, Palaska  
547 V, et al. The association between 171K polymorphism and resistance against  
548 scrapie affection in Greek dairy sheep. *Small Rumin Res.* 2018; 161:51-56.
- 549 38. Excoffier L, Laval G, Schneider S. Arlequin ver. 3.0: an integrated software  
550 package for population genetics data analysis. *Evol Bioinform Online.* 2005;  
551 1:47-50.
- 552 39. Felsenstein J. PHYLIP (Phylogeny Inference Package) version 3.695.  
553 Distributed by the author, Department of Genome Sciences and Department of  
554 Biology, University of Washington, Seattle. 2013.

- 555 40. Cañón J, García D, García-Atance MA, Obexer-Ruff G, Lenstra JA, Ajmone-  
556 Marsan P, et al. Geographical partitioning of goat diversity in Europe and the  
557 Middle East. *Anim Genet.* 2006; 37:327-334.
- 558 41. Amills M, Ramírez O, Tomàs A, Badaoui B, Marmi J, Acosta J, et al.  
559 Mitochondrial DNA diversity and origins of South and Central American  
560 goats. *Anim Genet.* 2009; 40:315-322.
- 561 42. Dawson M, Moore RC, Bishop SC. Progress and limits of PrP gene selection  
562 policy. *Vet Res.* 2008; 39.
- 563 43. Man WY, Lewis RM, Boulton K, Villanueva B. Predicting the consequences  
564 of selecting on PrP genotypes on PrP frequencies, performance and inbreeding  
565 in commercial meat sheep populations. *Genet Sel Evol.* 2007; 39:711–729.
- 566 44. Psifidi A, Basdagianni Z, Dovas CI, Arsenos G, Sinapis E,  
567 Papanastassopoulou M, et al. Characterization of the *PRNP* gene locus in  
568 Chios dairy sheep and its association with milk production and reproduction  
569 traits. *Anim Genet.* 2011; 42:406–414.
- 570 45. Sweeney T, Hanrahan JP. The evidence of associations between prion protein  
571 genotype and production, reproduction and health traits in sheep. *Vet Res.*  
572 2006; 39:28.
- 573 46. Álvarez L, Gutiérrez-Gil B, San Primitivo F, De La Fuente LF, Arranz JJ.  
574 Influence of prion protein genotypes on milk production traits in Spanish  
575 Churra sheep. *J Dairy Sci.* 2006; 89:1784-1791.
- 576 47. Casellas J, Caja G, Bach R, Francino O, Piedrafita J. Association analyses  
577 between the prion protein locus and reproductive and lamb weight traits in  
578 Ripolessa sheep. *J Anim Sci.* 2007; 85:592-597.

579 48. Lipsky S, Brandt H, Lühken G, Erhardt G. Analysis of prion protein genotypes  
580 in relation to reproduction traits in local and cosmopolitan German sheep  
581 breeds. Anim Reprod Sci. 2008; 103:69-77.

582 49. De Vries F, Hamann H, Drögemüller C, Ganter M, Distl O. Analysis of  
583 associations between the prion protein genotypes and production traits in East  
584 Friesian milk sheep. J Dairy Sci. 2005; 88:392-398.

585

586

## 587 **Supporting information**

588 **S1 Dataset. Genotypes at codons 146, 211 and 222 of the *PRNP* gene locus of**  
589 **individual Eghoria, Skopelos and Damascus goats obtained in the study.**

590

591 **S1 Table. Goat breeds with corresponding number of animals and haplotypic**  
592 **frequencies (%) at the *PRNP* gene locus (codon order 146, 211, 222) by**  
593 **country and literature reference.**

594

595 **S2 Table. Haplotypic frequencies (%) at the *PRNP* gene locus (codon order**  
596 **146, 211, 222).**

597