

Scotland's Rural College

Stabilization of organic carbon via chemical interactions with Fe and Al oxides in gley soils

Cloy, JM; Wilson, CA; Graham, MC

Published in:
Soil Science

DOI:
[10.1097/SS.0000000000000096](https://doi.org/10.1097/SS.0000000000000096)

Print publication: 01/01/2014

Document Version
Peer reviewed version

[Link to publication](#)

Citation for published version (APA):

Cloy, JM., Wilson, CA., & Graham, MC. (2014). Stabilization of organic carbon via chemical interactions with Fe and Al oxides in gley soils. *Soil Science*, 179(12), 547 - 560. <https://doi.org/10.1097/SS.0000000000000096>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29

The stabilisation of organic carbon via chemical interactions with Fe and Al oxides in gley soils

J. M. Cloy^{a*1}, C. A. Wilson^b and M. C. Graham^a

^a *School of GeoSciences, University of Edinburgh, West Mains Road, Edinburgh, UK*

^b *Biological and Environmental Sciences, School of Natural Sciences, University of Stirling, Stirling, UK*

*Dr. Cloy is corresponding author. Tel.: +44(0)131 535 4155. E-mail address: Joanna.Cloy@sruc.ac.uk

¹Present address: *SRUC, West Mains Road, Edinburgh, UK*

1 **Abstract**

2

3 The importance of soil organic carbon (SOC) stabilisation via chemical interactions with Fe and Al
4 oxide minerals within gley soils remains unclear. Changes in the proportions of Fe/Al oxides and SOC
5 and N contents associated with Fe/Al oxides within the profiles of gley soils under contrasting
6 hydrological regimes and freely draining control soils from Harwood Forest (northeast England) were
7 investigated. Sequential selective dissolution techniques were used to measure Fe/Al oxide crystallinity
8 and explore whether crystallinity differed between gleyed and freely draining soils. Extracts were
9 analysed using FT-IR spectroscopy to investigate the chemical characteristics of organic matter (OM)
10 associated with Fe/Al oxides. Strongly crystalline Fe oxides were the dominant (~ 50-80 % of total Fe
11 oxides present) mineral phase in gley mineral soils. Contrasting gley soil hydrological regimes
12 influenced total subsoil Fe, and total and weakly crystalline Al oxide concentrations. Also, within-
13 profile changes in strongly crystalline Fe oxide concentrations were linked to differences in
14 hydrological and redox conditions. A large proportion of SOC (generally 70-90 % of total) appeared to
15 be associated with Fe/Al oxides. Correlation plots, however, indicated that SOC contents were not
16 linearly related to amounts of total Fe and Al oxides, weakly crystalline Fe and Al oxides or strongly
17 crystalline Fe oxides. The lack of linear correlations observed for these acidic soils may be due to
18 contributions from other extractable soil components and factors such as high organic loadings and
19 insufficient amounts of Fe/Al oxides for interaction with SOC in topsoils and variable surface loadings
20 of different organic inputs at different soil depths. Subsoil C/N ratios (~18-21) were higher than those
21 in extracted subsoil residues (~9-17), suggesting that minerals other than Fe/Al oxides preferentially
22 adsorbed N-rich microbially processed compounds. The OM associated with weakly and strongly
23 crystalline Fe/Al-oxides was chemically different, the latter generally having greater hydroxyl,
24 aliphatic, carboxylate and/or phenolic character and less carbohydrate character than the former. This
25 research shows that interactions between Fe/Al oxides and SOM in redox-dynamic gley soils under
26 different hydrological regimes are complex and further investigations of SOC stabilisation in these
27 systems using selective dissolution and other complementary techniques are required.

28

29 Keywords: Gley soils, soil organic carbon, iron oxides, organo-mineral interactions, oxalate, dithionite

30

1 **1. Introduction**

2 The formation of stable pools of soil organic carbon (SOC) is a key process in the global terrestrial
3 carbon cycle. Since twice as much carbon is stored in the soil than in the atmosphere (Lal et al. 1995),
4 increasing interest in carbon storage has led to active research into SOC stabilisation. One pathway for
5 the formation of stable SOC, for which the mechanisms are still not fully understood, is via SOC-
6 mineral interactions. Soil organic matter (SOM) preferentially sorbs to metal oxyhydroxides and clay
7 minerals such as smectite (Guggenberger and Kaiser 2003) and the stabilisation of SOM through
8 organo-mineral interactions involving reactive sesquioxide and clay mineral surfaces has been reported
9 in many studies (Kaiser et al. 2002; Eusterhues et al. 2005 a and b; Wiseman and Püttmann 2006; von
10 Lützow et al. 2006; Wagai and Mayer 2007; Kalbitz and Kaiser 2008). Along with plant roots, fungal
11 hyphae and microbial- and plant-derived polysaccharides, weakly crystalline Fe and Al oxides are
12 important aggregate-binding agents (Oades, 1988; D'Angelo et al. 2009) and soil aggregate stability
13 has been found to increase with increasing Fe and Al oxide content (Krull et al. 2003). Weakly
14 crystalline minerals such as ferrihydrite ($\text{Fe}_2\text{O}_3 \cdot 0.5\text{H}_2\text{O}$) and amorphous $\text{Al}(\text{OH})_3$ have a high degree
15 of hydration, extensive surface area and variable charge and are thought to form stable organic-mineral
16 bonds through anion and inner-sphere ligand-exchange reactions. Crystallisation and dehydration
17 processes transform ferrihydrite into the strongly crystalline minerals goethite ($\alpha\text{-FeOOH}$) and
18 haematite ($\alpha\text{-Fe}_2\text{O}_3$) which have a lower surface area and charge density and consequently a lower
19 affinity for SOC (Torn et al. 1997). The amount of oxides and their crystallinity has been identified as a
20 controlling factor for SOC concentrations and turnover (Torn et al. 1997). For example, OM
21 mineralization in coniferous forest soils was found to be significantly reduced in the presence of short
22 range order minerals such as ferrihydrite and Al silicates such as allophane (Rasmussen et al. 2008).

23 Fe and Al oxides are often studied together because Fe oxides such as goethite and haematite undergo
24 isomorphic substitution of Fe with Al, but in contrast with Al, the stability of Fe oxides is affected by
25 the redox status of the soil. In general, there is a depth-related transition from oxidising conditions in
26 the upper sections to reducing conditions in the underlying sections of a soil. Gley soils, however, are
27 characterised by highly localised patterns of Fe oxides depletion and enrichment related to the
28 reduction of ferric (Fe^{3+}) and mobilisation of ferrous (Fe^{2+}) ions in anaerobic regions and the
29 reoxidation and subsequent precipitation as Fe^{3+} oxides in oxidising areas of the soil. These redox
30 processes give gley soils characteristic brown mottles which are indicative of localised re-oxidation of

1 Fe. Importantly, precipitates rich in both Fe and OM can also form, especially if interchanging redox
2 conditions prevail (Kalbitz and Kaiser 2008). As a consequence of the temporal and micro-spatial
3 variability in redox conditions, the stabilisation of SOM via interaction with Fe/Al oxides and therefore
4 the C turnover in gley soils is a highly complex process. The role of Fe and Al oxides in SOC dynamics
5 and especially the nature of the interactions between SOC and Fe/Al oxides in these gley soil systems
6 is not well-defined. The specific aims of this pedological and geochemical study are to:

- 7 i) determine changes in total SOC and N contents, proportions of Fe/Al oxides and SOC and
8 N contents associated with Fe/Al oxides within the profiles of gley (Stagnosol and
9 Gleysol) soils under contrasting hydrological regimes and freely draining control (gleyic
10 Cambisol) soils.
- 11 ii) characterise Fe/Al oxide crystallinity within the soil profiles using sequential selective
12 dissolution and explore whether Fe/Al oxide crystallinity increases in gleyed soils
13 experiencing more redox oscillations than freely draining soils.
- 14 iii) investigate chemical characteristics of OM associated with Fe/Al oxides within the soils.

15

16 The following questions will be addressed:

- 17 i) Do soil properties (e.g. SOC, N, total Al and Fe contents), proportions of Fe/Al oxides
18 (total, weakly and strongly crystalline) and SOC and N contents associated with Fe/Al
19 oxides differ significantly between gley soils?
- 20 ii) Do soil C/N ratios differ significantly in Stagnosols, Gleysols and gleyic Cambisols
21 before and after removal of SOC and N associated with Fe/Al oxides?
- 22 iii) Are whole and extractable SOC contents in Stagnosols, Gleysols and gleyic Cambisols
23 linearly correlated with Fe/Al oxide phases (total, weakly and strongly crystalline)?

24

25 **2. Materials and methods**

26 **2.1. Site and sampling**

27 The study site was Harwood Forest, Northumberland, UK (BNG NY 995 944), a commercially
28 managed Sitka Spruce (*Picea sitchensis* (Bong.) Carr.) plantation. This site has been the subject of
29 previous soil carbon and greenhouse gas investigations (Zerva et al. 2005; Ball et al. 2007;

1 Mojeremane et al., 2010). Mean annual rainfall is approximately 950 mm and mean air temperature is
2 7.6 °C. The forest covers an area of gentle slopes (< 10°) at altitudes of between 200 and 400 m above
3 ordnance datum. Soils studied here are described by Wilson et al. 2013. The underlying parent material
4 is dominated by till and drift deposits derived from Paleozoic sandstones, mudstones and shales; these
5 support Histosols, Gleysols, Stagnosols, Podzols and small areas of Cambisols (Avery, 1973; FAO,
6 2006). The chosen soils were histic Gleysol, histic Stagnosol, and gleyic Cambisol (FAO, 2006).
7 Inceptisols, Aquepts, Endoaquepts, Humic Endoaquepts (Order, Suborder, Great Group, Subgroup)
8 was the dominant soil type according to the US soil classification system (Soil Survey Staff, 2010). The
9 Gleysol profiles formed in sedimentary deposits on the margins of Fallowlees Loch, a small
10 groundwater basin lake, were affected seasonally by rises and falls in the groundwater levels. By
11 contrast, the Stagnosol profiles were situated on a gentle (5-10°) slope and drainage in the upper part of
12 the profile was restricted by the clay loam subsoils derived from the local glacial drift geology. The
13 Cambisol was located on an area of coarser textured sedimentary deposits alongside a small stream and
14 although the deeper subsoil exhibited some evidence of gleying, the upper 45 cm was relatively free-
15 draining (see Table 1 and Wilson et al. 2013 for further details). Soils were taken from grassy
16 clearances within the forest in spring 2009. Five replicate soil pits at randomly selected locations,
17 placed at least 5 m apart, were opened at each of the gley soil (Stagnosol and Gleysol) sites and one
18 soil pit was opened at the control (Cambisol) site. Mineral topsoil (10-20 cm depth) and subsoil (20-30
19 cm depth) samples of known volume were collected using a monolith tin (50 cm l x 15 cm b x 7 cm d).
20 Three replicate intact soil samples were taken at each soil depth for each of the pits and bagged
21 separately. For selected sites, single (not replicate) organic topsoil samples were taken at 0-10 cm
22 (surface vegetation not included). Soils were transported to the laboratory, weighed, air-dried as intact
23 soil sections, reweighed and passed through a 2-mm sieve.

24

25 **2.2. Physical analyses**

26 Bulk density, water content and percent water-filled pore space (WFPS) were determined for soil
27 samples (Robertson et al., 1999). Soil sub-samples were dried at 105 °C over 16 hours to correct values
28 for oven-dried moisture contents and where necessary, weights and estimated volumes of large stones
29 removed during sieving were recorded to calculate corrected bulk density values. WFPS was then
30 estimated for mineral soils according to Robertson et al. (1999) using a standard particle density value

1 of 2.65 g cm⁻³. For organic topsoils it was expected that particle densities would be considerably lower
2 than 2.65 g cm⁻³ and therefore WFPS was not calculated.

3 4 **2.3. Sequential extractions**

5 Acid ammonium oxalate extracts weakly crystalline Al³⁺ and Fe³⁺ oxides and SOM bound to these
6 phases but it does not solubilise crystalline oxides (von Lützow et al. 2007). Extraction with dithionite-
7 citrate-bicarbonate (DCB) disrupts Al³⁺ and Fe³⁺ bonds between minerals and SOM by reductive
8 dissolution and concurrently releases SOM from both weakly and strongly crystalline Al³⁺ and Fe³⁺
9 oxides into solution (von Lützow et al. 2007). If the weakly crystalline oxides have first been removed,
10 DCB can, however, be used as a selective extractant for strongly crystalline oxides. In this study,
11 weakly and then strongly crystalline Fe/Al oxides, along with the respective associated OM fractions,
12 were extracted from soil sub-samples for each replicate at each depth within each soil pit, by sequential
13 selective extraction using oxalate and then DCB. To isolate oxalate-soluble Al, Fe and bound OM, 0.02
14 M ammonium oxalate (pH 3) (Schwertmann 1964; van Oorschot & Dekkers 2001) was used in
15 preference to 0.2 M ammonium oxalate to minimise oxalate contamination of extracted OM and
16 remaining soil residues. To compare the efficiency of the 0.02 M and 0.2 M ammonium oxalate
17 extraction methods, test extractions using the conventional 0.2 M ammonium oxalate method were
18 carried out for selected single soil sub-samples from 10-20 cm and 20-30 cm depths for each pit (see
19 Supplementary Digital Content Table 1 for further details). Isolation of DCB-soluble Al, Fe and bound
20 OM from the oxalate-extracted soil residues was performed using the DCB method (Mehra & Jackson
21 1960; van Oorschot & Dekkers 1999). Extracted soil residues were then rinsed twice with deionised
22 water to remove contaminants and dried (30 °C; 48 hours) prior to total C and N analysis and total
23 digestion (see Section 2.4).

24 25 **2.4. Chemical analyses**

26 Soil pH values for sub-samples of each replicate were determined in soil solutions (air-dried
27 soil:deionised water ratio 1:2) using a Spectronic Model 3505 pH meter (Spectronic, Leeds, UK). OM
28 contents (calculated by loss on ignition at 450 °C for 16 hours) and total C and N concentrations were
29 determined (Carlo-Erba NA 1500 Elemental Analyser (Carlo-Erba, Milan, Italy)) for whole soils and
30 extracted soil residues (correcting for oven-dried % moisture). Total soil C was assumed to be equal to

1 total OC as the parent material is not calcareous (Zerva et al. 2005). The amount of OC and N
2 associated with Fe/Al oxides was determined by subtracting OC and N contents in the extracted soil
3 residues from those in the whole soils. Soil texture was measured for selected soil samples using a laser
4 diffraction Coulter counter (Coulter LS230). Representative soil samples < 2000 µm were pre-treated
5 to remove organic materials using 15% H₂O₂ and 0.1% Na₆P₆O₁₈ was added as a dispersant (Wilson
6 et al. 2009).

7
8 The Al and Fe concentrations in the oxalate and DCB extract solutions, and total Al and Fe
9 concentrations in whole soils and extracted soil residues (after microwave-assisted HNO₃/HBF₄ or
10 HNO₃/HF digestion using a modified US EPA 3052 method) were determined using a Perkin Elmer
11 Optima 5300 DV Inductively Coupled Plasma-Optical Emission Spectrometer (ICP-OES), equipped
12 with a gem-cone cross-flow nebuliser and AS 93 plus autosampler. Typical internal analytical precision
13 (±1 SD) for Al and Fe determination in extract and digest solutions was ≤ ± 5%. Calculated total and
14 extractable soil Al and Fe concentrations were corrected for oven-dried % moisture. Accuracy of total
15 concentration data was confirmed by quality control analyses of three certified reference materials
16 alongside samples; silty clay loam soil (CRM No. 7003), ombrotrophic peat (NIMT/UOE/FM001) and
17 coal (NBS SRM 1635). Determined concentrations on average were 85-95 % (Fe) and 92-95 % (Al) of
18 the certified values.

19
20 **2.5. Control extractions**

21 Control extractions using 0.049 M sodium sulfate (Na₂SO₄) were performed according to Wagai &
22 Mayer (2007) to account for levels of labile SOC (Kaiser et al. 2011) and Fe and Al that are soluble in
23 water and ion-exchangeable by oxysulfur anions, such as oxidation products of dithionite (Wagai &
24 Mayer 2007). These control extractions allowed differentiation between released particulate OC and
25 released OC associated with Fe/Al oxides. Selected single Harwood forest soil sub-samples from 0-10
26 cm, 10-20 cm and 20-30 cm depths for each pit were extracted. Extractable dissolved OC
27 concentrations were determined using a Shimadzu 6000 TOC (High Temperature Catalytic Oxidation)
28 analyser and extractable Fe and Al concentrations were determined using ICP-OES.

29

1 Test extractions using a 0.3 M solution of the organic-C based complexing agent sodium citrate
2 ($\text{Na}_3\text{C}_6\text{H}_5\text{O}_7$), as used in the DCB extraction method, were also performed for the selected sub-
3 samples. Sodium citrate extracts Fe/Al oxide-associated OM as well as some OM associated with other
4 mineral components (e.g. Al silicates). Also, like sodium sulfate, sodium citrate may extract labile
5 forms of OM. The purpose of these extractions therefore was to compare levels of SOC extracted by
6 sodium citrate with those extracted by the sodium sulfate control. Dissolved OC concentrations were
7 determined in sodium citrate extracts after blank subtraction to evaluate the proportion of SOC
8 associated with Fe/Al oxides and other mineral components and any extractable labile SOC.

9

10 **2.6. Purification of extracted OM and analysis using FT-IR spectroscopy**

11 Selected extract solutions were dialysed (dialysis tubing, MW cut-off 7000 Daltons, Medicell
12 International Ltd. dialysis membrane) against deionised water and water was changed every 24 hours
13 until the pH inside the dialysis tubing was approximately 6.5. Purification of extracts was necessary to
14 remove oxalate and citrate contaminants. Noisy FT-IR spectra obtained for undialysed oxalate and
15 DCB extracts, particularly in comparison to dialysed oxalate and DCB extracts, confirmed this (data
16 not shown). Blank oxalate and DCB extract solutions were also dialysed to check that purified extracts
17 changed from cloudy to clear. The purified extracted OM samples were freeze-dried and FT-IR spectra
18 were acquired in transmission mode with a Jasco FT/IT-460 Plus FT-IR spectrometer. Approximately
19 0.5-1 mg freeze-dried OM was mixed with KBr in a ratio of ~ 1:100. Background corrected spectra
20 were obtained in the range of $3600\text{-}500\text{ cm}^{-1}$, with 2 cm^{-1} resolution and 64 scans per sample. The
21 absorption bands were assigned according to Stevenson (1994). For comparison with extracted OM,
22 FT-IR spectra for Gleysol 10-20 cm and 20-30 cm whole soils were recorded on a Bruker Vertex 70
23 FT-IR spectrometer using a Diamond Attenuated Total Reflectance accessory.

24

25 **2.7. Statistical data analyses**

26 Measurement values for replicate sub-samples from each pit (experimental unit) were averaged or, in
27 the case of Gleysol and Stagnosol 0-10 cm soils, single samples from pits were averaged, and those
28 means were statistically analysed using MINITAB version 15. The significance of differences in soil
29 measurements at each soil depth for Gleysols and Stagnosols were tested using independent t-tests
30 (summary of results presented in Table 5). Data were checked for normality before t-tests were applied.

1 Differences between soil types were deemed significant if $P \leq 0.05$. Data for the Cambisols could not
2 be included in statistical data analyses because only one soil pit (i.e. one experimental unit) was
3 sampled. Paired t-tests were also used to compare data obtained for soil samples extracted using two
4 different oxalate concentrations (Supplementary Digital Content Table 1) and whole soil C/N ratios
5 with those of extracted soil residue C/N ratios in matched samples. Linear relationships between
6 measured SOC/extractable SOC contents and amounts of different extractable Fe/Al oxides (Fe_{DCB} ,
7 Al_{DCB} , Fe_O , Al_O , $\text{Fe}_{\text{DCB-O}}$ and $(\text{Fe} + \text{Al})_\text{O}$) were assessed using simple linear regression analysis
8 (Supplementary Digital Content Figs. 1 and 2).

9

10 **3. Results**

11

12 **3.1 Physical and chemical soil properties**

13 Visual observations confirmed that the three soil types exhibited different profiles and the gley
14 (Gleysol and Stagnosol) soils consisted of humified organic rich Ao surface layers approximately 10
15 cm deep overlying mottled mineral soils with A horizons approximately 10 cm deep. Soil profile
16 descriptions and physical and chemical characteristics of Harwood forest soils are summarised in Table
17 1. In the Gleysol, which is influenced by groundwater fluctuation, the surface organic horizon grades
18 into an organo-mineral A horizon. In the Stagnosol, which is dominated by a perched water table, the
19 distinction between the organic and mineral horizons is sharper with the organic horizon giving way to
20 a gley organo-mineral A horizon. All three profiles exhibited evidence of gleyic processes in the form
21 of Fe masses and nodules, but these were found to be more frequent and more distinct in the gley soils
22 than in the Cambisol. In the Stagnosol, Fe mottling occurred at a shallower depth (ca. 15 cm) than in
23 either the Gleysol or Cambisol, reflecting the presence of restricted drainage in the upper soil horizons
24 (Wilson et al., 2013).

25

26 Soil water contents were consistent with typical values previously measured for Harwood forest soils in
27 Spring (Ball *et al.*, 2007). Water and OM contents dropped markedly on going from the 0-10 cm (Ao
28 horizon) to the 10-20 cm (A horizon) with depth for all three soils but, as expected, Gleysol and
29 Stagnosol 0-10 cm soils had much higher water (~80-100 %) and OM (~24 %) contents than the freely
30 draining Cambisol control soils (water content ~50-70 % and OM contents ~5-9 %). WFPS values

1 (~50-80 %) were similar for all three mineral soils, being highest in the 10-20 cm mineral soils. Bulk
2 densities increased with depth from ~0.5 to ~1.0 g cm⁻³ due to increasing soil compaction and mineral
3 contents. All soils were acidic, albeit the Cambisol soils were slightly less acidic (pH ~5.1-5.4) than the
4 gley soils (pH ~3.9-4.7), and pH values remained fairly constant with depth (Table 1). Soil textures
5 were similar (sandy loam) for the 0-10 cm and 10-20 cm soils of all three soil types but clay contents in
6 the poorly draining Gleysol and Stagnosol 20-30 cm soils were higher (silt loam) than those for the
7 Cambisol (loamy sand) (Table 1).

8

9 **3.2. Soil C and N**

10 Concentrations of soil OC and N in Gleysols, Stagnosols and Cambisols are shown in Table 2. Soil OC
11 and N stocks were calculated for complete soil profiles from which 0-10, 10-20 and 20-30 cm soil
12 sections were collected. Total soil OC and N stocks from 0-30 cm in the Gleysol (5 pits) (92 ± 29 and
13 4.5 ± 1.3 t ha⁻¹, respectively) and Stagnosol (2 pits from which 0-10, 10-20 and 20-30 cm sections were
14 collected) (90 ± 20 and 4.8 ± 1.5 t ha⁻¹, respectively) soils were similar and soil OC and N stocks
15 decreased with depth (Fig. 1). Calculated total OC and N stocks from 0-30 cm for the Cambisol (1 pit)
16 (51 and 3.4 t ha⁻¹, respectively) soils were lower than those calculated for the gley soils due to the much
17 lower contribution to total OC and N stocks from the 0-10 cm Cambisol organic topsoil layer (Fig. 1).
18 Soil OC and N stocks at different depths for the Gleysols and Stagnosols were not significantly
19 different ($P > 0.05$, Table 5). Overall, calculated stocks are consistent with those measured for similar
20 soils collected from Harwood forest (Zerva et al. 2005; Ball et al. 2007). In all soils, soil OC and N
21 concentrations were lowest (~1.5-1.9 % and ~0.08-0.09 %, respectively) in the 20-30 cm soils. Gleysol
22 and Stagnosol OC and N concentrations decreased strongly with depth throughout the profile, whereas
23 for the Cambisol, highest values were found in the 10-20 cm mineral topsoil layer (Table 2). Soil C/N
24 ratios were highest in the Gleysols, suggesting that they were less humified than the Stagnosols and
25 Cambisols. With increasing depth, Gleysol and Stagnosol C/N ratios remained approximately constant
26 (~20-21 and ~18-19, respectively) but Cambisol C/N ratios varied (range ~13-20).

27

28 **3.3. Soil Al and Fe**

29 Total Al and Fe concentrations (mg g⁻¹) in Gleysols, Stagnosols and Cambisols are shown in Table 3.

30 Total Al and Fe concentrations increased with depth and were highest in the Gleysol 10-20 and 20-30

1 cm mineral soils. Stagnosol and Cambisol total Fe concentrations remained fairly constant with
2 increasing depth while total Al concentrations decreased (Table 3). Total Fe concentrations in the 20-
3 30 cm Gleysols were significantly higher ($P = 0.001$) than those found in the corresponding 20-30 cm
4 Stagnosols (Tables 3 and 5), suggesting that contrasting gley soil hydrological regimes had an effect on
5 total subsoil concentrations of the redox-sensitive element, Fe.

6

7 **3.4. Soil extractable elemental concentrations**

8 Test extractions using the conventional 0.2 M oxalate method (see Section 2.3) indicated that the Al
9 and Fe extraction by the 0.02 M oxalate method averaged 48 ± 19 % and 23 ± 9 %, respectively, of
10 those extracted by the conventional method (see Supplementary Digital Content Table 1). For this
11 reason, 0.2 M oxalate extraction Fe and Al values, rather than 0.02 M oxalate extraction values, were
12 used to calculate the amounts of weakly and strongly crystalline Fe and Al oxides present in these soils
13 (Table 4 and Supplementary Digital Content Table 1).

14 For clarity and to be consistent with terminology used in cited studies, the following abbreviations
15 will be used in this paper: Fe_O and Al_O correspond to oxalate extractable Fe and Al concentrations,
16 respectively, and represent the amounts of weakly crystalline Fe/Al oxides. Since sequential extractions
17 were carried out in this study using 0.02M oxalate followed by DCB extractions, DCB extractable Fe
18 (Fe_{DCB}) and Al (Al_{DCB}) correspond to total extractable Fe and Al concentrations (i.e. the sum of
19 sequential 0.02 M oxalate and DCB extractable Fe and Al concentrations) which represent the total
20 amounts of weakly and strongly crystalline Fe/Al oxides. Fe_{DCB-O} and Al_{DCB-O} correspond to $Fe_{DCB} -$
21 Fe_O and $Al_{DCB} - Al_O$, respectively, where Fe_O and Al_O are conventional 0.2 M oxalate extractable Fe
22 and Al concentrations (i.e. total Fe/Al oxide concentrations subtracting weakly crystalline Fe/Al oxide
23 concentrations). Thus, Fe_{DCB-O} and Al_{DCB-O} are measures of strongly crystalline Fe and Al oxides.

24

25 **3.4.1. Control extractable Fe, Al and C**

26 Control (Na_2SO_4) extractable SOC and Fe concentrations were detectable in only some of the selected
27 samples, whereas control extractable Al concentrations were detectable in most of them. As a
28 proportion of total extractable concentrations of SOC, Fe (Fe_{DCB}) and Al (Al_{DCB}), control extractable
29 concentrations of SOC, Fe and Al ranged from 0.6-3 %, 0.01-0.14 % and 0.08-13 %, respectively. For
30 Al, proportions greater than 10 % were only found in a fifth of the control samples. Therefore, control

1 extractable concentrations of SOC, Fe and Al concentrations were considered to be negligible or minor.
2 Control extractable concentrations were not subtracted from corresponding oxalate and DCB-
3 extractable SOC, Fe_{DCB} and Al_{DCB} concentrations, which were assumed to represent total Fe/Al oxide
4 associated concentrations. As a proportion of total extractable SOC, sodium citrate extractable SOC
5 ranged from 7-38 %. Sodium citrate extracted up to approximately 15 times more SOC than the sodium
6 sulfate controls confirming that the majority of the SOC extracted using the DCB procedure was
7 associated with Fe/Al oxides, although contributions from other mineral components were possible.

8

9 **3.4.2. Total Fe and Al oxide contents**

10 Fe_{DCB} and Al_{DCB} concentrations were calculated as the sum of sequential 0.02 M oxalate and DCB
11 extractable Al and Fe concentrations. Table 3 summarises Al_{DCB} and Fe_{DCB} concentrations for the
12 Gleysols, Stagnosols and Cambisols. Greatest concentrations of Fe_{DCB} and Al_{DCB} occurred in the
13 Gleysol 10-20 cm and 20-30 cm mineral soils and values were approximately two to three times greater
14 than those in the corresponding organic 0-10 cm Gleysols (Table 3). By contrast, Stagnosol and
15 Cambisol Al_{DCB} and Fe_{DCB} concentrations remained fairly constant or showed more gradual increases
16 with depth. Al_{DCB} constituted around a tenth of Fe_{DCB} in all three soils, the only exception being Al_{DCB}
17 in the 0-10 cm Gleysol, which constituted a fifth of Fe_{DCB} concentrations (Table 3). Fe_{DCB}
18 concentrations in the 10-20 cm and 20-30 cm Stagnosols were not significantly different from each
19 other ($P > 0.05$) and were not significantly different ($P > 0.05$) from those in the corresponding
20 Gleysols (Table 5). Al_{DCB} concentrations in the 10-20 cm and 20-30 cm Stagnosols, however, were
21 found to be significantly less ($P = 0.021$ and 0.032 , respectively) than those in the corresponding
22 Gleysols (Table 5).

23

24 **3.4.3. Weakly crystalline Fe and Al oxide contents**

25 Table 3 summarises Al_O and Fe_O for the Gleysols, Stagnosols and Cambisols. Based on extraction data
26 for selected soils, comparison of 0.02 M and 0.2 M oxalate-extractable concentrations of Al and Fe for
27 Gleysols, Stagnosols and Cambisols (Supplementary Digital Content Table 1), 0.02 M and 0.2 M
28 oxalate extractable Al and Fe concentrations generally exhibited the same trends for all three soil types.
29 Fe_O decreased with depth and, with the exception of the Stagnosol soils, Al_O increased with depth
30 (Table 3 and Supplementary Digital Content Table 1). Fe_O concentrations in the 10-20 cm and 20-30

1 cm Stagnosol soils were not significantly different ($P > 0.05$) than those in the corresponding Gleysols.
2 Al_O concentrations in the 20-30 cm Stagnosol soils, however, were significantly lower than those in the
3 corresponding Gleysols ($P = 0.022$) (Table 5), suggesting that less saturated gley soils had lower
4 weakly crystalline Al oxide concentrations than more saturated soils at the same depths.

5

6 **3.4.4. Strongly crystalline Fe and Al oxide contents**

7 Fe_{DCB-O} contents were calculated using 0.2 M oxalate extractable Fe concentrations determined for
8 selected 10-20 and 20-30 cm soil sections (Table 4 and Supplementary Digital Content Table 1). With
9 the exception of the 10-20 cm Stagnosols, the majority of the total Fe oxides present in these soils
10 occurred as strongly crystalline phases. By contrast, the majority of the total Al oxides present occurred
11 as weakly crystalline phases. Note that Al_{DCB-O} contents are not shown in Table 4 as in some cases Al_O
12 (0.2 M oxalate method) concentrations were equal to or greater than concentrations of $Al_{DCB} \cdot Fe_{DCB-O}$
13 concentrations were significantly lower in the 10-20 cm than the 20-30 cm Stagnosols ($P = 0.024$) but
14 not in the corresponding Gleysols ($P > 0.05$, Table 5). This suggests that differences in hydrological
15 and redox conditions within the Stagnosol profile influenced strongly crystalline Fe oxide
16 concentrations with higher levels occurring in the more oxidised 20-30 cm soil layer (see Section 4.1).

17

18 **3.5. Fe/Al oxide associated C and N**

19 Calculated 0.02 M oxalate and DCB extractable SOC and N concentrations (i.e. amounts of C and N
20 associated with Fe/Al oxides) for Gleysols, Stagnosols and Cambisols are shown in Table 2. As for
21 total SOC and N concentrations, extractable OC and N concentrations decreased with depth in the
22 Gleysols and Stagnosols. In general, for Gleysols at all depths and 10-20 cm and 20-30 cm in the
23 Stagnosols and Cambisol, 70-90 % of the SOC and soil N were associated with Fe/Al oxides with
24 much lower levels of soil OC and N being found in the 0-10 cm Stagnosol (48 %) and Cambisol (22 %)
25 soils (Table 2). Calculated soil OC and N concentrations associated with Fe/Al oxides for Gleysol and
26 Stagnosol profiles were not significantly different ($P > 0.05$) (Table 5) even though significantly
27 different amounts of Fe/Al oxides within these soils were found. This could be due to contributions
28 from SOC associated with other extractable mineral components during the selective dissolution
29 procedures.

30

1 **3.6. Extracted soil residue C and N contents**

2 Concentrations of SOC and total N in extracted soil residues for Gleysols, Stagnosols and Cambisols
3 are shown in Table 2. In all three soil types, soil residue OC and total N concentrations decrease with
4 increasing depth which, with the exception of the low whole soil OC and total N concentrations
5 observed for the 0-10 cm Cambisol, is consistent with trends observed for corresponding whole soil OC
6 and total N concentrations. Soil residue OC and total N concentrations and C/N ratios determined for
7 Gleysol and Stagnosol profiles were not significantly different ($P > 0.05$). For all three 10-20 cm and
8 20-30 cm soils, measured residue C/N ratios were significantly lower (paired t-test, $P < 0.05$) than the
9 corresponding whole soil C/N ratios (Table 2). In contrast, 0-10 cm Gleysol and Stagnosol soil residue
10 C/N ratios were higher (but not significantly, paired t-test, $P > 0.05$) than those found for the whole
11 soils (Table 2).

12 13 **3.7. Chemical characteristics of SOM associated with Fe/Al oxides**

14 Selected FT-IR spectra obtained for sequential oxalate and DCB extractable OM, for 10-20 cm and 20-
15 30 cm Gleysols, Stagnosols and Cambisols, are shown in Figures 2 and 3, respectively. Extracted OM
16 exhibited the typical absorption bands of humic substances in the following regions: 3400 cm^{-1} (H-
17 bonded OH groups), 2900 cm^{-1} (aliphatic C-H stretching), $1600\text{-}1560\text{ cm}^{-1}$ (C=O of amide groups,
18 quinone and/or ketones, aromatic C=C, COO^- symmetric stretching, N-H deformation and C=N
19 stretching) 1370 cm^{-1} (OH deformation and C-O stretching of phenolic OH, COO^- antisymmetric
20 stretching, C-H deformation of CH_2 and CH_3 groups) and 1000 cm^{-1} (C-O stretching of polysaccharide
21 or polysaccharide-like substances and/or Si-O stretches of silicates if present in extracts). In general,
22 absorption bands for DCB extractable OM were broader at 3400 cm^{-1} , stronger at 2900 and 1370 cm^{-1}
23 and weaker at 1000 cm^{-1} relative to those for oxalate extractable OM. In general, similar FT-IR spectra
24 trends were observed for all three soil types and soils from different depths (Figs 2 and 3). Gleysol 10-
25 20 cm and 20-30 cm whole soil FTIR spectra (Fig. 4) confirm that the oxalate and DCB extraction
26 procedures extracted the expected important organic moieties.

27 28 **4. Discussion**

29 **4.1 Proportions of Fe/Al oxides in gley soil profiles**

1 For the soils studied here, it was found that total extractable Al and Fe oxides comprised 5-11 % and
2 40-90 % of the total soil Al and Fe, respectively (Table 3), indicating that Fe oxides are more dominant
3 than Al oxides in these soils. Similarly Bascomb (1968) found that considerable amounts of Fe oxides,
4 particularly well-crystallised Fe oxides, were present in a range of gley and well-drained brown earth
5 soil profiles from the UK. Also, Gasparatos et al. (2004) found that 78 % of the total Fe content in
6 concretions from imperfectly drained Greek alfisol soils existed in the form of Fe oxides. In the
7 Gleysols and Stagnosols studied here, highest Fe oxide contents were found in the zones of maximum
8 gleying, i.e. at depths of 10-20 and 20-30 cm (Tables 1 and 4), supporting findings from a similar study
9 of gley soils from the Yorkshire Warplands (Bloomfield 1952). Strongly crystalline oxides comprised
10 50-80 % of the total soil Fe oxides in all the Harwood forest mineral soils (Table 4), confirming the
11 dominance of strongly crystalline Fe oxides such as goethite in these temperate acid soils (Eusterhues
12 et al., 2003).

13 Interestingly, the Stagnosol 10-20 cm soils had noticeably lower (not significantly lower, $P = 0.059$,
14 see Table 5) amounts of total Fe oxides compared with corresponding Gleysols. Differences could not
15 be tested statistically but Stagnosol 10-20 cm total Fe oxide concentrations were also noticeably lower
16 than those in the Cambisol 10-20 cm soils. Only approximately one fifth of the total Fe oxides present
17 in these 10-20 cm Stagnosol soils comprised strongly crystalline Fe oxides. These trends could be
18 explained by contrasting hydrological regimes and soil redox conditions in Gleysol and Stagnosol
19 profiles. As supported by colour and mottling observations (Table 1), it is likely that the Stagnosol 10-
20 20 cm soils were more strongly reduced and the 20-30 cm soils were more strongly oxidised than the
21 corresponding Gleysol horizons. The significantly lower Al_{DCB} concentrations in the 10-20 cm and 20-
22 30 cm Stagnosol soils compared with those in the corresponding Gleysols (Section 3.4.2) is surprising
23 because Al oxides are less likely to be influenced by contrasting gley soil redox processes than Fe
24 oxides. Also, pH values for both Gleysols and Stagnosols were very similar so changes in soil pH
25 cannot explain these differences. This finding could be attributable to Al substitution within the Fe
26 oxides in the Gleysols and Gleysol sesquioxides having less dominance of Fe than Stagnosol
27 sesquioxides.

28 Thompson et al. (2006) directly demonstrated that Fe oxide crystallinity increased during soil redox
29 oscillations. Redox reactors were used to study soils subjected to a series of reduction-oxidation cycles

1 and the cumulative response of soil solids during multiple redox oscillations included a decrease in
2 concentrations of oxalate extractable Al and Fe and a transformation of short range order Fe (present in
3 the studied soils as nano-goethite) to micro-crystalline goethite and micro-crystalline haematite
4 (Thompson et al. 2006). For the Harwood Forest soils studied here, with the exception of the Stagnosol
5 10-20 cm soils, concentrations of strongly crystalline Fe oxides were indeed higher (and therefore
6 oxalate extractable Fe concentrations were lower) in the redox-dynamic gley soils than the freely
7 draining Cambisol soils (Tables 3 and 4) but this was not the case for Al oxides. Field (Table 1) and
8 micromorphological (Wilson et al., 2013) evidence support the taxonomic classifications of Gleysol
9 and Stagnosol but statistically significant differences between the Gleysol and Stagnosol with regard to
10 SOC storage and Fe oxide crystallinity were generally not found. Micromorphological examination of
11 these gley soils, teamed with Scanning Electron Microscopy-Energy Dispersive X-ray Spectrometry,
12 suggested that hydrological differences were not great enough to differentially affect the development
13 of redoximorphic conditions and therefore levels of Fe oxide crystallinity (Wilson et al., 2013).

14

15 **4.2 Calculated Fe_O/Fe_{DCB} ratios for gley soils**

16 The active Fe ratio (Fe_O/Fe_{DCB}) is a measure of the proportion of weakly crystalline Fe oxides within
17 total Fe oxides, providing a useful indication of the degree of crystallinity of Fe oxides in soils in
18 general. It is useful because it reflects the different environmental conditions (e.g. redox) under which
19 Fe oxides have been precipitated and transformed (Mansfeldt et al. 2012). Soils with a low proportion
20 of weakly crystalline Fe components typically have Fe_O/Fe_{DCB} ratios <0.2 and $(Fe + Al)_O < 10 \text{ mg g}^{-1}$
21 (Mikutta et al. 2006). As the 0.02 M oxalate extraction method was found to be less efficient than the
22 conventional 0.2 M oxalate extraction method, Fe_O/Fe_{DCB} ratios and mean $(Fe + Al)_O$ concentrations
23 were calculated for samples extracted using 0.2 M oxalate (Table 4 and Supplementary Digital Content
24 Table 1). Calculated Fe_O/Fe_{DCB} ratios for all soils at depths of 10-20 and 20-30 cm suggest that, with
25 the exception of the 10-20 cm Gleysols which had ratios <0.2 and were found to be significantly lower
26 than those in the corresponding Stagnosols ($P = 0.001$, Table 5), these soils generally contained high
27 proportions of weakly crystalline Fe components. Mean $(Fe + Al)_O$ concentrations for the soils at
28 depths of 10-20 and 20-30 cm were all below 10 mg g^{-1} (Table 4) and there were no significant
29 differences between Gleysol and Stagnosol mineral soils ($P > 0.05$, Table 5), but these low values are

1 likely to be attributable to the low concentrations of weakly crystalline Al (Al_2O_3) relative to those of Fe
2 (Fe_2O_3) found in these Harwood forest soils.

3 4 **4.3 OM accumulation and proportions of OM associated with Fe/Al oxides in gley soil profiles**

5 Several mechanisms may be responsible for the chemical interactions between OM and Fe/Al oxides in
6 gley soils. The chemical nature of the organic material (e.g. functional group content, hydrophobicity,
7 molecular size), the type of mineral (e.g. goethite, hematite) and specific surface area of mineral
8 particles are key factors (Krull et al. 2003). The Gleysols, Stagnosols and Cambisols studied here were
9 acidic ($\text{pH} \leq 5$) (Table 1) and are therefore expected to be rich in Fe and Al oxide surfaces with
10 protonated hydroxyl groups which are important for C stabilisation (Eusterhues et al. 2005a; Kiem and
11 Kögel-Knabner 2002). Interactions between these Fe/Al oxide surfaces and OM can be strong (e.g.
12 inner-sphere ligand exchange interactions between OH groups on Fe and Al oxides surfaces and OM
13 with carboxyl and phenolic OH groups) and/or weak (e.g. electrostatic interactions such as van der
14 Waals forces between protonated/positively charged functional groups on both Fe/Al oxides and OM
15 surfaces at low pH or outer-sphere interactions) (von Lützow et al. 2006). Soil OC and N stocks
16 generally decreased with depth (Fig. 1), consistent with typical trends of decreasing SOM contents with
17 depth but greater fractions of total SOM are likely stabilised by minerals in subsurface horizons due to
18 intimate associations of humified OM with minerals at these depths (Kaiser et al. 2002). A recent ^{14}C
19 study estimated total soil C mean residence times of between 181 and 305 years for organic topsoils
20 and 1072 and 1381 years for mineral subsoils from Harwood Forest (Burns 2011), suggesting that
21 mineral stabilisation mechanisms are important in Harwood Forest subsoils.

22 Oxalate and DCB-extractable SOC concentrations were broadly similar for all three of the 10-
23 20 cm (1.7–2.8 %) and 20-30 cm (1.3–1.5 %) soils and compared with the organic 0-10 cm soils,
24 higher levels of soil OC and N were associated with Fe/Al oxides (Table 2). These findings further
25 confirm the importance of organo-Fe/Al oxide associations (Wilson et al. 2013; Falsone et al. 2013)
26 and strong correlations between the presence of SOM and the presence of Fe pedofeatures in mineral
27 soils from all three soil types. The lower levels of OC and N associated with Fe/Al oxides in the
28 organic topsoils is likely to be attributable to the OM in these soils being fresher and less
29 decomposed/humified than OM in underlying mineral soils (as indicated by C/N ratio trends in Table
30 2), and therefore containing fewer reactive functional groups (e.g. COOH groups (see Section 4.4))

1 with which to interact with soil minerals (Kaiser et al. 2011). Also, Rasse et al. (2005) suggested that
2 the interaction of root-derived organic acids with mineral surfaces in subsurface soils with low OM
3 contents may be more effective than in OM-rich surface soils because subsurface soil mineral surfaces
4 are not yet saturated with OM. Therefore, another explanation for the low proportions of total OM
5 associated with Fe and Al oxides in the Harwood Forest 0-10 cm organic topsoils could be the high
6 loadings of OM and insufficient amounts of reactive Fe/Al oxide surfaces for the OM to interact
7 (Tables 1 - 3). Overall, for these Harwood Forest soils, whole SOC contents appear to vary
8 independently of Fe and Al oxide contents (Tables 2 and 3) which could be attributable to variable
9 surface loadings, depending on different organic inputs with depth, as found by Kaiser et al. (2002) for
10 two contrasting acidic forest soil profiles.

11 Enrichment and depletion patterns of OM associated with Fe/Al oxides in gley soils are controlled by
12 hydrological regimes (low mineralisation activity due to water saturation and decreasing oxygen
13 availability being an important mechanism for OM enrichment) and redox-facilitated transport
14 processes (see Section 1). Although speculative, the transport of dissolved OM and Fe/Al through
15 lateral subsurface water movement may also have occurred in the Stagnosol soils as they were situated
16 on a gentle (5-10°) slope (see Section 2.1 and Wilson et al. 2013). However, as mentioned in Section
17 4.1, the absence of significant differences between SOC storage in the Gleysols and Stagnosols studied
18 here (Section 3.2) may be due to hydrological differences not being great enough to differentially affect
19 the development of redoximorphic conditions and therefore contrasting OM enrichment and depletion
20 patterns in the whole gley soils studied here were not detected. Falsone et al. (2013) conducted a
21 micromorphological study of the SOM in the gley mineral soils considered here and found that the
22 transformation of SOM and/or the formation of organic-inorganic associations were important C
23 stabilisation mechanisms. Indeed small water retention pores were found to be associated with Fe-rich
24 amorphous organic features, indicating that soil porosity coupled with water stagnation affected Fe-
25 SOM interactions and the accumulation of transformed organic compounds in these gleyed soils
26 (Falsone et al. 2013).

27

28 **4.4 Preferential affinity of organic compounds for Fe/Al oxides**

29 Stagnosol, Gleysol and gleyic Cambisol 10-20 cm and 20-30 cm mineral soil C/N ratios were found to
30 be significantly different before and after removal of SOC and N associated with Fe/Al oxides. Oxalate

1 and DCB extracted soil residue C/N ratios (i.e. after removal of SOC and N associated with Fe/Al
2 oxides) determined for Stagnosol, Gleysol and gleyic Cambisol mineral soils, were significantly lower
3 than corresponding whole soil C/N ratios (Section 3.6). This could indicate that more labile, less
4 decomposed OM was preferentially extracted by the oxalate and DCB methods, implying that this type
5 of OM is associated with Fe/Al oxides in mineral soils but it is worth noting that some OM not
6 associated with Fe/Al oxides is likely to have also been extracted (see Section 3.4.1). The lack of
7 similar significant differences for topsoil (0-10 cm) C/N ratios (Section 3.6) is most likely due to high
8 OM loadings of fresh OM and insufficient amounts of reactive Fe/Al oxide surfaces for the OM to
9 interact (see Section 4.3). Sorption of OM to mineral surfaces is not homogeneous and fractionation of
10 organic compounds occurs during the sorption process (Kaiser and Zech 2000). Spielvogel et al. (2008)
11 reported that a specific fraction of the OC pool consisting of O/N-alkyl C and microbially-derived
12 carbohydrates is preferentially stabilised by association with Fe and Al minerals in acid subsoils.
13 Another explanation for the trends observed for whole and residue subsoil C/N ratios could be that
14 mineral phases unextractable by oxalate and DCB (e.g. minerals other than oxides and Al silicates)
15 preferentially adsorb N-rich microbially processed compounds. It has been suggested that positively
16 charged amino groups account for sorption of N-containing substances to minerals (Hedges and Oades
17 1997). Under the acidic Harwood Forest soil conditions, amino groups and Fe/Al oxides are likely to be
18 predominantly positively charged and capable of forming strong inner-sphere interactions and/or weak
19 interactions with each other, but positively charged amino groups may be particularly attracted to
20 negatively charged clay minerals such as illite (Kaiser and Zech 2000). Preferential adsorption of N-
21 rich OM on mineral surfaces and therefore low C/N ratios was observed by Wagai et al. (2009) for
22 heavy mineral-dominant SOM fractions isolated via density fractionation. It is also worth noting,
23 however, that the unextractable N-rich OM was likely to contain occluded particulate OM released
24 after chemical destruction of aggregates.

25

26 The OM associated with weakly and strongly crystalline Fe/Al oxides in the Harwood soils studied
27 here had high hydroxyl, aliphatic, carbonyl, aromatic, amide, phenolic, carboxylate and carbohydrate
28 content (Figs 2 and 3). Differences in the chemical nature of the OM associated with weakly and
29 strongly crystalline Fe/Al oxides were apparent and in general, the OM associated with strongly
30 crystalline Fe/Al oxides has greater hydroxyl, aliphatic, carboxylate and phenolic character and less

1 carbohydrate character than OM associated with weakly crystalline Fe/Al oxides. These findings
2 support those of previous studies. For example, it has been suggested that Fe oxides are strongly
3 selective for hydrophobic aromatic compounds and high MW OM fractions (e.g. Kaiser &
4 Guggenberger 2007) but hydrophilic OM, which contains more labile carbohydrates, adsorbs weakly to
5 Fe oxide surfaces (Spielvogel et al. 2008). Also, Eusterhues et al. (2005a) found that C associated with
6 Fe oxides is oxidation resistant aliphatic C.

7

8 **4.5 Correlations between SOC contents and Fe/Al oxide phases**

9 Correlation plots for the Harwood forest soils studied here indicated that SOC contents were not
10 linearly related to amounts of total Fe and Al oxides, weakly crystalline Fe and Al oxides or strongly
11 crystalline Fe oxides. Using values determined for each of the Harwood 10-20 cm and 20-30 cm soil
12 profiles, no correlations ($R^2 < 0.03$) were found between total SOC contents and Fe_O (0.2 M), Al_O (0.2
13 M), Fe_{DCB-O} (0.2 M) and $(Fe + Al)_O$ (see Supplementary Digital Content Fig. 1). Also, no strong
14 positive correlations ($R^2 < 0.04$) between either total or extractable SOC contents and amounts of
15 Fe_{DCB} and Al_{DCB} for all soils at all depths (see Supplementary Digital Content Fig. 2). This is in
16 contrast to other studies where close correlations between SOC contents and various measures of Al
17 and Fe oxides in a range of soils have been found. For example, Oades (1988) demonstrated a positive
18 relationship between SOC contents and the presence of substantial amounts of Al and Fe oxides. Kaiser
19 & Guggenberger (2000) found strong positive correlations between SOC and dithionite-extractable
20 (dithionite-only rather than DCB-extractable) Fe contents in topsoils and illuvial subsoils from 34
21 forest soils. Positive correlations between SOC content and non-crystalline Fe oxides in volcanic
22 subsoils (Torn et al. 1997) and between SOC and oxalate extractable Fe and Al ($(Fe + Al)_O$) contents in
23 French loamy soils across a range of depths (Rasse et al. 2006) have also been reported. Mikutta et al.
24 (2006) even found that mineral-associated stable OC (isolated via density fractionation) related
25 positively to the content of crystalline Fe oxides in soils with low amounts (Fe_O/Fe_{DCB} ratios < 0.2) of
26 more reactive weakly crystalline components.

27 Soil texture may be important since Kiem and Kogel-Knabner (2002) found positive linear
28 correlations between OC and weakly and strongly crystalline Fe oxide contents for sandy soils but not
29 loamy soils and it was suggested that, although stabilisation of OC via interactions of OM with Fe
30 oxide surfaces was important for both sandy and loamy soils, this mechanism may be particularly

1 dominant in sandy soils. Interestingly, Kiem and Kogel-Knabner (2002) found that strongly crystalline
2 Fe oxides made up 60 to 70 % of total Fe oxides in these loamy arable soils, consistent with findings
3 for these sandy loam, silt loam and loamy sand Harwood soils (see Section 3.5) (Table 1). Indeed
4 Eusterhues et al. (2005b) found positive correlations between organo-mineral associated C (OC_{min}),
5 determined using density fractionation, and Fe_{DCB} and Fe_O concentrations in sandy, acid forest subsoils
6 and it was suggested that topsoil organo-mineral associations did not follow the same trend as the
7 underlying subsoils because they contained too much OC or insufficient amounts of Fe oxides for the
8 OM to interact, as indicated by OC_{min}/Fe_{DCB} ratio values of ~ 1.5 and 1.4 for subsoils and values up to
9 ~ 14 for topsoils. Poor correlations for the Harwood Forest organic 0-10 cm soils are also likely to be
10 attributable to higher organic loadings of fresh less decomposed OM and/or insufficient Fe/Al oxides
11 (e.g. for the less organic Cambisols). Extractable OC/Fe_{DCB} ratios calculated for Gleysol and Stagnosol
12 0-10 cm, 10-20 cm and 20-30 cm soils were highly variable (see Table 3), and ratios calculated for all
13 three soil types are greater (especially in the Gleysol and Stagnosol topsoils) than the maximum
14 sorptive capacity of 0.22 g OC per g Fe for Fe oxide phases at pH 4-7 determined from sorption
15 experiments (Kaiser and Guggenberger 2006). This indicates that these acidic, OM-rich soils contained
16 organo-Fe associations in the form of precipitated complexes and that mechanisms other than simple
17 sorption are responsible for stabilising OC, consistent with previous findings for similar low pH
18 organic soils (Wagai and Mayer 2007; Wagai et al. 2011). Indeed no or negative correlations between
19 Fe_{DCB} and TOC in neutral soils have been explained by the decreased number of positively charged
20 surface hydroxyls (Kögel-Knabner et al. 2008). Kaiser et al. (2011), however, found that positive
21 relationships between oxalate extractable Al and Fe and SOC contents were strongest for near-neutral
22 arable soils rather than acidic forest subsurface soils such as those studied here.

23 Alongside Fe and Al oxides, other mineral components which contribute to SOM sorption and
24 stabilisation can possibly be removed during the selective dissolution extraction procedures (see
25 Section 3.4.1): oxalate removes Al and Fe in organic complexes (i.e. metal-humus complexes) (Mikutta
26 et al. 2006) and oxalate and DCB dissolve short range order Al silicates (Eusterhues et al. 2005b).
27 Therefore the possible role of other soil components that were extracted alongside Fe/Al oxides and
28 SOM associated with Fe/Al oxides, mainly metal-humus complexes and short range order Al silicates,
29 should be considered. For example, stable OM in acidic soil environments has been found to be
30 predominantly associated with Fe oxides and short-range order Al silicates (Kögel-Knabner et al.

1 2008). The soils under investigation here were not subjected to pyrophosphate extractions (D'Angelo et
2 al. 2009) or chemical pre-treatments with sodium persulfate ($\text{Na}_2\text{S}_2\text{O}_8$) (Eusterhues et al. 2005b) to
3 selectively remove organic Fe and Al complexes in soil samples prior to oxalate extraction. D'Angelo
4 et al (2009) measured both oxalate and pyrophosphate extractable OC and Fe in a range of agricultural
5 and forest soils and found that pyrophosphate extractable forms of Fe were more important for
6 retaining SOC than oxalate extractable forms of Fe. Overall, the correlation approach does not clarify
7 the role of Fe/Al oxides in preserving OC in these gley soils. The interacting factors mentioned above,
8 along with the possibility that SOC contents varied independently of Fe and Al oxide contents as a
9 result of variable surface loadings of different organic inputs at different soil depths (see Section 4.3),
10 could provide explanations for the lack of correlations observed. However, future work involving
11 particle-size in combination with density fractionation and/or pyrophosphate extractions or chemical
12 pre-treatments to remove organic Fe and Al complexes prior to sequential oxalate and DCB extractions
13 could be useful.

14

15 **5. Conclusions**

16 This study has confirmed that total Fe oxides (70-90 % of the total Fe present) in the gley 20-30 cm
17 soils of Harwood forest comprised mainly of strongly crystalline Fe oxide phases (50-80 % of total Fe
18 oxides present). Large amounts of SOC were associated with Fe/Al oxides (generally 70-90 % of total
19 SOC) and therefore, as found in previous microanalytical investigations of these soils, these mineral
20 surfaces appear to play a role in stabilising OC in gley soils. Absence of significant differences
21 between SOC storage in the gley soils studied here, however, may be due to hydrological differences
22 not being great enough to differentially affect the development of redoximorphic conditions and
23 therefore contrasting OM enrichment and depletion patterns. Total OC and extractable OC contents
24 were not found to be linearly related to Fe/Al oxide contents for any of the soil profiles and the lack of
25 linear correlations observed is possibly due to a variety of factors including high loadings of fresh less
26 decomposed OM, especially for the organic 0-10 cm soils, and in some cases insufficient amounts of
27 Fe/Al oxides for OM to interact. Also, variable surface loadings of different organic inputs at different
28 soil depths and contributions of organo-Fe associations in the form of precipitated complexes, as
29 indicated by high OC/Fe ratios in these OM-rich acidic soils may contribute to the lack of linear
30 correlation. Furthermore, contributions from other extractable soil components (e.g. Fe- and other

1 metal-humus complexes, short range order Al silicates) are likely to be important for OM stabilisation.
2 Drawing on statistical test results, significant differences observed suggested that contrasting gley soil
3 hydrological regimes had an effect on total subsoil Fe concentrations, and total and weakly crystalline
4 Al oxide concentrations. Also, significantly different concentrations of strongly crystalline Fe oxides
5 between 10-20 and 20-30 cm Stagnosol soils suggested that the Stagnosols possibly experienced
6 stronger within-profile differences in hydrological conditions and redox oscillations than Gleysols.
7 Geochemical investigations also indicated that mineral phases other than Fe/Al oxides in mineral
8 subsoils preferentially adsorb N-rich microbially processed compounds and that OM associated with
9 weakly and strongly crystalline Fe/Al-oxides in the soils studied had high hydroxyl, aliphatic, carbonyl,
10 aromatic, amide, phenolic, carboxylate and carbohydrate content. Overall this work shows that
11 interactions between Fe/Al oxides and SOM in redox-dynamic gley soils under different hydrological
12 regimes are complex and further investigations of SOC stabilisation in these systems using selective
13 dissolution and other complementary techniques are required.

14

15 **Acknowledgements**

16 We gratefully acknowledge the Natural Environment Research Council (Grant Ref. No.
17 NE/G010102/1) and SAGES for funding this project.

18

1 **References**

2 Avery, B. W. 1973. Soil classification in the soil survey of England and Wales. *Journal of Soil Science*
3 24: 324-338.

4

5 Ball, T., K. A. Smith, and J. B. Moncrieff. 2007. Effect of stand age on greenhouse gas fluxes from a
6 Sitka spruce [*Picea sitchensis* (Bong.) Carr.] chronosequence on a peaty gley soil. *Global Change Biol.*
7 13: 2128-2142.

8

9 Baldock, J. A., and J. O. Skjemstad. 2000. Role of the soil matrix and minerals in protecting natural
10 organic materials against biological attack. *Org. Geochem.* 31: 697-710.

11

12 Bascomb, C. L. 1968. Distribution of pyrophosphate-extractable iron and organic carbon in soils of
13 various groups. *J Soil Sci.* 19: 251-268.

14

15 Bloomfield, C. 1952. The distribution of iron and aluminium oxides in gley soils. *J. Soil Sci.* 3: 167-
16 171.

17

18 Burns, N. R. 2011. Soil organic matter stability and the temperature sensitivity of soil respiration, PhD
19 Thesis, University of Edinburgh.

20

21 D'Angelo, E. M., C. A. Kovzelove, and A. D. Karathanasis. 2009. Carbon sequestration processes in
22 temperate soils with different chemical properties and management histories. *Soil Sci.* 174: 45-55.

23

24 Eusterhues, K., C. Rumpel, M. Kleber, and I. Kögel-Knaber. 2003. Stabilisation of soil organic matter
25 by interactions with minerals as revealed by mineral dissolution and oxidative degradation. *Organic*
26 *Geochem.* 34: 1591-1600.

27

28 Eusterhues, K., C. Rumpel, and I. Kögel-Knaber. 2005a. Stabilization of soil organic matter isolated
29 via oxidative degradation. *Organic Geochem.* 36: 1567-1575.

30

1 Eusterhues, K., C. Rumpel, and I. Kögel-Knaber. 2005b. Organo-mineral associations in sandy acid
2 forest soils: importance of specific surface area, iron oxides and micropores. *Eur J Soil Sci.* 56: 753-
3 763.

4

5 Falsone, G., C. A. Wilson, J. M. Cloy, M. C. Graham, and E. Bonifacio. 2013. Relating microfeatures
6 of soil organic matter to C stabilisation: optical microscopy, SEM-EDS, abiotic oxidation, *Biol. Fertil.*
7 *Soils.* 50: 623-632.

8

9 FAO. 2006. World reference base for soil resources 2006: a framework for international classification,
10 correlation and communication. *World Soil Resources Report*, 103, FAO, Rome.

11

12 Gasparatos, D., C. Haidouti, and D. Tarenidis. 2004. Characterization of iron oxides in Fe-rich
13 concretions from an imperfectly-drained Greek soil: a study by selective-dissolution techniques and x-
14 ray diffraction. *Arc. Agron. Soil Sci.* 50: 485-493.

15

16 Guggenberger, G., and K. Kaiser. 2003. Dissolved organic matter in soils. Challenging the paradigm of
17 sorptive preservation. *Geoderma* 113: 293-310.

18

19 Hedges, J. I., and J. M. Oades. 1997. Comparative organic geochemistries of soils and marine
20 sediments. *Organic Geochem.* 27: 319-361.

21

22 Kaiser, K., and W. Zech. 2000. Dissolved organic matter sorption by mineral constituents of subsoil
23 clay fractions. *J. Plant Nutr. Soil Sci.* 163: 531-535.

24

25 Kaiser, K., K. Eusterhues, C. Rumpel, G. Guggenberger, and I. Kögel-Knaber. 2002. Stabilization of
26 organic matter by soil minerals – investigations of density and particle-size fractions from two acid
27 forest soils. *J. Plant Nutr. Soil Sci.* 165: 451-459.

28

29 Kaiser, K., and G. Guggenberger. 2007. Sorptive stabilization of organic matter by microporous
30 goethite-sorption into small pores vs. surface complexation. *Geochim. Cosmochim. Acta* 71: 25-35.

1
2 Kaiser, M., K. Walter, R. H. Ellerbrock, and M. Sommer. 2011. Effects of land use and mineral
3 characteristics on the organic carbon content, and the amount and composition of Na-pyrophosphate-
4 soluble organic matter. *Eur. J. Soil Sci.* 62: 226-236.
5
6 Kalbitz, K., and K. Kaiser. 2008. Contribution of dissolved organic matter to carbon storage in forest
7 mineral soils. *J. Plant Nutr. Soil Sci.* 171: 52-60.
8
9 Kiem, R., and I. Kögel-Knabner. 2002. Refractory organic carbon in particle-size fractions of arable
10 soils II: organic carbon in relation to mineral surface area and iron oxides in fractions < 6 μm . *Organic*
11 *Geochem.* 33: 1699-1713.
12
13 Kögel-Knabner, I., G. Guggenberger, M. Kleber, E. Kandeler, K. Kalbitz, S. Scheu, K. Eusterhues, and
14 P. Leinweber. 2008. Organo-mineral associations in temperate soils: integrating biology, mineralogy,
15 and organic matter chemistry. *J. Plant Nutr. Soil Sci.* 171: 61-82.
16
17 Krull, E. S., J. A. Baldock, and J. O. Skjemstad. 2003. Importance of mechanisms and processes of the
18 stabilisation of soil organic matter for modelling carbon turnover. *Fun. Plant Biol.* 30: 207-222.
19
20 Lal, R., J. Kimble, E. Levine, and C. Whitman. 1995. World soils and greenhouse effect: An overview.
21 In Lal *et al.* (eds.), *Soils and Global Change*. Lewis Publishers, Boca Raton.
22
23 Mansfeldt, T., S. Schuth, W. Häusler, F. E. Wagner, S. Kaufhold, and M. Overesch. 2012. Iron oxide
24 mineralogy and stable iron isotope composition in a Gleysol with petrogleyic properties. *J. Soils*
25 *Sediments* 12: 97-114.
26
27 Mojeremane, W., R. M. Rees, and M. Mencuccini. 2010. Effects of site preparation for afforestation on
28 methane fluxes at Harwood Forest, NE England. *Biogeochemistry* 97: 89-107.
29

- 1 Mehra, O. P., and M. L. Jackson, M.L. 1960. Iron oxide removal from soils and clays by dithionite-
2 citrate systems buffered with sodium bicarbonate. *Clays Clay Min.* 7: 317-327.
3
- 4 Mikutta, R., M. Kleber, M. S. Torn, and R. Jahn. 2006. Stabilization of soil organic matter: association
5 with minerals or chemical recalcitrance. *Biogeochemistry* 77: 25-56.
6
- 7 Oades, J. M. 1988. The retention of organic matter in soils. *Biogeochemistry* 5: 35-70
8
- 9 Rasmussen, C., R. J. Southard, and W. R. Horwath. 2008. Litter type and soil minerals control
10 temperate forest soil carbon response to climate change. *Global Change Biol.* 14: 2064-2080.
11
- 12 Rasse, D. P., J. Mulder, C. Moni, and C. Chenu. 2006. Carbon turnover kinetics with depth in a French
13 loamy soil. *Soil Sci. Soc. Am. J.* 70: 2097-2105.
14
- 15 Rasse, D. P., C. Rumpel, and M. F. Dignac. 2005. Is soil carbon mostly root carbon? Mechanisms for a
16 specific stabilisation. *Plant Soil* 269: 341-356.
17
- 18 Robertson, G. P., D. C. Coleman, and C. S. Bledsoe. 1999. *Standard Soil Methods for Long-term*
19 *Ecological Research.* Oxford University Press, Oxford.
20
- 21 Schwertmann, U. 1964. Differenzierung der Eisenoxide des Bodens durch photochemische Extraktion
22 mit saurer Ammoniumoxalatlösung. *Z. Pflanzenernährung. Düngung Bodenkunde* 105: 194-202.
23
- 24 Soil Survey Staff. 2010. *Keys to Soil Taxonomy*, 11th ed. USDA-Natural Resources Conservation
25 Service, Washington DC.
26
- 27 Spielvogel, S., J. Prietzel, and I. Kögel-Knaber. 2008. Soil organic matter stabilization in acidic forest
28 soils is preferential and soil type-specific. *Eur. J. Soil Sci.* 59: 674-692.
29
- 30 Stevenson, F. J., *Humus Chemistry*, Wiley, New York.

1
2 Thompson, A., O. A. Chadwick, D. G. Rancourt, and J. Chorover. 2006. Iron-oxide crystallinity
3 increases during soil redox oscillations. *Geochim. Cosmochim. Acta* 70: 1710-1727.
4
5 Torn, M. S., S. E. Trumbore, O. A. Chadwick, and P. M. Vitousek, and D. M. Hendricks. 1997.
6 Mineral control of soil organic carbon storage and turnover. *Nature* 389: 170-173.
7
8 van Oorschot, I. H. M., and M. J. Dekkers. 1999. Dissolution behaviour of fine-grained magnetite and
9 maghemite in the citrate-bicarbonate-dithionite extraction method. *Earth Planet Sci. Lett.* 167: 283-295.
10
11 van Oorschot, I.H.M., and M. J. Dekkers. 2001. Selective dissolution of magnetic iron oxides in the
12 acid-ammonium oxalate/ferrous iron extraction method–I. Synthetic samples. *Geophys. J. Int.* 145:
13 740-748.
14
15 von Lützw, M., I. Kögel-Knabner, K. Ekschmitt, E. Matzner, G. Guggenberger, B. Marschner, and H.
16 Flessa. 2006. Stabilization of organic matter in temperate soils: mechanisms and their relevance under
17 different soil conditions – a review. *Eur. J. Soil Sci.* 57: 426-445.
18
19 von Lützw, M., I. Kögel-Knabner, K. Ekschmitt, H. Flessa, G. Guggenberger, E. Matzner, and B.
20 Marschner. 2007. SOM fractionation methods: relevance to functional pools and to stabilisation
21 mechanisms. *Soil Biol. Biochem.* 39: 2183-2207.
22
23 Wagai, R., L. M. Mayer, and K. Kitayama. 2009. Nature of the “occluded” low-density fraction in soil
24 organic matter studies: a critical review. *Soil Sci. Plant Nutr.* 55: 13-25.
25
26 Wagai, R., and L. M. Mayer. 2007. Sorptive stabilization of organic matter in soils by hydrous iron
27 oxides. *Geochim. Cosmochim. Acta* 71: 25-35.

1 Wagai, R., L. M. Mayer, K. Kitayama, and Y. Shirato. 2011. Association of organic matter with iron
2 and aluminium across a range of soils determined via selective dissolution techniques coupled with
3 dissolved nitrogen analysis. *Biogeochemistry* doi:10.1007/s1053301196525, 112: 95-109.
4

5 Wilson, C. A., D. A. Davidson, and M. S. Cresser. 2009. An evaluation of the site specificity of soil
6 elemental signatures for identifying and interpreting former functional areas. *J. Archeol. Sci.*, 36: 2327-
7 2334.
8

9 Wilson, C. A., J. M. Cloy, M. C. Graham, and L. McKenna. 2013. A microanalytical study of iron,
10 aluminium and organic matter relationships in soils with contrasting hydrological regimes. *Geoderma*,
11 202-203: 71-81.
12

13 Wiseman, C. L. S., and W. Püttmann. 2006. Interactions between mineral phases in the preservation of
14 soil organic matter. *Geoderma* 134: 109-118.
15

16 Zerva, A., T. Ball, K. A. Smith and M. Mencuccini. 2005. Soil carbon dynamics in a Sitka spruce
17 (*Picea sitchensis* (Bong.) Carr.) chronosequence on a peaty gley. *Forest Ecol. Man.* 205: 227-240.
18
19
20

1 **Figure captions**

2

3 **Fig. 1** Soil C and N stocks (t ha^{-1}) at different depths in the Harwood forest Gleysols, Stagnosols and
4 Cambisols. Arithmetic mean value \pm SD of values acquired from five different soil pit profiles are
5 indicated for Gleysols and Stagnosols.

6

7 **Fig. 2** FT-IR spectra for oxalate extracts from a) 10-20 cm and b) 20-30 cm Gleysol, Stagnosol and
8 Cambisol Harwood forest soils.

9

10 **Fig. 3** FT-IR spectra for dithionite-citrate-bicarbonate (DCB) extracts from a) 10-20 cm and b) 20-30
11 cm Gleysol, Stagnosol and Cambisol Harwood forest soils.

12

13 **Fig. 4** FT-IR spectra for Harwood forest Gleysol 10-20 cm and 20-30 cm whole soils before extraction.

14

15

1	List of Supplementary Digital Content
2	
3	Table 1.doc
4	
5	Figures 1 and 2.doc
6	

Figure 1




Figure 2





Figure 3




Figure 4




Table 1: Soil profile descriptions, physical and chemical characteristics of Harwood forest soils based on the mean of five replicate soil profiles (only one profile for the Cambisol) and three replicate sub-samples for each profile (figures in brackets show one standard deviation).

Soil type (sampling depth (cm))	Soil horizon (horizon depth (cm))	Hydrological regime, elevation and grid reference	Visual observations	Water content (% of mass)	Water- filled pore space (%)	Bulk density (g cm ⁻³)	pH	Organic matter content (%)	Particle size distribution (%)			Texture classification
									Sand 60-2000 µm	Silt 2-60 µm	Clay < 2 µm	
Gleysol												
0-10 cm	Ao 0-10 cm	Poorly draining, permanent saturation at depth from ground water. <i>Elevation:</i> 300 m 55° 14' 49" N, 1° 58' 3" W	Dark brown.	82 (23) ^a	-	0.50 (0.03) ^a	4.1 (0.7) ^a	24 (15) ^a	51 ^c	46 ^c	3 ^c	Sandy loam
10-20 cm	A 10-20 cm		Dark greyish brown. Few distinct orange mottles.	37 (4)	60 (48)	0.89 (0.41)	4.5 (1.1)	6.6 (2.0)	53 ^c	44 ^c	3 ^c	Sandy loam
20-30 cm	Bg 20-35+ cm		Brownish grey. Frequent orange mottles.	24 (5)	47 (18)	1.39 (0.54)	4.3 (0.5)	4.4 (1.6)	41 ^c	52 ^c	7 ^c	Silt loam
Stagnosol												
0-10 cm	Ao 0-9 cm	Poorly draining, upper soil horizons saturated by seasonally perched water table. <i>Elevation:</i> 240 m 55° 12' 23" N, 2° 0' 24" W	Blackish grey.	100 (0)	-	0.44 (0.14) ^b	3.9 (0.1) ^b	24 (6) ^b	54 ^c	44 ^c	2 ^c	Sandy loam
10-20 cm	A(g) 9-20 cm		Mottled grey. Frequent distinct mottles.	49 (16)	69 (13)	0.95 (0.21)	4.3 (0.4)	9.2 (3.3)	55 ^c	42 ^c	3 ^c	Sandy loam
20-30 cm	Bg 20-45+ cm		35 (6)	54 (18)	0.99 (0.17)	4.7 (0.8)	5.3 (1.1)	28 ^c	67 ^c	5 ^c	Silt loam	
Cambisol												
0-10 cm	Ah 0-3 cm and A 3-12 cm	Freely draining. <i>Elevation:</i> 240 m 55° 12' 22" N, 2° 0' 22" W	Dark brown. Brown.	66 ^c	-	0.62 ^c	5.4 ^c	8.5 ^c	53 ^c	45 ^c	2 ^c	Sandy loam
10-20 cm	B(g) 12-32+ cm		Greyish brown. Indistinct mottles.	66 (30)	82 (9)	1.08 (0.13)	5.3 (0.1)	6.8 (0.1)	64 ^c	34 ^c	2 ^c	Sandy loam
20-30 cm	B(g)		45 (0)	68 (6)	0.96 (0.06)	5.1 (0.0)	5.4 (0.3)	72 ^c	27 ^c	1 ^c	Loamy sand	

^aValues determined as mean of single soil samples from five replicate profiles, ^bvalues determined as mean of single soil samples from two replicate profiles and ^cvalues determined for single soil samples from one profile.

Table 2: Soil organic carbon and total N concentrations determined for whole soils and extracted soil residues, corresponding C/N ratios and calculated soil C and N concentrations associated with Fe/Al oxides for Harwood forest soils based on the mean of five replicate soil profiles (only one profile for the Cambisol) and three replicate sub-samples for each profile (figures in brackets show one standard deviation).

Soil type (sampling depth (cm))	Whole soil OC concentration (%)	Soil residue OC concentration after oxalate and DCB extraction (%)	Calculated oxalate and DCB extractable SOC concentration (%) <i>[% of total whole soil OC]</i>	Whole soil total N concentration (%)	Soil residue total N concentration after oxalate and DCB extraction (%)	Calculated oxalate and DCB extractable soil N concentration (%) <i>[% of total whole soil N]</i>	Whole soil C/N ratio	Soil residue C/N ratio after oxalate and DCB extraction
Gleysol								
0-10 cm	7.9 (3.5) ^a	2.6 (2.8) ^a	5.3 (2.8) ^a <i>[70 (20)]</i>	0.40 (0.17) ^a	0.081 (0.053) ^a	0.35 (0.18) ^a <i>[78 (10)]</i>	20 (3) ^a	28 (10) ^a
10-20 cm	3.5 (1.0)	0.70 (0.25)	2.8 (0.9) <i>[79 (7)]</i>	0.17 (0.03)	0.038 (0.012)	0.13 (0.04) <i>[76 (11)]</i>	21 (3)	19 (7)
20-30 cm	1.9 (1.1)	0.41 (0.27)	1.5 (0.8) <i>[78 (6)]</i>	0.090 (0.037)	0.025 (0.011)	0.064 (0.032) <i>[69 (11)]</i>	21 (3)	17 (7)
Stagnosol								
0-10 cm	10 (3) ^b	5.6 (4.0) ^b	4.5 (1.4) ^b <i>[48 (27)]</i>	0.53 (0.13) ^b	0.27 (0.19) ^b	0.26 (0.06) ^b <i>[53 (25)]</i>	19 (0) ^b	21 (0) ^b
10-20 cm	2.7 (0.8)	0.46 (0.15)	2.2 (0.8) <i>[81 (9)]</i>	0.14 (0.03)	0.041 (0.011)	0.10 (0.03) <i>[70 (6)]</i>	18 (3)	12 (3)
20-30 cm	1.5 (0.4)	0.28 (0.10)	1.3 (0.3) <i>[81 (4)]</i>	0.093 (0.033)	0.028 (0.009)	0.071 (0.014) <i>[72 (4)]</i>	18 (6)	9 (2)
Cambisol								
0-10 cm	2.2 ^c	1.7 ^c	0.47 ^c <i>[22]</i>	0.13 ^c	0.12 ^c	0.014 ^c <i>[11]</i>	17 ^c	14 ^c
10-20 cm	2.1 (0.2)	0.32 (0.19)	1.7 (0.4) <i>[84 (11)]</i>	0.16 (0.02)	0.026 (0.011)	0.14 (0.03) <i>[83 (10)]</i>	13 (1)	12 (3)
20-30 cm	1.7 (0.2)	0.21 (0.07)	1.5 (0.2) <i>[87 (5)]</i>	0.083 (0.047)	0.033 (0.024)	0.050 (0.064) <i>[46 (40)]</i>	20 (13)	9 (6)

^aValues determined as mean of single soil samples from five replicate profiles, ^bvalues determined as mean of single soil samples from two replicate profiles and ^cvalues determined for single soil samples from one profile.

Table 3. Mean total Al and Fe concentrations (mg g^{-1}), mean oxalate-extractable concentrations of Al (Al_O) and Fe (Fe_O) (mg g^{-1}) and mean total 0.02M oxalate and DCB-extractable Al (Al_DCB) and Fe (Fe_DCB) concentrations (mg g^{-1}) and oxalate and DCB-extractable OC: Fe_DCB ratios for Harwood forest soils. Means are based on the mean of five replicate soil profiles (only one profile for the Cambisol) and three replicate sub-samples for each profile (figures in brackets show one standard deviation).

Soil type (sampling depth (cm))	Total soil Al concentration (mg g^{-1})	Total soil Fe concentration (mg g^{-1})	Al_O (mg g^{-1}) (0.02 M oxalate)	Fe_O (mg g^{-1}) (0.02 M oxalate)	Al_DCB (mg g^{-1}) (0.02 M oxalate and DCB) [% of total soil Al]	Fe_DCB (mg g^{-1}) (0.02 M oxalate and DCB) [% of total soil Fe]	Extractable OC: Fe_DCB ratio (range across Gleysol and Stagnosol profiles)
Gleysol							
0-10	14 (7) ^a	13 (4) ^a	0.37 (0.06) ^a	1.8 (0.5) ^a	0.97 (0.26) ^a [8.3 ± 3.3]	5.4 (1.3) ^a [43 ± 9]	4 - 22
10-20	19 (4)	17 (7)	0.54 (0.19)	1.6 (0.4)	1.6 (0.4) [8.8 ± 3.3]	16 (7) [66 ± 10]	0.74 – 6.7
20-30	25 (11)	23 (4)	1.1 (0.4)	1.6 (0.4)	2.1 (0.8) [11 ± 6]	16 (6) [66 ± 17]	0.72 – 2.9
Stagnosol							
0-10	15 (4) ^b	12 (6) ^b	0.50 (0.13) ^b	1.6 (0.4) ^b	1.2 (0.1) ^b [7.8 ± 2.7]	8.3 (4.1) ^b [69 ± 0]	3 - 10
10-20	18 (5)	10 (3)	0.44 (0.11)	1.3 (0.1)	0.90 (0.10) [5.4 ± 1.7]	6.8 (2.2) [69 ± 3]	1.1 – 7.1
20-30	11 (4)	11 (4)	0.43 (0.13)	1.2 (0.3)	1.0 (0.2) [11 ± 5]	11 (5) [84 ± 5]	0.27 – 3.0
Gleyic Cambisol							
0-10	16 ^c	13 ^c	0.27 ^c	0.80 ^c	0.87 ^c [5.4]	8.4 ^c [63]	0.56
10-20	15 (2)	13 (1)	0.31 (0.04)	1.0 (0.2)	1.0 (0.0) [7.0 ± 0.9]	12 (3) [90 ± 17]	1.5
20-30	11 (4)	13 (2)	0.30 (0.01)	0.88 (0.12)	0.87 (0.17) [8.0 ± 4.7]	10 (5) [75 ± 35]	1.7

^aValues determined as mean of single soil samples from five replicate profiles, ^bvalues determined as mean of single soil samples from two replicate profiles and ^cvalues determined for single soil samples from one profile.

Table 4. Total 0.02 M oxalate plus DCB extractable concentrations of Al (Al_{DCB}) and Fe (Fe_{DCB}) ($mg\ g^{-1}$), total extractable minus 0.2 M oxalate extractable Fe concentrations (Fe_{DCB-O}) ($mg\ g^{-1}$), $Fe_O:Fe_{DCB}$ ratios and $(Fe + Al)_O$ concentrations using 0.2 M oxalate method for selected Harwood forest soil samples (see Supplementary Material Table 1). Gleysol and Stagnosol measurements are the mean of single soil samples from five replicate profiles and Cambisol measurements are values determined for single soil samples from one profile (figures in brackets show one standard deviation).

Soil type (sampling depth (cm))	Al_{DCB} (0.02 M oxalate and dithionate extractable Al) ($mg\ g^{-1}$) ^a	Fe_{DCB} (0.02 M oxalate and dithionate extractable Fe) ($mg\ g^{-1}$) ^a	Fe_{DCB-O} (Total extractable Fe minus 0.2 M oxalate extractable Fe) ($mg\ g^{-1}$)	$Fe_O:Fe_{DCB}$ ratio (0.2 M oxalate method)	$(Fe + Al)_O$ ($mg\ g^{-1}$) (0.2 M oxalate method)
Gleysol					
10-20	1.4 (0.6)	18 (10)	13 (10)	0.32 (0.14)	5.8 (1.9)
20-30	2.5 (0.9)	19 (10)	16 (11)	0.27 (0.20)	5.2 (1.3)
Stagnosol					
10-20	0.89 (0.04)	7.2 (2.0)	1.5 (0.8)	0.81 (0.09)	6.8 (1.5)
20-30	1.2 (0.20)	15 (6)	9.9 (5.3)	0.37 (0.10)	6.2 (1.5)
Gleyic Cambisol					
10-20	1.0	14	6.4	0.35	8.4
20-30	1.0	13	7.1	0.29	7.0

^aFor direct comparison, values quoted here are different from those in Table 3 because they represent values measured in the samples selected for test extractions.

Table 5. Summary of independent t-test results indicating whether differences in soil measurements at each depth for Gleysols and Stagnosols were statistically significant, where differences were deemed significant if $P \leq 0.05$, s = significant, ns = not significant.

Soil measurement	Gleysol (n = 5) and Stagnosol (n = 2) Ao 0-10 cm P value	Gleysol (n = 5) and Stagnosol (n = 5) A 10-20 cm P value	Gleysol (n = 5) and Stagnosol (n = 5) Bg 20-30 cm P value
Whole SOC concentration (%)	0.475 (ns)	0.174 (ns)	0.500 (ns)
Whole soil total N concentration (%)	0.382 (ns)	0.199 (ns)	0.900 (ns)
Whole soil C/N ratio	0.481 (ns)	0.302 (ns)	0.379 (ns)
Whole SOC stock (t ha^{-1})	0.995 (ns)	0.115 (ns)	0.229 (ns)
Whole soil total N stock (t ha^{-1})	0.744 (ns)	0.645 (ns)	0.172 (ns)
Total soil Al concentration (mg g^{-1})	0.894 (ns)	0.645 (ns)	0.056 (ns)
Total soil Fe concentration (mg g^{-1})	0.860 (ns)	0.128 (ns)	0.001 (s)
Al _{DCB} (0.02 M oxalate and DCB) (mg g^{-1})	0.221 (ns)	0.021 (s)	0.032 (s)
Fe _{DCB} (0.02 M oxalate and DCB) (mg g^{-1})	0.510 (ns)	0.059 (ns)	0.210 (ns)
Al _O (0.02 M oxalate) (mg g^{-1})	0.522 (ns)	0.376 (ns)	0.022 (s)
Fe _O (0.02 M oxalate) (mg g^{-1})	0.372 (ns)	0.158 (ns)	0.087 (ns)
Fe _{DCB-O} (Total extractable Fe minus 0.2 M oxalate extractable Fe) (mg g^{-1})	-	0.059 (ns)	0.327 (ns)
Fe _O :Fe _{DCB} ratio (0.2 M oxalate method)	-	0.001 (s)	0.357 (ns)
(Fe + Al) _O (0.2 M oxalate method) (mg g^{-1})	-	0.380 (ns)	0.308 (ns)
Oxalate and DCB extractable SOC concentration (%)	0.618 (ns)	0.295 (ns)	0.571 (ns)
Oxalate and DCB extractable soil N concentration (%)	0.398 (ns)	0.311 (ns)	0.707 (ns)