

Scotland's Rural College

Predicting the effect of rotation design on N, P, K balances on organic farms using the NDICEA model

Smith, LG; Tarsitano, D; Topp, CFE; Jones, SK; Gerrard, CL; Pearce, BD; Williams, AG; Watson, CA

Published in:
Renewable Agriculture and Food Systems

DOI:
[10.1017/S1742170515000381](https://doi.org/10.1017/S1742170515000381)

Print publication: 01/01/2015

Document Version
Peer reviewed version

[Link to publication](#)

Citation for published version (APA):

Smith, LG., Tarsitano, D., Topp, CFE., Jones, SK., Gerrard, CL., Pearce, BD., Williams, AG., & Watson, CA. (2015). Predicting the effect of rotation design on N, P, K balances on organic farms using the NDICEA model. *Renewable Agriculture and Food Systems*, 31, 471 - 484. <https://doi.org/10.1017/S1742170515000381>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

1 **Predicting the effect of rotation design on N, P, K balances on organic farms**

2 **using the NDICEA model**

3 Laurence G. Smith^{1,2}, Davide Tarsitano³, Cairistiona F. E. Topp³, Stephanie K. Jones³,
4 Catherine L. Gerrard¹, Bruce D. Pearce¹, Adrian G. Williams^{2*} and Christine A. Watson⁴

5 **Abstract**

6 The dynamic model NDICEA (Nitrogen Dynamics In Crop rotations in Ecological Agriculture) was used
7 to assess the Nitrogen (N), phosphorus (P) and potassium (K) balance of long term organic cropping
8 trials and typical organic crop rotations on a range of soil types and rainfall zones in the UK. The
9 measurements of soil N taken at each of the organic trial sites were also used to assess the
10 performance of NDICEA. The modelled outputs compared well to recorded soil nitrogen levels, with
11 relatively small error margins. NDICEA therefore seems to be a useful tool for UK organic farmers.
12 The modelling of typical organic rotations has shown that positive N balances can be achieved,
13 although negative N balances can occur under high rainfall conditions and on lighter soil types as a
14 result of leaching. The analysis and modelling also showed that some organic cropping systems rely
15 on imported sources of P and K to maintain an adequate balance and large deficits of both nutrients
16 are apparent in stockless systems. Although the K deficits could be addressed through the buffering
17 capacity of minerals, the amount available for crop uptake will depend on the type and amount of
18 minerals present, current cropping and fertilisation practices and the climatic environment. A P
19 deficit represents a more fundamental problem for the maintenance of crop yields and the organic

¹ The Organic Research Centre, Hamstead Marshall, RG20 0HR, UK

² School of Environment, Energy and AgriFood, Cranfield University, Bedford, MK43 0AL, UK

³ Crop and Soils Systems, Scotland's Rural College (SRUC), West Mains Rd, Edinburgh, EH9 3JG, UK

⁴ Crop and Soils Systems, Scotland's Rural College (SRUC), Craibstone Estate, Aberdeen, AB21 9YA, UK

1 sector currently relies on mined sources of P which represents a fundamental conflict with the
2 International Federation of Organic Agriculture Movements (IFOAM) organic principles.

3 *Corresponding author: adrian.williams@cranfield.ac.uk

4• Keywords: organic farming; nutrients; sustainability; crop rotation

5 **Introduction**

6
7 Organic cropping systems focus on feeding the soil, rather than the plant, to build long-term system
8 health and resilience^[1]. This approach results in a reliance on fertility building ley periods and the
9 application of composts and manures, which supply a source of nutrition for the growing crops
10 whilst potentially improving the soil microbial life and organic matter contents^[1, 2]. The length of
11 the ley period can vary from short term (12-18 months) to long term (around 5 years), but typically
12 the ley is kept for about 18 months to 3 years. In Europe, organic farmers most frequently use grass-
13 clover mixes for their leys, with white clover (*Trifolium repens*) and red clover (*T. pratense*) being
14 popular legume species and perennial ryegrass (*Lolium perenne*) and Italian ryegrass (*L. multiflorum*)
15 as commonly chosen grass species^[3]. The crops following the ley period make use of the built-up
16 fertility, although the ley period can also remove fertility, in particular K in conserved grass (silage).
17 Cropping following the ley phase often includes rotational use of over-winter green manures and
18 cover crops such as cereal rye (*Secale cereale*) and vetch (*Vicia satvia*) to reduce losses and to supply
19 additional N through biological fixation^[1]. The use of these approaches on organic farms creates
20 systems in which the nitrogen supplied is in a less available form, compared with conventional
21 systems using mineral fertiliser^[4]. The supply of available nitrogen in organic systems can therefore
22 be a limiting factor for the maintenance of crop yields^[5-7]. In addition, poor synchronicity between
23 the supply and demand for nitrogen can lead to leaching and gaseous losses, particularly following
24 ley cultivation. Nevertheless, this is also an issue for conventional farmers, particularly following
25 periods of high rainfall^[6, 8, 9]. Under organic management, the surplus of N following ley cultivation

1 can be followed by an N deficit later in the crop rotation^[5]. Although this shortage can be resolved
2 through the application of organic composts, manures, and/or through the use of short-term green
3 manures, it can be difficult to match the N supplied from such sources with crop demand. A reliance
4 on such methods can therefore contribute to lower nitrogen use efficiencies compared to non-
5 organic systems applying targeted mineral N^[4, 6]. Despite the challenges of N availability and
6 synchronicity of N supply/demand on organic farms, Berry *et al.*^[10] found positive N balances in a
7 comparison of nine organic farms, and reported that the farms were probably sustainable in terms
8 of N supply and offtake. However, the same study found that phosphorus (P) and potassium (K)
9 levels were in deficit within the stockless systems assessed, and that only farms with large manure
10 returns from stock fed with bought-in feed had a positive or neutral K budget. Korsæth *et al.*^[11] and
11 Torstensson *et al.*^[4] also found P and K deficits within organic arable cropping and mixed dairy
12 farming systems in Norway and Sweden respectively.

13 The research presented here aimed to assess the effect of rotation design on the supply and
14 offtake of nitrogen (N), phosphorus (P) and potassium (K) on organic farms using the dynamic model
15 NDICEA (Nitrogen Dynamics In Crop rotations in Ecological Agriculture). Five hypotheses were posed
16 at the outset of this study. First, NDICEA can effectively calculate the course of mineral-N over a
17 range of organic crop rotations. Second, the N supplied through biological fixation in stockless
18 organic rotations is sufficient to support crop offtake. Third, organic cropping systems incorporating
19 livestock manure applications are able to maintain a positive or neutral N, P and K balance. Fourth,
20 organic rotations will typically rely on imported P to maintain a balance of this nutrient. Fifth, a
21 deficit of K is a common feature in the overall nutrient balance of typical organic crop rotations.

22

1 **Methods**

2 The NDICEA model^(17,18) was applied to assess the supply and demand of nitrogen (N), phosphorus
3 (P) and potassium (K) within a range of stocked (i.e. with manure) and stockless rotations applied at
4 experimental organic farms in the UK. In addition, typical organic rotations were drawn from the
5 literature.

6 **Model description**

7 NDICEA is a dynamic, target-oriented model with crop yield and crop quality parameters, e.g. dry
8 matter, N, P and K contents, used as a basis for crop uptake calculations. Mineralisation of nitrogen
9 from soil organic matter and organic inputs such as manure and compost is also calculated, factoring
10 in the effects of weather, irrigation and soil type, although the model does not account for
11 volatilisation losses during composting / storage of manure. The model uses a daily time step,
12 utilising site specific weather data (rainfall, temperature, evapotranspiration) and user-defined soil
13 and crop parameters. Although the model contains default values for a range of soil types these
14 values can be automatically adjusted through the addition of data on measured soil mineral nitrogen
15 and soil organic matter within the user interface. Following the entry of these values calibration of
16 the model takes place through the implementation of an algorithm that selects an optimum
17 parameter value from a range of plausible values for such variables as N leaching, denitrification and
18 water holding capacity^[12]. Within this study measured values of soil mineral nitrogen and soil
19 organic matter were used to calibrate the model runs and improve the accuracy of the assessments
20 (see Table 3). A repeat calculations function within the model also allows the user to assess the
21 longer term impacts of rotations both in terms of the nutrient supply and the effect on organic
22 matter stocks. The focus of the model is on nitrogen dynamics. For P and K, a simpler farm-gate
23 balance approach is taken (i.e. **only crop offtake and atmospheric deposition is calculated**, based on
24 the user-defined input parameters and/or default values). The calculations for P and K are also

1 unaffected by changing the soil type or daily rainfall and evapotranspiration values within the model.
2 The wide range of cover crops and green manure-options within the NDICEA interface makes the
3 tool particularly applicable for organic farmers, however the tool can also be used to improve
4 understanding of nitrogen dynamics under non-organic management^[12]. Under both organic and
5 non-organic management model performance will be improved by calibration, with a higher number
6 of measurements improving the accuracy of the estimates of N supply and losses^[13].

7 **Description of sites and cropping systems**

8 The model was run using crop, soil and weather data from the UK Government funded
9 organic conversion trials held at ADAS Terrington^[14], Warwick University's Hunts Mill site^[15] as well
10 as other long term trials at Elm Farm Research Centre (EFRC)^[16], Scotland's Rural College (SRUC
11 Tulloch and Woodside^[17]) and a grazing only trial at the Institute of Biological, Environmental and
12 Rural Sciences (IBERS) at the University of Aberystwyth (Ty Gwyn)^[18]. Please see Table 1 and Figure
13 1 for more information on the trials.

14 Soils data from each site were collected from project reports, site records and published
15 literature^[14-18]. The bulked soil samples at each site were taken along a W transect twice each year in
16 the case of Elm Farm^[16], Warwick University^[15] and ADAS Terrington^[14] (after sowing and harvest)
17 and once per year at the SRUC sites^[19] and at Ty Gwyn^[18] (January and April respectively). Samples
18 were analysed for available P (Modified Morgan's solution at SRUC sites and Olsen's method at
19 other sites), available K (Modified Morgan's solution at SRUC sites and ammonium nitrate extraction
20 at ADAS and Elm Farm), mineral nitrogen (potassium chloride solution) and organic matter (loss on
21 ignition). Soil samples were taken at a range of depths. At Elm Farm separate topsoil (0-15cm) and
22 subsoil (15-30cm) samples were assessed for the above parameters. At Warwick University
23 assessments were carried on samples from 0-30cm and 30-60cms. At Ty Gwyn and mineral nitrogen
24 was sampled to 80cm in 15cm and 20cm increments respectively, although only the first sample

1 layer was assessed for P, K and organic matter. At ADAS Terrington all samples were taken to 90cm
2 in 15cm increments. At Woodside, the mineral nitrogen was sampled to 45cm in increments of
3 15cm, and at Tulloch, the mineral nitrogen was sampled to 30cm in increments of 15cm.

4 The rotations applied at EFRC and ADAS Terrington were managed as stockless systems,
5 although phosphate fertilisers permitted under organic standards were applied. The EFRC trial
6 received lime up to a maximum rate of 2 tonnes per hectare per year. Lime was similarly applied at
7 the ADAS site in order to keep the pH between 6 and 6.5. All of the red clover leys at each site were
8 managed through cutting and mulching. The Hunts Mill plot trials included both 'with manure' and
9 'without manure' treatments. Both sites at Hunts Mill received a single application of green waste
10 compost at a rate of 20 tonnes per hectare. At the Ty Gwyn organic dairy unit, manure was
11 deposited at a rate consistent with 2 Livestock Units (LSU) per hectare and lime was applied at a
12 mean rate of 0.7 t/ha over 3 years. At both of the SRUC sites (i.e. Tulloch and Woodside) total
13 annual manure applications were based on 2.8 LSU/ha for the period 1991-1998. In addition ground
14 limestone and potassium sulphate (K_2SO_4) were applied to all Woodside plots in 1991, at a rate of
15 3.75 t/ha and 150 kg/ha respectively. All grass-clover leys at SRUC sites were managed through a
16 combination of grazing with sheep and cutting for silage as described in Taylor et al.^[17].

17 In addition to data from the organic trials, information on typical organic rotations was
18 gathered based on examples within the Organic Farm Management Handbook^[20] and following
19 guidelines given to organic farmers with respect to the proportion of fertility building leys to
20 exploitative phase^[1,20] (see Table 2 for description of the rotations used). Manure application rates
21 for the typical stocked cropping systems were derived using typical livestock numbers for cropping
22 farms (i.e. 0.3 Grazing Livestock Units per Utilisable Agricultural hectare) reported within a sample of
23 approximately 30 organic farms included within the FBS (Farm Business Survey) for England and
24 Wales^[21-23]. Rock phosphate application rates were derived with expert input from the Institute of
25 Organic Training and Advice (IOTA) registered advisers.

1 The rotations were chosen to represent a range of stocked and stockless organic cropping
2 systems. To assess the representativeness of the rotations crop areas were compared to those
3 reported for a stratified sample of 30 organic farms included within the Organic Farm Income
4 Reports published by Aberystwyth University and The Organic Research Centre ^[21-23]

5
6 As shown in Figure 2, the typical rotations are broadly representative of the crop areas
7 reported on actual organic farms within the Farm Income Reports' matched sample. Although there
8 are some differences by crop type (e.g. both stocked simple and stockless simple containing a high
9 percentage of cereal crops) the differences are generally in the region of 15-20%. In view of the wide
10 variation between the rotations on individual farms, this is an acceptable margin of error and the
11 rotations applied here can be considered to be broadly representative of organic cropping farms.

12

13 **Model application**

14 The model was applied to assess the effect of rotation design on the supply and offtake of
15 nitrogen (N), phosphorus (P) and potassium (K) on the above experimental sites and within typical
16 organic rotations. The measured changes in soil organic matter over time were small for most of the
17 sites assessed (data not shown). It was therefore necessary to run the model a number of times to
18 ensure a minimal gain or loss of soil organic matter and to avoid erroneous conclusions. Based on
19 measured data and results from long term experiments ^[24, 25] a uniform, near steady-state was
20 assumed to have been reached once the annual change in soil organic matter was less than 2% of
21 the total organic matter pool (expressed in kg/ha) over the rotation. **Two runs of the model were**
22 **implemented for each site, an uncalibrated run, using only the basic, user-adjusted soil and crop**
23 **parameters, and a calibrated run following the input of measured amounts of soil mineral N and soil**
24 **organic matter to make the model automatically adjust advanced soil parameters such as N leaching**

1 **and denitrification factors**. The root mean square errors (RMSE) were calculated based on the size of
2 the deviation between the measured soil N and modelled soil N values and the number of samples
3 at each site. The observed N values were used therefore in the calibrated runs as both inputs to the
4 model and as comparator for assessing model performance. Following the calculation of N, P and K
5 balances for the trial sites, a further application of the model was implemented for the typical
6 organic rotations described above, using the same soil and weather conditions as the trial sites.

7 **Results and discussion**

8 The modelled estimates for Soil Mineral Nitrogen (SMN) were compared with the sampled
9 soil N values from each of the sites to test the ability of the model to simulate the measured
10 rotations. An NPK balance for each of the rotations was then calculated.

11 **Comparison of the NDICEA model's estimates for soil N with measured mineral N values** 12 **at each of the trial sites**

13 A RMSE of 20 kg N ha⁻¹ or less was proposed by Van der Burgt *et al.*^[12] to represent
14 acceptable model performance for practical purposes. This could be achieved for most of the sites,
15 although in some cases (e.g. EFRC and Hunts Mill) the modelled results are above this value (see
16 calibrated model output in Table 3). The higher errors at EFRC could be a result of the small number
17 of measurements (i.e. 7). The high errors for the stocked rotations at Hunts Mill could be explained
18 by the fact that soil N measurements were taken soon after application of manures (the modelled
19 values for this phase of the rotation were more than 100 kg N/ha lower than the recorded values). It
20 may also be possible that the model is underestimating the nitrogen supplied or that the deeper
21 samples at Hunts Mill (0-60cm) resulted in a mixing of topsoil and subsoil layers, and a subsequent
22 overestimate of the mineral N content in the subsoil layer.

1 **NPK balance for each of the rotations applied at the organic trial sites**

2 Modelled nutrient balances derived from NDICEA are presented in Table 4 for each of the stockless
3 rotations. The results include an estimate of the change in soil organic matter, with a negative
4 number indicating mining of existing reserves and a positive number indicating an assimilation of N
5 to soil organic matter (i.e. an increase in organic matter stocks).

6 Table 4 illustrates that the amount of N supplied through biological fixation could potentially
7 support the crop removal at the EFRC trial, although due to losses from the system, in particular
8 leaching, much of this N is lost. The negative values for organic N indicated a mining of organic
9 matter over the course of the experiment. This decline was observed through field measurements
10 with organic matter levels dropping from 32 g/kg⁻¹ to 25 g/kg⁻¹ of soil, probably as a result of starting
11 the trial after a 5-6 year ley. Organic matter levels would be expected to rise again on return to a
12 longer-term ley period, as is standard organic practice. The increase in organic matter levels from
13 the implementation of a one-year fertility building ley was reflected in the NDICEA model, which
14 showed a rise during this period, although the subsequent decline more than offset the gain. When
15 a nutrient demanding crop was introduced into the rotation (e.g. potatoes in EFRC B) the deficit for
16 all three nutrients increased to the extent that further use of external inputs (e.g. composts or
17 manures) would be required, in particular for P and K. The high P and K offtake of beans similarly
18 contributed to the large deficit of these two nutrients within the balance for rotation EFRC C.

19 A similar picture is presented for the rotation at ADAS Terrington. Despite the large
20 contribution of N through fixation, there are considerable losses from leaching and denitrification,
21 resulting in a negative N balance. It may be possible to address this problem through better use of
22 over-winter cover crops (cover crops performed poorly over the course of the experiment with only
23 one crop yielding over 1 t DM/ha). The deficits for P within this trial are also unsustainable in the
24 long-term and would need to be addressed through imports or reducing the exploitative phase of

1 the rotation. The K deficits for this site are also substantial, however it is possible that weathering
2 of K stocks in the mineral pool could redress this^[26].

3 Lower rates of leaching were found for both Hunts Mill plots, although considerable deficits
4 of P and K were also found despite the addition of green waste compost on area 1. The K deficits
5 could be addressed through the buffering capacity of K-bearing minerals^[26] however the P deficit
6 represents a more fundamental issue for the maintenance of crop yields in the longer term^[27]. The
7 results for Hunts Mill area 6 also illustrate that it is possible to maintain a fairly balanced system with
8 regard to N through the effective use of late summer/autumn sown green manures (i.e. without the
9 use of a ley/break crop), although the overall deficit for N may result in a reduction in offtake or the
10 need to use imported composts or manure.

11 Most of the stocked rotations were found to be more balanced with regard to N and P
12 supply and loss (see Table 5). However, all of the SRUC sites faced a large K deficit, due to the high
13 offtake from grass/clover silage, the potato crop (only at Woodside) and the use of straw for
14 bedding, which were not offset by the manure application. Similar results were found within the
15 nutrient balances for Tulloch and Woodside calculated by Watson et al.^[28] although lower offtake
16 was estimated within this study due to the lower assumptions on P and K content within NDICEA.
17 Despite the K deficit, no trend in K levels was found over time at Tulloch or Woodside, although the
18 soil samples were restricted to the first 30cm and 45cm due to the presence of indurated layers at
19 deeper levels, largely impenetrable to soil augers or crop roots. It is possible that soil K levels at
20 these sites were being supplemented by reserves within parent material, in addition to potential
21 inputs from crop residues (these inputs are ignored in standard soil K measurements and in part
22 explains why test values are unrelated to crop K balances^[26]). These and other factors lead Khan et
23 al.^[26] to suggest that measurements of available soil K are an unreliable indicator and that producers
24 should use strip trials to determine site-specific fertiliser management. At both Woodside sites, the
25 rate of N leaching was higher than at Tulloch, despite a lower annual rainfall. This was in part

1 related to the lighter soil texture and the relatively low yield of the grass/clover leys at Woodside 37
2 (4-6 t DM/ha, Taylor *et al.*^[17]) which was related to the high soil moisture deficit. The low
3 grass/clover yield at Woodside 37 also led to a negative N balance overall, due to a lower rate of
4 biological N fixation. Much of the excess nitrogen at the other SRUC sites was locked up as organic
5 matter (illustrated as a positive value under 'Change in organic N' in Table 5). This was observed
6 within the trial through a small increase in soil organic matter levels observed at Tulloch, although
7 the measured organic matter levels at Woodside remained relatively constant. Volatilization rates
8 were low across all of the stocked rotations in Table 5 as a result of incorporating applied manure on
9 the same day as application on the trial sites.

10 **Nutrient (NPK) balance for typical rotations applied using site conditions of the organic** 11 **trials**

12 Nutrient balances are presented in Table 6 for each of the typical rotations described in Table 2.
13 The results presented below are mean values across all six sites and associated soil/weather
14 conditions.

15 The stocked complex rotation described above seems to represent a well-balanced system
16 with regard to N, and P supply and offtake. However, the model predicted a relatively large K deficit
17 with offtake exceeding supply. As discussed earlier, this could be addressed through K delivery from
18 the weathering of minerals depending on the underlying geology, climatic conditions and site
19 management^[26, 29], or through imported compost and/or mineral sources. The higher proportion of
20 nutrient-demanding crops (e.g. potatoes) within the stocked simple rotation creates a larger K deficit
21 compared to the stocked complex example. As with the stocked experimental sites in Table 5,
22 volatilization rates were low for all of the stocked rotations in Table 6, as a result of selecting same-
23 day incorporation of manure applied within NDICEA, and the low stocking density (i.e. 0.3 LSU per
24 hectare). The volatilisation losses would be expected to increase if incorporation was delayed for
25 any reason.

1 The stockless complex rotation has a deficit for all three nutrients. Two years of a red clover ley plus
2 one year of spring beans did not provide enough N to support four years of crop offtake due to a
3 high rate of leaching and denitrification. The presence of nutrient demanding crops contributes to
4 the deficit (i.e. potatoes lead to a high N and K demand and beans to a high K offtake). The stockless
5 simple rotation faces less of a deficit with respect to N, due to a higher input of biologically fixed N
6 from the inclusion of peas which have a higher rate of N fixation than beans within NDICEA, and the
7 use of the grass/vetch over-winter green manures following the spring crops. In addition there is an
8 absence of nutrient demanding crops (e.g. potatoes) however the rate of leaching is still high. The
9 relatively low deficit of P within all of the typical rotations is a result of the application of rock
10 phosphate. All of the modelled rotations would face a P deficit on a similar scale to the K balance
11 without the use of this input.

12 **Implications for improved organic management**

13 In common with previous studies, the work presented here found considerable rates of N leaching
14 within the rotations assessed ^[4, 10, 30]. In some cases, this exceeded the amount lost by product
15 removal (e.g. the stockless simple rotation described in Table 2). High rates of leaching under
16 organic management are related to difficulties associated with matching crop N demand with N
17 availability, particularly following incorporation of the ley when N availability exceeds demand ^{[4, 31,}
18 ^{32]}. The use of organic manures can also make it difficult to predict N availability, compared with
19 applications of mineral fertiliser ^[33], making it more difficult to maximise N recovery and crop yields
20 under organic management ^[34]. As a result of these factors, lower nitrogen use efficiency has been
21 reported for organic cropping in comparisons with conventional systems ^[4, 31].

22 The effective use of over winter green manures and undersowing of leys in cereal crops will
23 help to reduce losses and thus enhance overall N efficiency ^[4, 35] and the lowest rates of leaching
24 within this study were found for the rotations incorporating undersown crops and cover-crops (e.g.
25 ADAS Terrington, Warwick University Hunts Mill). Poor cover-crop establishment (e.g. at the ADAS

1 and Elm Farm experiment) was experienced as a result of competition from weeds and slow
2 emergence which reduced the benefit obtained^[36]. Poor cover crop establishment can also be
3 related to competition from the cash crop, adverse weather conditions and low soil temperatures at
4 the time of sowing^[37]. In particular, the occasional occurrence of poorly performing cover crops
5 presents an important challenge for the long-term sustainability of stockless systems, which rely on
6 keeping the N supplied through biological fixation within the system. Although with careful rotation
7 design, such systems are, in theory, sustainable from a nitrogen management perspective^[38], in
8 practice these systems appear to be highly vulnerable to poor establishment during the cover-
9 cropping period.

10 The use of cover crops is not limited to organic farms, and higher nitrogen use efficiencies
11 can be obtained by using this method alongside targeted mineral fertiliser application(s) to meet
12 crop demand (and thus increase yield) whilst minimising losses^[4]. Such tightly controlled systems
13 could represent a suitable approach to developing highly N-efficient production systems, through a
14 combination of organic practices and targeted fertiliser application^[33, 39]. Similar targeted
15 approaches could still be used on organically-managed land, through the use organic fertilisers with
16 a high N availability (e.g. poultry manure and digestate from slurry based anaerobic digestion) to
17 supply readily available N at key points in the rotation^[5, 40]. However the application of such
18 sources can increase the occurrence of nitrophilous weeds and their use within organic systems has
19 been questioned as the high N availability leads to feeding the plant instead of the soil^[41] and a
20 reduction in the amount of organic matter applied in the case of digestate^[42, 43]. The use of
21 perennial crops can also help to reduce leaching in organic systems through keeping the soil covered
22 and improving N synchrony^[44, 45], although lower yields, weed susceptibility and pest and disease
23 management issues may limit uptake^[46]. A lack of technical information, suitable varieties and
24 socioeconomic constraints (e.g. lower consumer demand compared to staple annual crops) also limit
25 the potential for a wider adoption of perennial cropping^[46, 47].

1 Organic farmers can also reduce N leaching considerably through improved management of
2 manures and slurries. **In particular careful storage, application timing and choice of application**
3 **method will help to maximise N recovery and minimise losses where slurries and manures are**
4 **applied** ^[48, 49]. Manure analysis can also improve on farm nutrient use efficiency and help to reduce
5 losses by improving understanding of nutrient supply from organic sources ^[50]. In some regions,
6 there may be opportunities for farmers to work together to measure the nutrient use efficiency of
7 their systems through a combination of manure and livestock dietary analysis combined with soil
8 sampling ^[51, 52]. Such participatory approaches can be effective at allowing for improvement options
9 to be identified and for the fine-tuning of production systems. Again the use of such methods is not
10 restricted to organic farms, however the inability of such farms to access manufactured N fertiliser
11 makes the implementation of such measures all the more important for the effective prediction of N
12 supply.

13 With regard to phosphorus, the modelled systems able to achieve a sustainable balance
14 were using external inputs of rock phosphate to offset losses. **Although rock phosphate can help to**
15 **offset losses, a reliance on this source may result in limited P bioavailability to meet crop demand,**
16 **due to slow rates of solubilisation** ^[53]. In addition, the use of such a fertiliser clearly does not fit well
17 with the IFOAM organic principles ^[54] which emphasise the importance of reducing inputs to increase
18 the long-term sustainability of farming systems. Despite this aim, the use of imported manure,
19 straw and/or rock phosphate is common on organic farms, particularly for the supply of P and K ^{[55-}
20 ^{57]}. In many cases, manure and straw is sourced from conventional farms, which has led to the
21 conclusion that organic farms are being 'propped-up' by conventional agriculture, and that as a
22 result a large-scale conversion to organic management would be unsustainable ^[55-57]. Organic
23 monogastric systems (in particular poultry) also often require imported feed (e.g. soy) to supply
24 protein and essential amino acids ^[58] and so these systems **are supplemented** by internationally
25 imported P and K.

1 The use of household waste and sewage sludge on organic farms could represent a possible
2 solution to reduce the reliance on conventional manure and/or rock phosphate on organic cropping
3 farms, in particular for the sustainable supply of P. Source separated urine also presents an
4 opportunity to apply readily available nitrogen and phosphorus^[57, 59, 60]. The use of such sources
5 clearly fits with the organic ideal of closing the system as far as possible^[42], although in this case the
6 'system' expands beyond the farm gate to the consumer^[56]. Although there have been many cases
7 of household waste recycling on organic farms to supply nutrients to the soil^[61-63], the use of sewage
8 sludge or urine is strictly prohibited on organic land in Europe, despite the fact that its use seems to
9 be a rational and scientifically supported method of closing the nutrient cycle. Developments in the
10 area of struvite (magnesium ammonium phosphate) recovery from waste water treatment plants
11 could present a possible solution, allowing for application of a refined and slow release mineral
12 fertiliser product, however this product is not currently on the list of permitted fertiliser within the
13 European Commission organic regulation^[64]. This is an area that needs further scrutiny from a
14 scientifically based perspective as it would appear that historical concerns about the toxic effects of
15 applying urine and sewage sludge to agricultural land may no longer be justified^[60, 65], although
16 public perception concerning the risks to human health remains an issue in some areas^[66].

17 Increasing the co-operative use of manure between (organic) livestock and arable farmers has also
18 been suggested as a possible route for reducing the use of conventional manures on organic land
19 and, within farming in general, the co-operative use of manure between specialised livestock and
20 arable holdings could contribute to the prevention of stockpiling of nutrients and associated losses
21 on intensive livestock holdings^[67, 68]. In particular this approach has been encouraged in Denmark
22 by a decision to phase out the use of conventional manure and straw on organic land by 2021, partly
23 in recognition of the conflict between principle and practice and partly to prevent the import of
24 genetically modified organisms (GMOs) into organic systems via manure^[42]. In addition, the
25 transition strategy in Denmark has highlighted the importance of crop rotation design (in particular

1 to improve understanding on nutrient supply and losses), the development of crop cultivars for low-
2 nutrient environments, and the development of biogas plants that can run on plant-based feedstock
3 (in particular grass/clover harvested from leys) in recognition of the limited supply of organic
4 manure ^[42, 69].

5 Potassium deficits were observed across all of the rotations however on many soils, this
6 does not present an issue given vast reserves of mineral K within parent material which may be
7 released for plant uptake by weathering^[26]. Despite this potential, Holmqvist et al.^[70] found that
8 weathering and bioavailability from the mineral fraction can vary greatly (between 3 and 80 kg K
9 ha⁻¹yr⁻¹ on a range of soil types in Norway, Sweden and Scotland) although the modelled predictions
10 in this study did not take into account the dynamic and localised biological weathering by plant roots
11 illustrated by x-ray diffraction studies (e.g. Hinsinger et al.^[71] in Khan et al.^[26]) and the potential
12 contribution of mycorrhizal fungi to K availability^[72]. Nevertheless improved knowledge of site-
13 specific geochemical and mineralogical data in addition to soil rhizosphere interactions, could be a
14 useful aid to develop site-specific fertiliser recommendations and nutrient balances ^[70, 73]. With
15 respect to mineral reserves of K on this sites assessed in this study, only EFRC, IBERS and ADAS
16 Terrington could be expected to supply a considerable amount of K from the clay fraction^[74],
17 although sand- and silt-sized muscovite and biotite can also be a major source of plant-available K on
18 lighter soils^[75] and the presence of these and other K-bearing minerals may have offset some or all
19 of the K offtake at Hunts Mill and the Scottish sites^[76]. Despite the high deficits, there was no
20 apparent trend in available K levels over time at most of the experimental sites considered, although
21 Hunts Mill showed a slight decline over the course of the study and the K measurements at Tulloch
22 (taken in the winter) may have been affected by the preceding silage crop ^[28]. Other studies have
23 demonstrated a decline in soil P and K levels following conversion to organic management ^[4, 77] and
24 positive yield responses have been observed following K applications in long-term experiments in
25 Australia and the UK^[78] and within rice production systems, following several years of intensive

1 cropping^[79]. It is thus important to use nutrient budgets together with soil analysis to help
2 understand the buffering capacity of soils and the management of P and K on individual fields. It
3 should also be remembered that the bank-balance (i.e. supply minus offtake) concept of nutrient
4 management can have major limitations, as N fertilisation in excess of crop removal can lead to a
5 depletion of soil carbon reserves by enhancing microbial decomposition^[80, 81]. This approach can
6 also lead to an uneconomical fertiliser usage in the case of K that may also have an adverse effects
7 on soil quality and productivity^[26] although a range of management factors (e.g. N supply and tillage
8 system) can mask the effect of K fertilisation on crop yield^[78]. It has also been suggested that crop
9 yield and quality reductions following K fertiliser application are more likely to be related to K-Mg
10 and K-Ca antagonism in plant uptake and/or K immobilisation in the soil^[78], rather than toxicity in
11 the plant and root zone, or a depletion of the soil structure^[26].

12 In summary, it is clear from the analysis and modelling within this study that most typical
13 organic cropping systems in the UK will require nutrient inputs to maintain an N, P and K balance, .
14 It should also be remembered that most organic farms import fewer nutrients than their
15 conventional counterparts^[4, 82, 83]. Although this approach naturally leads to lower yields, and can
16 lead to lower nitrogen efficiencies within cropping systems^[4], it can also offer a useful way to
17 balance production and environmental concerns^[84, 85]. For example organic farms often require less
18 fossil energy on a per hectare or kilogram of product basis, in particular through the absence of
19 imported mineral N fertiliser^[86]. The use of grass/clover leys and manures for fertility building on
20 organic farms also contributes to greater soil organic carbon concentrations and stocks on
21 organically managed land^[87]. In addition organic methods (e.g. use of clover and other legumes to
22 supply N) can be used effectively on conventional farms to increase efficiencies and reduce the
23 environmental impacts of the agriculture sector as a whole^[39, 88, 89].

24

1 **Conclusion**

2

3 An assessment of the NDICEA model has found that it is a useful tool for UK organic farmers to
4 assess the amount of nitrogen supplied and lost through their rotations, although the model should
5 be calibrated to improve accuracy for UK conditions where measured crop N, P, K, soil-N and
6 organic matter values are available. The modelling of the N, P and K balance within organic trials
7 found that in most cases sufficient N is being supplied through biological fixation to support the
8 cropping, although leaching in higher rainfall areas and on lighter soil types may prevent the N from
9 becoming available to the crop(s). The study has also shown that careful rotation design is
10 particularly important within stockless organic systems to reduce losses and avoid the requirements
11 for external inputs as far as possible. Although adequate nitrogen balances are theoretically
12 achievable within stockless organic cropping systems, these systems are highly vulnerable to cover
13 crop failure, poor crop yields and low rates of N fixation within the fertility building period. Negative
14 P and K balances were found for most of the experimental stockless systems and the typical
15 stockless rotations modelled within this study. For phosphorus, the systems seem to be dependent
16 on imported rock phosphate for the maintenance of a small surplus or deficit. The much larger K
17 deficits could be addressed through weathering and subsequent bioavailability of mineral K stocks,
18 depending on site and management conditions. On soils with naturally low K deposits within parent
19 material, K inputs in the form of fertiliser or feed may be required to offset removal, or a reduction
20 in K demanding crops (e.g. potatoes) may be necessary.

21 N, P and K balances on organic farms are a useful method for exploring the extent to which
22 organic methods can be applied effectively to improve nutrient use efficiencies within agricultural
23 systems. It is likely that the greatest nitrogen use efficiencies can be achieved through a
24 combination of organic production methods (e.g. use of cover crops and clover to supply N)
25 combined with conventional farming practices (e.g. use of mineral fertiliser at key points in the

1 rotation to meet crop demands fully and increase yields). In addition, the need to obtain minerals
2 from sustainable sources leads to the conclusion that deriving these from suitably defined
3 wastewater treatment could close the nutrient loop for organic farms, but this would require a
4 change in international standards.

7 Acknowledgements

8
9 We would like to thank Geert-Jan van der Burgt for his help with the use of the NDICEA model and
10 the anonymous reviewers whose helpful comments greatly improved the quality of the manuscript.
11 Funding from the Department for Environment, Food and Rural Affairs and the Scottish Government
12 is gratefully acknowledged. The part time PhD for which this study forms a part is supported by an
13 education grant awarded by the Ratcliff Foundation.

15 References

- 16 1. Lampkin, N. 2002. Organic Farming. Old Pond Publishing Ltd., Ipswich.
- 17 2. Watson, C.A., Atkinson, D., Gosling, P., Jackson, L.R., and Rayns, F.W. 2002. Managing soil
18 fertility in organic farming systems. *Soil Use and Management* 18: 239-247.
- 19 3. Döring, T., Baddeley, J., Hatch, D., Marshall, A., Pearce, B., Roderick, S., Stobart, R., Storkey,
20 J., Watson, C., and Wolfe, M. 2013. Using legume-based mixtures to enhance the nitrogen
21 use efficiency and economic viability of cropping systems. Home Grown Cereals Authority:
22 Warwickshire, UK.
- 23 4. Torstensson, G., Aronsson, H., and Bergström, L. 2006. Nutrient use efficiencies and leaching
24 of organic and conventional cropping systems in Sweden. *Agronomy Journal* 98: 603-615.

- 1 5. Berry, P.M., Sylvester-Bradley, R., Philipps, L., Hatch, D.J., Cuttle, S.P., Rayns, F.W., and
2 Gosling, P. 2002. Is the productivity of organic farms restricted by the supply of available
3 nitrogen? *Soil Use and Management* 18: 248-255.
- 4 6. Dawson, J.C., Huggins, D.R., and Jones, S.S. 2008. Characterizing nitrogen use efficiency in
5 natural and agricultural ecosystems to improve the performance of cereal crops in low-input
6 and organic agricultural systems. *Field Crops Research* 107: 89-101.
- 7 7. de Ponti, T., Rijk, B., and van Ittersum, M.K. 2012. The crop yield gap between organic and
8 conventional agriculture. *Agricultural Systems* 108: 1-9.
- 9 8. Patil, R.H., Laegdsmand, M., Olesen, J.E., and Porter, J.R. 2010. Effect of soil warming and
10 rainfall patterns on soil N cycling in Northern Europe. *Agriculture, Ecosystems &*
11 *Environment* 139: 195-205.
- 12 9. Liang, X., Xu, L., Li, H., He, M., Qian, Y., Liu, J., Nie, Z., Ye, Y., and Chen, Y. 2011. Influence of N
13 fertilization rates, rainfall, and temperature on nitrate leaching from a rainfed winter wheat
14 field in Taihu watershed. *Physics and Chemistry of the Earth* 36: 395-400.
- 15 10. Berry, P., Stockdale, E., Sylvester-Bradley, R., Philipps, L., Smith, K., Lord, E., Watson, C., and
16 Fortune, S. 2003. N, P and K budgets for crop rotations on nine organic farms in the UK. *Soil*
17 *Use and Management* 19: 112-118.
- 18 11. Korsæth, A. 2012. N, P, and K budgets and changes in selected topsoil nutrients over 10
19 years in a long-term experiment with conventional and organic crop rotations. *Applied and*
20 *Environmental Soil Science Volume 2012*: 17.
- 21 12. Van der Burgt, G., Oomen, G., Habets, A., and Rossing, W. 2006. The NDICEA model, a tool to
22 improve nitrogen use efficiency in cropping systems. *Nutrient Cycling in Agroecosystems* 74:
23 275-294.

- 1 13. van der Burgt, G.J. and Rietberg, P. 2012. Enhancement of the NDICEA decision support tool
2 for reducing N losses in commercial farming. In WP2, Deliverable 2.6, EC Framework 7
3 project N-TOOLBOX, KBBE-2008-1-2-08. L.D. Louis Bolk Institute: Netherlands.
- 4 14. Cormack, W.F. 2006. Crop performance in a stockless arable organic rotation in eastern
5 England. *Biological Agriculture & Horticulture* 24: 1-20.
- 6 15. Lennartsson, M. 2000. Conversion to organic field vegetable production (Defra project
7 OF0126T). Garden Organic: Ryton, UK.
- 8 16. Welsh, J.P., Philipps, L., and Cormack, W.F. 2002. The long-term agronomic performance of
9 organic stockless rotations. In J. Powell (ed.) *Proceedings of the UK Organic Research*
10 *Conference*. Organic Centre Wales, University of Wales Aberystwyth, Aberystwyth, Wales.
11 p.47-50.
- 12 17. Taylor, B., Younie, D., Matheson, S., Coutts, M., Mayer, C., Watson, C., and Walker, R. 2006.
13 Output and sustainability of organic ley/arable crop rotations at two sites in northern
14 Scotland. *The Journal of Agricultural Science* 144: 435-447.
- 15 18. Haggard, R. and Padel, S. 1996. Conversion to organic milk production (Defra project OF0103).
16 University of Wales Aberystwyth: UK.
- 17 19. Watson, C., Baddeley, J., Edwards, A., Rees, R., Walker, R., and Topp, C. 2011. Influence of
18 ley duration on the yield and quality of the subsequent cereal crop (spring oats) in an
19 organically managed long-term crop rotation experiment. *Organic Agriculture* 1: 147-159.
- 20 20. Lampkin, N., Measures, M., and Padel, S. 2011. 2011/12 Organic Farm Management
21 Handbook. The Organic Research Centre, Elm Farm: Newbury, UK.
- 22 21. Moakes, S. and Lampkin, N. 2010. Organic farm incomes in England and Wales 2008/09
23 (Defra project OF0373). University of Wales Aberystwyth and The Organic Research Centre:
24 Aberystwyth and Newbury, UK.

- 1 22. Moakes, S. and Lampkin, N. 2011. Organic farm incomes in England and Wales 2009/10
2 (Defra project OF0373). University of Wales Aberystwyth and The Organic Research Centre:
3 Aberystwyth and Newbury, UK.
- 4 23. Moakes, S., Lampkin, N., and Gerrard, C.L. 2012. Organic farm incomes in England and Wales
5 2010/11 (Defra project OF0373). University of Wales Aberystwyth and The Organic Research
6 Centre: Aberystwyth and Newbury, UK.
- 7 24. Melillo, J.M., Aber, J.D., Linkins, A.E., Ricca, A., Fry, B., and Nadelhoffer, K.J. 1989. Carbon
8 and nitrogen dynamics along the decay continuum: plant litter to soil organic matter. *Plant
9 and Soil* 115: 189-198.
- 10 25. Johnston, A.E., Poulton, P.R., and Coleman, K. 2009. Soil organic matter: its importance in
11 sustainable agriculture and carbon dioxide fluxes. In D. Sparks (ed.) *Advances in Agronomy*.
12 Academic Press, San Diego. p.1-57.
- 13 26. Khan, S., Mulvaney, R., and Ellsworth, T. 2014. The potassium paradox: Implications for soil
14 fertility, crop production and human health. *Renewable Agriculture and Food Systems* 29: 3-
15 27.
- 16 27. Cordell, D., Drangert, J.-O., and White, S. 2009. The story of phosphorus: global food security
17 and food for thought. *Global Environmental Change* 19: 292-305.
- 18 28. Watson, C., Younie, D., Stockdale, E., Cormack, W., Cook, S., Green, M., Holland, J., Leake, A.,
19 and Welsh, J. 2000. Yields and nutrient balances in stocked and stockless organic rotations in
20 the UK. *Aspects of Applied Biology* 62: 261-268.
- 21 29. Simonsson, M., Andersson, S., Andrist-Rangel, Y., Hillier, S., Mattsson, L., and Öborn, I. 2007.
22 Potassium release and fixation as a function of fertilizer application rate and soil parent
23 material. *Geoderma* 140: 188-198.

- 1 30. Kirchmann, H., Bergström, L., Kätterer, T., Mattsson, L., and Gesslein, S. 2007. Comparison of
2 long-term organic and conventional crop–livestock systems on a previously nutrient-
3 depleted soil in Sweden. *Agronomy Journal* 99: 960-972.
- 4 31. Bergström, L., Kirchmann, H., Aronsson, H., Torstensson, G., and Mattsson, L. 2008. Use
5 efficiency and leaching of nutrients in organic and conventional cropping systems in Sweden.
6 In H. Kirchmann and L. Bergström (eds.) *Organic Crop Production—Ambitions and Limitations*.
7 Springer, Netherlands. p.143-159.
- 8 32. Aronsson, H., Torstensson, G., and Bergström, L. 2007. Leaching and crop uptake of N, P and
9 K from organic and conventional cropping systems on a clay soil. *Soil Use and Management*
10 23: 71-81.
- 11 33. Cassman, K.G., Dobermann, A., and Walters, D.T. 2002. Agroecosystems, nitrogen-use
12 efficiency, and nitrogen management. *AMBIO: A Journal of the Human Environment* 31: 132-
13 140.
- 14 34. Seufert, V., Ramankutty, N., and Foley, J.A. 2012. Comparing the yields of organic and
15 conventional agriculture. *Nature* 485: 229-232.
- 16 35. Kaffka, S. and Koepf, H.H. 1989. A case study on the nutrient regime in sustainable farming.
17 *Biological Agriculture & Horticulture* 6: 89-106.
- 18 36. ADAS. 2006. Assessing the sustainability of a stockless arable organic rotation. Defra project
19 OF0318. ADAS UK Ltd: UK.
- 20 37. Snapp, S.S., Swinton, S.M., Labarta, R., Mutch, D., Black, J.R., Leep, R., Nyiraneza, J., and
21 O'Neil, K. 2005. Evaluating cover crops for benefits, costs and performance within cropping
22 system niches. *Agronomy Journal* 97: 322-332.
- 23 38. Schmutz, U., Rayns, F., and Firth, C. 2007. Balancing fertility management and economics in
24 organic field vegetable rotations. *Journal of the Science of Food and Agriculture* 87: 2791-
25 2793.

- 1 39. Godfray, H. 2014. The challenge of feeding 9–10 billion people equitably and sustainably.
2 The Journal of Agricultural Science 152: 1-7.
- 3 40. Möller, K. and Stinner, W. 2009. Effects of different manuring systems with and without
4 biogas digestion on soil mineral nitrogen content and on gaseous nitrogen losses (ammonia,
5 nitrous oxides). European Journal of Agronomy 30: 1-16.
- 6 41. Möller, K. 2009. Influence of different manuring systems with and without biogas digestion
7 on soil organic matter and nitrogen inputs, flows and budgets in organic cropping systems.
8 Nutrient Cycling in Agroecosystems 84: 179-202.
- 9 42. Oelofse, M., Jensen, L., and Magid, J. 2013. The implications of phasing out conventional
10 nutrient supply in organic agriculture: Denmark as a case. Organic Agriculture 3: 41-55.
- 11 43. Stinner, W., Möller, K., and Leithold, G. 2008. Effects of biogas digestion of clover/grass-leys,
12 cover crops and crop residues on nitrogen cycle and crop yield in organic stockless farming
13 systems. European Journal of Agronomy 29: 125-134.
- 14 44. Di, H. and Cameron, K. 2002. Nitrate leaching in temperate agroecosystems: sources, factors
15 and mitigating strategies. Nutrient Cycling in Agroecosystems 64: 237-256.
- 16 45. Cox, T., Bender, M., Picone, C., Tassel, D.V., Holland, J., Brummer, E., Zoeller, B., Paterson, A.,
17 and Jackson, W. 2002. Breeding perennial grain crops. Critical Reviews in Plant Sciences 21:
18 59-91.
- 19 46. Pimentel, D., Cerasale, D., Stanley, R.C., Perlman, R., Newman, E.M., Brent, L.C., Mullan, A.,
20 and Chang, D.T.-I. 2012. Annual vs. perennial grain production. Agriculture, Ecosystems &
21 Environment 161: 1-9.
- 22 47. Valdivia, C., Barbieri, C., and Gold, M.A. 2012. Between forestry and farming: policy and
23 environmental implications of the barriers to agroforestry adoption. Canadian Journal of
24 Agricultural Economics/Revue canadienne d'agroeconomie 60: 155-175.

- 1 48. Webb, J., Pain, B., Bittman, S., and Morgan, J. 2010. The impacts of manure application
2 methods on emissions of ammonia, nitrous oxide and on crop response—A review.
3 Agriculture, Ecosystems & Environment 137: 39-46.
- 4 49. Smith, K., Beckwith, C., Chalmers, A., and Jackson, D. 2002. Nitrate leaching following
5 autumn and winter application of animal manures to grassland. Soil Use and Management
6 18: 428-434.
- 7 50. Watson, C., Öborn, I., Eriksen, J., and Edwards, A. 2005. Perspectives on nutrient
8 management in mixed farming systems. Soil Use and Management 21: 8.
- 9 51. Verhoeven, F., Reijs, J., and Van Der Ploeg, J. 2003. Re-balancing soil-plant-animal
10 interactions: towards reduction of nitrogen losses. NJAS-Wageningen Journal of Life Sciences
11 51: 147-164.
- 12 52. Le Gal, P.Y., Dugué, P., Faure, G., and Novak, S. 2011. How does research address the design
13 of innovative agricultural production systems at the farm level? A review. Agricultural
14 Systems 104: 714-728.
- 15 53. Edwards, A.C., Walker, R.L., Maskell, P., Watson, C.A., Rees, R.M., Stockdale, E.A., and Knox,
16 O.G. 2010. Improving bioavailability of phosphate rock for organic farming. Genetic
17 Engineering, Biofertilisation, Soil Quality and Organic Farming. Springer. p.99-117.
- 18 54. International Federation of Organic Agriculture Movements (IFOAM). 2006. IFOAM organic
19 principles. IFOAM: Bonn, Germany.
- 20 55. Nowak, B., Nesme, T., David, C., and Pellerin, S. 2013. To what extent does organic farming
21 rely on nutrient inflows from conventional farming? Environmental Research Letters 8:
22 044045.
- 23 56. Oelofse, M., Høgh-Jensen, H., Abreu, L.S., Almeida, G.F., El-Araby, A., Hui, Q.Y., and de
24 Neergaard, A. 2010. A comparative study of farm nutrient budgets and nutrient flows of

- 1 certified organic and non-organic farms in China, Brazil and Egypt. Nutrient cycling in
2 agroecosystems 87: 455-470.
- 3 57. Kirchmann, H., Kätterer, T., and Bergström, L. 2008. Nutrient supply in organic agriculture. In
4 H. Kirchmann and L. Bergström (eds.) *Organic Crop Production - Ambitions and Limitations*.
5 Springer, Netherlands. p.89-116.
- 6 58. Dekker, S.E.M., de Boer, I.J.M., Vermeij, I., Aarnink, A.J.A., and Koerkamp, P.W.G.G. 2011.
7 Ecological and economic evaluation of Dutch egg production systems. *Livestock Science* 139:
8 109-121.
- 9 59. Germer, J., Addai, S., and Sauerborn, J. 2011. Response of grain sorghum to fertilisation with
10 human urine. *Field Crops Research* 122: 234-241.
- 11 60. Karak, T. and Bhattacharyya, P. 2011. Human urine as a source of alternative natural
12 fertilizer in agriculture: A flight of fancy or an achievable reality. *Resources, conservation and
13 recycling* 55: 400-408.
- 14 61. Luske, B. and van der Kamp, J. 2009. Carbon sequestration potential of reclaimed desert soils
15 in Egypt. *Louis Bolk Institute and Soil and More International: Netherlands*.
- 16 62. Darnhofer, I. 2005. Organic farming and rural development: some evidence from Austria.
17 *Sociologia Ruralis* 45: 308-323.
- 18 63. Altieri, M., Companioni, N., Cañizares, K., Murphy, C., Rosset, P., Bourque, M., and Nicholls,
19 C. 1999. The greening of the "barrios": Urban agriculture for food security in Cuba.
20 *Agriculture and Human Values* 16: 131-140.
- 21 64. European Council. 2007. Council Regulation (EC) No 834/2007 of 28 June 2007 on organic
22 production and labelling of organic products and repealing Regulation (EEC) No 2092/91.
23 European Commission: Brussels.

- 1 65. Smith, S. 2009. Organic contaminants in sewage sludge (biosolids) and their significance for
2 agricultural recycling. *Philosophical Transactions of the Royal Society A: Mathematical,*
3 *Physical and Engineering Sciences* 367: 4005-4041.
- 4 66. Robinson, K.G., Robinson, C.H., Raup, L.A., and Markum, T.R. 2012. Public attitudes and risk
5 perception toward land application of biosolids within the south-eastern United States.
6 *Journal of Environmental Management* 98: 29-36.
- 7 67. Wilkins, R. 2008. Eco-efficient approaches to land management: a case for increased
8 integration of crop and animal production systems. *Philosophical Transactions of the Royal*
9 *Society B: Biological Sciences* 363: 517-525.
- 10 68. El-Hage Scialabba, N. and Müller-Lindenlauf, M. 2010. Organic agriculture and climate
11 change. *Renewable Agriculture and Food Systems* 25: 11.
- 12 69. Jørgensen KF and Kristensen E. 2010. Fælles strategi for udfasning af konventionel gødning
13 og halm i økologisk landbrugsproduktion [Common strategy for the phasing out of manure
14 and straw in organic agriculture]. Organic Denmark and Danish Agriculture and Food Council:
15 Copenhagen.
- 16 70. Holmqvist, J., Øgaard, A.F., Öborn, I., Edwards, A., Mattsson, L., and Sverdrup, H. 2003.
17 Application of the PROFILE model to estimate potassium release from mineral weathering in
18 Northern European agricultural soils. *European Journal of Agronomy* 20: 149-163.
- 19 71. Hinsinger, P., Dufey, J., and Jaillard, B. 1991. Biological weathering of micas in the
20 rhizosphere as related to potassium absorption by plant roots. In B. McMichael and H.
21 Persson (eds.) *Plant Roots and Their Environment*. Elsevier, Amsterdam, The Netherlands.
22 p.98-105.
- 23 72. Hoffland, E., Giesler, R., Jongmans, T., and Van Breemen, N. 2002. Increasing feldspar
24 tunneling by fungi across a north Sweden podzol chronosequence. *Ecosystems* 5: 11-22.

- 1 73. Andrist-Rangel, Y., Edwards, A., Hillier, S., and Öborn, I. 2007. Long-term K dynamics in
2 organic and conventional mixed cropping systems as related to management and soil
3 properties. *Agriculture, Ecosystems & Environment* 122: 413-426.
- 4 74. Buckman, H.O. and Brady, N.C. 1984. *The Nature and Properties of Soils*. 9th ed. Macmillan,
5 New York, USA.
- 6 75. Mengel, K., Rahmatullah, and Dou, H. 1998. Release of potassium from the silt and sand
7 fraction of loess-derived soils. *Soil Science* 163: 805-813.
- 8 76. Andrist-Rangel, Y., Hillier, S., Öborn, I., Lilly, A., Towers, W., Edwards, A.C., and Paterson, E.
9 2010. Assessing potassium reserves in northern temperate grassland soils: A perspective
10 based on quantitative mineralogical analysis and aqua-regia extractable potassium.
11 *Geoderma* 158: 303-314.
- 12 77. Loes, A.K. and Øgaard, A.F. 1997. Changes in the nutrient content of agricultural soil on
13 conversion to organic farming in relation to farm-level nutrient balances and soil contents of
14 clay and organic matter. *Acta Agriculturae Scandinavica, Section B - Soil & Plant Science* 47:
15 201-214.
- 16 78. Bar-Yosef, B., Magen, H., Johnston, A.E., and Kirkby, E.A. 2015. Potassium fertilization:
17 paradox or K management dilemma? *Renewable Agriculture and Food Systems* 30: 115-119.
- 18 79. Greenland, D.J. 1997. *The Sustainability of Rice Farming*. CAB International, Wallingford, UK.
- 19 80. Mulvaney, R.L., Khan, S.A., and Ellsworth, T.R. 2009. Synthetic Nitrogen Fertilizers Deplete
20 Soil Nitrogen: A Global Dilemma for Sustainable Cereal Production *Journal of Environmental*
21 *Quality* 38: 2295-2314.
- 22 81. Khan, S., Mulvaney, R., Ellsworth, T., and Boast, C. 2007. The myth of nitrogen fertilization
23 for soil carbon sequestration. *Journal of Environmental Quality* 36: 1821-1832.

- 1 82. Hansen, B., Kristensen, E.S., Grant, R., Høgh-Jensen, H., Simmelsgaard, S.E., and Olesen, J.E.
2 2000. Nitrogen leaching from conventional versus organic farming systems—a systems
3 modelling approach. *European Journal of Agronomy* 13: 65-82.
- 4 83. Trydeman Knudsen, M., Sillebak Kristensen, I., Berntsen, J., Molt Petersen, B., and Steen
5 Kristensen, E. 2006. Estimated N leaching losses for organic and conventional farming in
6 Denmark. *The Journal of Agricultural Science* 144: 135-149.
- 7 84. Gomiero, T., Pimentel, D., and Paoletti, M.G. 2011. Environmental impact of different
8 agricultural management practices: conventional vs. organic agriculture. *Critical Reviews in*
9 *Plant Sciences* 30: 95-124.
- 10 85. Francis, C. and Porter, P. 2011. Ecology in sustainable agriculture practices and systems.
11 *Critical Reviews in Plant Sciences* 30: 64-73.
- 12 86. Smith, L.G., Williams, A.G., and Pearce, B.D. 2015. The energy efficiency of organic
13 agriculture: A review. *Renewable Agriculture and Food Systems* 30: 280-301.
- 14 87. Gattinger, A., Muller, A., Haeni, M., Skinner, C., Fliessbach, A., Buchmann, N., Mäder, P.,
15 Stolze, M., Smith, P., and Scialabba, N.E.-H. 2012. Enhanced top soil carbon stocks under
16 organic farming. *Proceedings of the National Academy of Sciences* 109: 18226-18231.
- 17 88. Pretty, J.N., Noble, A.D., Bossio, D., Dixon, J., Hine, R.E., Penning de Vries, F.W.T., and
18 Morison, J.I.L. 2005. Resource-conserving agriculture increases yields in developing
19 countries. *Environmental Science & Technology* 40: 1114-1119.
- 20 89. Gaudin, A., Westra, S., Loucks, C.E., Janovicek, K., Martin, R.C., and Deen, W. 2013.
21 Improving resilience of northern field crop systems using inter-seeded red clover: A review.
22 *Agronomy* 3: 148-180.

23

24

1

2 Captions for Tables and Figures:

3

4 **Figure 1.** Approximate location and site parameters for each of the long-term organic trial sites. OM
5 = organic matter content of the soil (% loss on ignition). Rainfall amounts are mean values over the
6 course of the trial(s).

7 **Table 1.** Crop Rotations used at each of the experimental sites (C = Carrots, G/C = Grass White

8 Clover, P = Potatoes, RC = Red clover, SBA = Spring barley, SB = Spring beans, SO = Spring oats, S =
9 Swede, SW = Spring wheat, WB = Winter beans, WO = Winter oats, WW = Winter wheat).

10

11 **Table 2:** Typical organic rotations assessed within this study (G/WC = Grass/ white clover,

12 RC/G = Red clover/grass, SO - Spring oats, SW = Spring wheat, SB = Spring beans, WO =

13 Winter oats, WW = Winter wheat, WR = Winter rye, P = Potatoes, PE = Peas)

14

15 **Figure 2.** Comparison of the land use by crop type for the typical rotations used within this study to
16 data for 30 'Cropping Farms' collected with the FBS-based Organic Farm Incomes Reports (2010-
17 2012). Error bars = standard error.

18 **Table 3.** Comparison of the error found in calculating soil N (kg/N/ha) produced by

19 uncalibrated and calibrated runs of NDICEA using data collected for the rotations applied at
20 each experimental site (RMSE = Root Mean Square Error across all measurements, n =

21 number of soil mineral N samples used for calibration of the model).

22

1 **Table 4.** Nutrient balance of stockless organic rotations/trial sites expressed in kg/ha/year

2 **Table 5.** Nutrient balance of stocked organic rotations/trial sites expressed in kg/ha/yr.

3 **Table 6.** Nutrient balance of typical organic rotations/trial sites expressed in kg/ha/yr. se = standard
4 error (se is only calculated for N, P and K balances are unaffected by the site conditions).

5

6

7

8

9

10

11

12

13

14

15

16

SRUC Woodside, Elgin,
Annual rainfall: 641 mm
OM: 6%
Soil type: Light sandy loam
pH: 6.2

SRUC Tulloch, Craibstone,
Aberdeen
Annual rainfall: 902mm
OM: 9%
Soil type: Sandy loam
pH: 5.7

Garden Organic/Warwick
Univ. Hunts Mill site,
Warwickshire
Annual rainfall: 548mm
OM: 1.8%
Soil type: Light sandy loam
pH: 6.3

ADAS Terrington, Norfolk
Annual rainfall: 622mm
OM: 2.5%
Soil type: Silty clay loam
pH: 7.5

IBERS Ty Gwyn Organic
Dairy Unit
Annual rainfall: 884 mm
OM: 6.5%
Soil type: Silty clay loam
pH: 5.8

The Organic Research Centre,
Elm Farm, Berkshire
Annual rainfall: 712 mm
OM: 3.2%
Soil type: Clay loam
pH: 6.5






Table 2: Typical organic rotations assessed within this study (G/WC = Grass/ white clover, RC/G = Red clover/grass, SO = Spring oats, SW = Spring wheat, SB = Spring beans, WO = Winter oats, WW = Winter wheat, WR = Winter rye, P = Potatoes, PE = Peas)

Rotation	Course									
	1	2	3	4	5	6	7	8	9	10
Stocked 'complex'	G/WC	G/WC	G/WC	WW	WO	RC/G	RC/G	P	SB	SW
Stocked 'simple'	RC/G	RC/G	WW	P	WW	WR				
Stockless 'complex'	RC/G	RC/G	P	WO	SB	SW				
Stockless 'simple'	RC/G	WW	PE	SO						

Table 3: Comparison of the error found in calculating soil N (kg/N/ha) produced by uncalibrated and calibrated runs of NDICEA using data collected for the rotations applied at each experimental site (RMSE = Root Mean Square Error across all measurements, n = number of soil mineral N samples used for calibration of the model)

Site / experiment	n	RMSE uncalibrated model	RMSE calibrated model
EFRC A	7	48.8	16.6
EFRC B	7	57.6	48.7
EFRC C	7	30.4	21.5
ADAS Terrington	10	10.9	6.3
Warwick, Hunts Mill Area 1 with FYM	12	47.7	44.1
Warwick, Hunts Mill Area 6 with FYM	12	22.3	19.3
Warwick, Hunts Mill Area 1 no FYM	12	41.0	38.8
Warwick, Hunts Mill Area 6 no FYM	12	23.7	18.6
SRUC Woodside W37	30	12.8	11.6
SRUC Woodside W50	30	14.1	11.8
SRUC Tulloch T50	5	22.0	18.9
SRUC Tulloch T67	5	13.4	6.0
Ty Gwyn	2	8.3	7.5

Table 4: Nutrient balance of stockless organic rotations/trial sites expressed in kg/ha/year

	EFRC A			EFRC B			EFRC C			ADAS			HuntsMill Area 1			HuntsMill Area 6		
	N	P	K	N	P	K	N	P	K	N	P	K	N	P	K	N	P	K
	Fertiliser applied		8			8			8			8		8	5	18		
Deposition	20		4	20		4	20		4	30	1	5	20		3	20		3
Biological fixation	44			59			59			84			42			21		
Total supply	64	8	4	79	8	4	79	8	4	114	9	5	70	5	22	41	0	3
Volatilisation	0			0			0						0			0		
Denitrification	47			54			44			26			24			7		
Leaching	57			57			52			28			18			15		
Product Removal	36	8	6	63	17	32	55	14	26	77	15	45	35	9	55	33	10	62
Total loss	140	8	6	174	17	32	151	14	26	131	15	45	77	9	55	55	10	62
Nutrient balance	-76	0	-2	-95	-9	-28	-72	-6	-22	-17	-6	-40	-7	-4	-33	-14	-10	-59
Change in soil organic N	-93			-100			-60			0			-1			-13		
Change in soil mineral N	17			5			-12			-17			-6			-1		

Table 5: Nutrient balance of stocked organic rotations/trial sites expressed in kg/ha/yr

	Tulloch T50			Tulloch T67			Woodside W50			Woodside W37			Hunts Mill Area 1			Hunts Mill Area 6		
	N	P	K	N	P	K	N	P	K	N	P	K	N	P	K	N	P	K
Fertiliser applied	63	17	63	83	22	81	63	17	62	38	10	37	48	13	53	62	16	61
Deposition	12	1	7	12	1	7	12	1	7	12	1	7	20		4	20		3
Biological fixation	57			109			112	0	0	60			25			9		
Total supply	132	18	69	204	24	88	187	18	69	110	11	44	93	13	57	91	16	65
Volatilisation	7			8			6			5			5			5		
Denitrification	3			3			12			11			15			14		
Leaching	26			30			49			45			29			29		
Product Removal	59	17	96	57	36	188	78	19	92	61	17	79	43	10	59	42	11	71
Total loss	95	17	96	98	36	188	145	19	92	122	17	79	92	10	59	90	11	71
Nutrient balance	37	1	-27	106	-12	-100	42	0	-23	-12	-5	-35	1	4	-2	1	4	-6
Change in organic N	32			101			43			6			1			1		
Change in mineral N	5			5			1			-18			0			0		

Table 6: Nutrient balance of typical organic rotations/trial sites expressed in kg/ha/yr. se = standard error (se is only calculated for N, P and K balances are unaffected by the site conditions)

	Stocked complex				Stocked simple				Stockless complex				Stockless simple			
	N	se (+/-)	P	K	N	se (+/-)	P	K	N	se (+/-)	P	K	N	se (+/-)	P	K
Fertiliser applied	32	0	12	23	26	0	14	21			8				9	
Deposition	17	3		5	17	2		5	17	3		5	16	2		5
Biological fixation	142	9			117	9			93	10			130	11		
Total supply	190	9	12	28	160	10	14	26	109	10	8	5	146	11	9	5
Volatilisation	2	0			2	0			0	0			0	0		
Denitrification	21	2			17	2			31	3			26	2		
Leaching	35	9			39	9			50	10			57	11		
Product Removal	79	0	13	37	72	0	14	42	49	0	10	37	55	0	10	16
Total loss	137	8.7	13	37	130	10	14	42	130	10	10	37	138	11	10	16
Nutrient balance	53	7	-1	-9	30	9	0	-16	-21	3	-2	-32	8	3	-1	-11
Change in organic N	51	8			28	8			-3	10			20	9		
Change in mineral N	2	8			2	7			-18	9			-12	7		