

Scotland's Rural College

Genetic dissection of photoperiod response based on GWAS of pre-anthesis phase duration in spring barley

Alqudah, Ahmad M; Sharma, Rajiv; Pasam, Raj K; Graner, Andreas; Kilian, Benjamin; Schnurbusch, Thorsten

Published in:
PLoS ONE

DOI:
[10.1371/journal.pone.0113120](https://doi.org/10.1371/journal.pone.0113120)

Print publication: 24/11/2014

Document Version
Publisher's PDF, also known as Version of record

[Link to publication](#)

Citation for published version (APA):

Alqudah, A. M., Sharma, R., Pasam, R. K., Graner, A., Kilian, B., & Schnurbusch, T. (2014). Genetic dissection of photoperiod response based on GWAS of pre-anthesis phase duration in spring barley. *PLoS ONE*, 9(11), Article e113120. <https://doi.org/10.1371/journal.pone.0113120>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

CORRECTION

Correction: Genetic Dissection of Photoperiod Response Based on GWAS of Pre-Anthesis Phase Duration in Spring Barley

The PLOS ONE Staff

There are errors in the formatting of [Table 1](#). Please see the correct [Table 1](#) here.

The following errors appear in the body of the article.

In the second paragraph of the Introduction, *PSEUDO-RESPONSE REGULATOR* (*HvPRR37*) should be *PSEUDO-RESPONSE REGULATOR 37* (*HvPRR37*).

There are errors in the sixth sentence of the Materials and Methods subsection titled “Phenotyping.” The correct sentence is: To avoid any mineral deficiency each pot was additionally fertilized with 1.5 gram of solid fertilizer (that constitute minerals 17:11:10/N: P: K).

There are errors in the fourth sentence of the Results subsection titled “Identification of marker-trait association within the *ppd-H1*-carrying group.” The correct sentence is: However, *HvCO1* resides in a chromosomal region, which physically contains three other *CO*-like family members (*HvCO12/ HvCO13/HvM* and *HvCO15*) and the circadian clock-related genes, *HvLHY/HvCCA1*.

There are errors in the twelfth sentence of the fourth paragraph in the Discussion subsection titled “Heading time genetic network models.” The correct sentence is: Similar conclusions can be drawn from gene clusters on chromosome 4H, which besides *HvCO16* also includes *HvPRR59*, *HvPhyB* and *HvPrr73* (see 4H; 51.1–51.4 cM), 5HS (including *HvC*)3, *HvTFL1*, *HvCMF13*; 43.7–51.6 cM) and 1HS (*HvCMF10*; 41.1–48.2 cM).

There are errors in the fourteenth sentence of the fourth paragraph in the Discussion subsection titled “Heading time genetic network models.” The correct sentence is: Another group-specific effect was found for SNPs around the *FT*-like homolog *HvFT2* and the circadian clock-related gene *HvGI* (3HS; 45.8–52.0 cM).

There are errors in the third sentence of the sixth paragraph in the Discussion subsection titled “Heading time genetic network models.” The correct sentence is: Although its genetic position strongly suggests that the strongest associations reside very close to *HvCO1* (supported by 13 SNP markers; see Figure 9), we cannot completely rule out that linkage with other *CO* genes in this region like *HvCO12/ HvCO13/HvM* and *HvCO15* (Figure 7) contribute to this effect.



CrossMark
click for updates

 OPEN ACCESS

Citation: The PLOS ONE Staff (2015) Correction: Genetic Dissection of Photoperiod Response Based on GWAS of Pre-Anthesis Phase Duration in Spring Barley. PLoS ONE 10(4): e0123748. doi:10.1371/journal.pone.0123748

Published: April 7, 2015

Copyright: © 2015 The PLOS ONE Staff. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Table 1. Spike row-type and origins of spring barley accessions with photoperiod-sensitive (*Ppd-H1*) and reduced photoperiod sensitivity (*ppd-H1*).

Origin [‡]	Photoperiod-sensitive(<i>Ppd-H1</i>)		Reduced photoperiod sensitivity (<i>ppd-H1</i>)		Total
	Two-rowed	Six-rowed	Two-rowed	Six-rowed	
WANA	12	21	11	1	45
EU	10	6	80	12	108
EA	0	28	2	6	36
AM	6	12	4	7	29
Total	28	67	97	26	218
	95		123		

[‡] WANA: West Asia and North Africa, EU: Europe, EA: East Asia, AM, Americas.

doi:10.1371/journal.pone.0123748.t001

Reference

1. Alqudah AM, Sharma R, Pasam RK, Graner A, Kilian B, Schnurbusch T (2014) Genetic Dissection of Photoperiod Response Based on GWAS of Pre-Anthesis Phase Duration in Spring Barley. PLoS ONE 9(11): e1113120. doi: [10.1371/journal.pone.0113120](https://doi.org/10.1371/journal.pone.0113120) PMID: [25420105](https://pubmed.ncbi.nlm.nih.gov/25420105/)