

Scotland's Rural College

Factors affecting ewe longevity on sheep farms in three European countries

McLaren, A; McHugh, Noirin; Lambe, NR; Pabiou, Thierry; Wall, Eamon; Boman, Inger Anne

Published in:
Small Ruminant Research

DOI:
[10.1016/j.smallrumres.2020.106145](https://doi.org/10.1016/j.smallrumres.2020.106145)

Print publication: 01/08/2020

Document Version
Peer reviewed version

[Link to publication](#)

Citation for published version (APA):
McLaren, A., McHugh, N., Lambe, NR., Pabiou, T., Wall, E., & Boman, I. A. (2020). Factors affecting ewe longevity on sheep farms in three European countries. *Small Ruminant Research*, 189, [106145].
<https://doi.org/10.1016/j.smallrumres.2020.106145>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

1 Factors affecting ewe longevity on sheep farms in three European countries

2
3 A. McLaren^a, N. McHugh^b, N.R. Lambe^a, T. Pabiou^c, E. Wall^c and I.A. Boman^d

4
5 ^a*Scotland's Rural College, Hill & Mountain Research Centre, Kirkton Farm, Crianlarich,*
6 *FK20 8RU, UK.*

7 ^b*Teagasc, Animal and Grassland Research and Innovation Centre, Moorepark, Fermoy,*
8 *County Cork, Ireland.*

9 ^c*Sheep Ireland, Highfield House, Shinagh, Bandon, County Cork, Ireland.*

10 ^d*The Norwegian Association of Sheep and Goat Breeders, PO Box 104, N-1431, Ås, Norway.*

11 12 **Abstract**

13 The ability to identify ewes that can outperform their contemporaries, in terms of how long they remain
14 productive in the flock, will help towards improving flock efficiency and profitability. The main
15 objectives of this study were to: 1) identify the main reasons for mortality or culling within diverse
16 sheep production systems in Ireland, Norway and UK; 2) investigate the influence of early life factors
17 on ewe longevity within each of these systems; and 3) determine whether common approaches or
18 recommendations could be employed to improve ewe longevity. The main reasons for mortality or
19 culling were, in addition to old age, mastitis (Irish and Norwegian sheep) and tooth loss (UK hill sheep).
20 In each country, there were significant differences in age at last lambing due to the year the ewe was
21 born (but in no consistent pattern), and due to her flock of birth ($P < 0.05$). From the Norwegian data,
22 there was some indication ewes from younger dams lambed for the last time at a younger age, however,
23 this trend was not seen in the Irish or UK data. Ewes born as singletons, in the Irish data, lambed for
24 the last time at an older age than those that had been born in larger litters, although this was not observed
25 in the other data sets. Age at first lambing and some breed proportions (proportion of Texel and Suffolk
26 particularly) of the animal (both not fitted in the Norwegian or UK analyses) were found to have a
27 highly significant ($P < 0.0001$) effect on age at last lambing in the Irish analyses. The results suggest that
28 longevity is influenced by a range of different factors and the early life predictors investigated could

29 not be used to provide consistent recommendations across countries, production systems and breeds
30 that would influence ewe longevity. One common definition or solution to select ewes for longer
31 productive life in divergent sheep flocks may not be appropriate.

32 **Keywords (3-6 words): longevity, culling, ewe, sheep**

33 **1. Introduction**

34 Ewe longevity has long been regarded by sheep breeders across Europe, and internationally, as an
35 economically important trait in a breeding ewe flock, due to the potential to reduce culling rates and
36 female replacement costs. The ability to identify ewes that are able to outperform their contemporaries,
37 in terms of how long they remain productive in the flock, will help towards improving flock efficiency
38 and profitability. Fewer unproductive animals on farm will also reduce greenhouse gases emissions per
39 kilogram of lamb produced, providing an additional environmental benefit (Jones et al., 2013).

40 Production systems for meat sheep vary considerably across Europe, due to differences in
41 environmental and climatic conditions, breed types, subsidy systems and markets, amongst other
42 factors. Therefore, different pressures are exerted on breeding ewes, and their flock-masters, which can
43 influence mortality rates and culling decisions. As a result, research outputs and information on ewe
44 longevity from one country, breed or system may not be relevant or applicable in others.

45 Ewe longevity has been investigated across several studies in different countries. However, definitions
46 of longevity vary across studies, countries and systems. Nevertheless, there may be common ways of
47 defining ewe longevity as a trait that would allow comparisons across populations in different systems
48 with differing levels of recording. One of the most common traits relating to ewe longevity that has
49 been investigated in the literature is productive life, determined by the number of years the ewe remains
50 in the flock (Borg et al., 2009). Unlike the data available from research flocks, which may record
51 cull/death dates and reasons, data collected from commercially recorded flocks seldom provides
52 sufficient detail to assess different causes of ewes leaving the flock. However, the age of the ewe at her
53 last recorded lambing event is a longevity trait that could potentially be investigated across research

54 and commercial data sets from different sources, in an attempt to better understand commonalities and
55 assess potential to share knowledge across sheep populations.

56 Results from studies that have investigated the genetic control of ewe longevity suggest that ewe
57 productive life is lowly heritable, with heritability estimates ranging from 0.03-0.13 (Borg et al., 2009;
58 Brash et al., 1994; Conington et al., 2001; Lambe et al., 2008; Lee et al., 2015; McLaren et al., 2017;
59 Zishiri et al., 2013). In addition to estimating variance components for ewe longevity, Conington et al.
60 (2001) also produced some of the first economic weightings for longevity, relating to different types of
61 farming system in the UK. However, there are few international examples of longevity traits included
62 in national breeding indices, or as stand-alone breeding values, using commercially recorded data
63 available from performance recording schemes (Ireland and New Zealand: Santos et al., 2015; UK
64 Lleyns: McLaren et al., 2017), and breeding values for ewe longevity are not yet widely available in
65 most countries.

66 Before attempting to move to a genetic solution for increasing ewe longevity across countries, it is
67 important to understand the main causes of ewe mortality or culling (voluntary or involuntary) within
68 systems, and to attempt to identify early life predictors. This information may help us to understand
69 whether the same trait is being compared across populations and whether common solutions to extend
70 ewe longevity are possible or appropriate.

71 The objectives of this study were to: 1) identify the main reasons for mortality or culling within diverse
72 sheep production systems in Ireland, Norway and UK; 2) investigate the influence of early life factors
73 on ewe longevity within each of these systems; and 3) determine whether common approaches or
74 recommendations could be employed to increase ewe longevity across different European sheep
75 production systems.

76 **2. Materials and methods**

77 *2.1. Recording mortality and culling reasons*

78 Currently, the national sheep performance recording systems in both Norway and Ireland provide
79 farmers the opportunity to record reasons for death or culling. This has not been the case in national

80 UK breeding programmes, but detailed information is available from research flocks, which provide an
81 indication of the common reasons for animals leaving similar commercial-type flocks.

82 The Norwegian Sheep Recording System (SRS) was established in the 1950's and is voluntary. It is
83 now hosted by Animalia – Norwegian Meat and Poultry Research Centre (Animalia, 2018). The
84 minimum requirements for recording are: animals joining or leaving the flock (dead or alive); lambs
85 born; and an autumn weight on all lambs. Slaughter data on individuals are automatically transferred
86 from the abattoirs. It is also compulsory for members of ram circles to be members of the SRS and, in
87 recent years, an increasing number of traits (i.e. mating information, birth weights, teat size and reasons
88 for a ewe leaving the flock) are now required to be recorded. Approximately 40% of all sheep flocks,
89 and 52% of all ewes, in Norway report to SRS. The most commonly recorded breed is the Norwegian
90 White Sheep (NWS), making up 69% of the ewes in the SRS (Animalia, 2018).

91 Sheep Ireland, and the Sheep Ireland database, were formed in 2009. Performance data, from just over
92 700 commercial and pedigree ram breeding flocks throughout Ireland, are currently recorded through
93 the LambPlus performance recording service (<https://www.sheep.ie>). Growth, lambing and litter size
94 are the main elements of the genetic evaluation and sheep breeders can use the Terminal and
95 Replacement overall indexes to implement breeding improvement (Pabiou et al., 2014). Traits are
96 recorded across a wide range of pure- and cross-bred animals, but Texel, Suffolk, Charollais, Belclare,
97 and Vendeen sired animals provide the bulk of those performance recorded. Ewe longevity is not
98 currently a trait included in the Sheep Ireland genetic indexes but a major focus is now being put on
99 building a large dataset of accurate culling and death reasons from all performance recording flocks.

100 For the current study, death and cull data for the UK analyses were from performance recorded Scottish
101 Blackface ewes, based on SRUC's hill and upland research flocks, one in the Pentland Hills, near
102 Edinburgh, the other in the Western Highlands, near Crianlarich. These flocks have been performance
103 recorded since 1999 and the data used to develop a multi-trait selection index suitable for hill sheep
104 breeds (Conington et al., 2001). The selection index created has gone on to form the basis of the
105 commercial Hill-2 Index, which is currently available to commercial UK hill sheep breeders through

106 the Signet Sheepbreeder Service (<http://www.signetfbc.co.uk/>). Although longevity was included as a
107 trait in the selection index developed by Conington et al. (2001), it is not currently included in the Hill-
108 2 index. However, details were available for every death or cull across both research flocks and, as these
109 flocks are recorded as part of the Signet Sheepbreeder Service, these were deemed to be representative
110 of the data that would be submitted from similar UK hill-type flocks.

111 Data from these three sources were assessed to identify the main reasons for ewes, over one year of age,
112 leaving the flock. Details of the full range of options provided for recording death or cull reasons are
113 given in the supplementary material. The data available from each country included: 113,319 records
114 from Norwegian White Sheep ewes, reared on SRS ram circle flocks, recorded between 2011 and 2016;
115 23,880 records from pure- and cross-bred ewes reared on commercial Irish flocks, recorded between
116 2010 and 2016; and 3,224 records from Scottish Blackface hill ewes recorded on SRUC research flocks
117 between 2003 and 2016.

118 *2.2. Early life factors affecting ewe longevity*

119 The definition of ewe longevity used in these analyses was the age (in years) at the last recorded lambing
120 event. The data available from each country included: Norway, n=55,703 records, from Norwegian
121 White Sheep ewes born between 2001 and 2012; Ireland, n =14,213 records, from ewes born between
122 2004 and 2011; and UK, n=3,173 records, from Scottish Blackface ewes born between 2002 and 2011.

123 *2.2.1. Statistical analyses*

124 A number of different fixed effects and covariates were tested using generalized linear models in
125 Genstat (VSN International, 2013) or SAS® software (SAS Institute, 2013). Stepwise regression was
126 then used to identify a suitable final model for the analyses for each country's data set.

127 The common effects used across all countries included ewe birth year (Norway 12 levels, 2001-2012;
128 Ireland 8 levels, 2004-2011; UK 10 levels, 2002-2011), flock of birth (Norway 415 flocks; Ireland 368
129 flocks; UK 2 flocks), the age of the ewe's dam (Norway 6 levels, 1 to ≥ 6 ; Ireland 7 levels, 1 to ≥ 7 ; UK
130 5 levels, 2 to ≥ 6), litter size of the ewe when she was born, her birth weight and her live weights recorded

131 between 6-8 weeks old and between 14-20 weeks old (as well as squared live weights in the Norwegian
132 and UK analyses, to test if live weight had a non-linear quadratic effect on age at last lambing). All
133 weights were fitted as covariates. In the UK and Irish data, live weights recorded between 6-8 weeks
134 old were corrected to 56 and 40 days old respectively, as these are the recommended ages for weighing
135 lambs within these breeding programmes. Weights were corrected to a fixed age by linear regression of
136 live weight on age, using the predicted mean weight at the specified age plus the residual value for each
137 individual lamb. Similarly, for those recorded between 14-20 weeks old, the live weights were corrected
138 to 111 and 100 days old respectively. The Norwegian live weight data was live weight gain from birth
139 to the respective weighing event, corrected to 42 and 140 days old. An interaction term between ewe
140 birth year and flock were fitted in the Norwegian and UK models, but was dropped from the Irish
141 analyses due to non-significance. The Irish model also included age of the ewe at her first lambing event
142 (>8 and <18 months of age, or ≥ 18 and <28 months of age), and the breed composition for each animal
143 for the main recorded breeds (Texel, Suffolk, Charollais, Belclare or Vendeen), since the data contained
144 both pure- and cross-bred animals. In the Norwegian and UK data sets, ewe age at first lambing was
145 not fitted in the models, for the reasons explained below.

146 The final model used across the datasets in each country was therefore:

147 Age at last lambing = Ewe birth year + Flock of birth + Dam age + Ewe birth weight + Birth litter size
148 + Live weight (6-8 weeks old) + [Live weight (6-8 weeks old)]^{2*} + Live weight (14-20 weeks old) +
149 [Live weight (14-20 weeks old)]^{2*} + (Ewe birth year x Flock of birth)* (+ *age at first lambing* + *breed*
150 *proportion Texel* + *breed proportion Suffolk* + *breed proportion Belclare* + *breed proportion*
151 *Charollais* + *breed proportion Vendeen*)**

152 *Fitted in the UK and Norwegian analyses only

153 ** Fitted in the Irish analysis only

154 An overall summary of the data available for the effects fitted in the models, are shown in Table 1.
155 When an effect was identified as a significant source of variation (and included less than 10 factors),
156 least-squares means (LSMs) and standard errors of the means (SEMs) were estimated. Differences

157 between the means were then tested using a t-test to identify significant differences between factors,
158 within country.

159 Other important points to note are that in the Norwegian and Irish analyses, any ewes that missed a
160 lambing event during their lifetime (i.e. had a lambing interval over 600 days at any point in their life)
161 were removed completely from the analyses. Also, all animals in the Norwegian dataset had to have a
162 lambing event at 1 year old to be included in the analyses, whereas the Irish data included ewes that
163 lambed for the first time as a 1 or 2 year old. The UK data did not include any ewes that lambed as a 1
164 year old (which is uncommon in UK hill flocks), and did include ewes that missed one lambing year
165 (either through barrenness or lamb loss).

166

167 **3. Results**

168 *3.1 Reasons for mortality or culling*

169 The main reasons submitted for ewes leaving the flock (those representing over 5% of records in each
170 data set), in each country, are given in Table 2. The proportion of ewes culled due to old age ranged
171 from 12.4% to 23.5%. A common problem identified in both Norway and Ireland was mastitis (19.9%
172 and 13.5% of ewes culled, respectively), whereas in the UK data set, the highest proportion of animals
173 were culled due to problems associated with their teeth (38.9%). In addition to mastitis, a relatively
174 high proportion of animals in the Norwegian data set were culled for udder/teat damage or poor udder
175 conformation (16.9%), with a smaller proportion culled due to udder-related problems in the Irish data
176 (5.7%). Although not shown in Table 2, mastitis only accounted for 3.4% of the reasons provided in the
177 UK data. Poor body condition contributed to the most common culling reasons in the Irish (5.7%) and
178 UK (6.8%) data sets, whilst reproductive issues accounted for 6.3% and 5.4% of culls in the Norwegian
179 and UK data sets, respectively. Ewes that were known to have died, but no reason was attributed, made
180 up 19.9% and 3.8% of the reasons in Ireland and UK data sets, respectively. Additionally, ewes
181 identified as being sold straight to an abattoir for slaughter made up 15.9% of the Irish data. Ewes
182 identified as being missing, and therefore presumed dead, accounted for 6.8% of the UK data. The
183 remaining reasons, not listed in Table 2 but given in the supplementary material, when combined, made
184 up 44.5% (n=19 reasons), 12.0% (n=7 reasons) and 18.6% (n=9 reasons) of the Norwegian, Irish and
185 UK data, respectively. These remaining reasons included lambing associated problems (e.g. prolapses,
186 abortions, poor mothering ability and lambing difficulties), physical reasons (e.g. bad feet or legs), poor
187 genetics (estimated breeding values and/or overall index values) and health related issues (other than
188 mastitis). Norway also had a number of records associated with overall management reasons (e.g. food
189 shortages) and predators (of which there were 8 different sub-reasons, e.g. taken by a wolverine, wolf
190 or lynx).

191

192

193

194 *3.2 Early life factors affecting age at last lambing*

195 The summary statistics for age at last lambing in the data sets from Norway, Ireland and the UK are
196 given in Table 3.

197 The frequency of the data available from each country, for age at last lambing, is shown in Fig 1. The
198 distributions vary across countries. In Norway, last lambing events were fairly evenly spread across
199 ages – ranging from 12-20% from 1 to 6 years old. Just over 20% of the records available were for ewes
200 who's first lambing was also their last lambing (at 1 years old), and 18.7% of the records were from
201 ewes that were 6 years old. In the Irish data, the distribution appeared to be more normal in shape across
202 ages from 1-7 years old, with 6.4% and 12.4% of the age at last lambing records for ewes aged 1 or 2
203 years old and a peak of around 21% at 4 and 5 years old. However, it should be noted that the average
204 age at first lambing was 1.88, therefore not all ewes had the opportunity to lamb as a 1 year old. In the
205 UK data, just over 20% of the age at last lambing data records were from 2 and 3 year old ewes
206 (combined). The most common age for ewes to leave the flock was at 5 years old (42.6% of the UK
207 data records).

208

209 **Fig. 1** here

210

211 Of the early life predictors tested in the models, only the birth year of the ewe and her flock of birth
212 significantly ($P<0.05$) affected age at last lambing across all three analyses (Table 4). The age of the
213 ewe's dam and the interaction between birth year of the ewe and the flock of birth had significant
214 ($P<0.05$) effects on age at last lambing in both the Norwegian and UK analyses. Other significant effects
215 ($P<0.05$) included birth litter size of the ewe in the Norwegian and Irish analyses and the weight of the
216 ewe recorded between 6-8 weeks and at 14-20 weeks (squared) of age in the Irish and UK analyses,
217 respectively. Age at first lambing and the proportion of Texel, Suffolk and Belclare breed in the animal
218 (not fitted in the Norwegian or UK analyses) were found to have a highly significant ($P<0.001$) effect
219 on age at last lambing in the Irish analyses. The amount of variance accounted for by the combined

220 terms in the models (R^2) and the residual means standard error (RMSE) values are also given in Table
221 4.

222
223 In order to investigate the direction and magnitude of the significant effects on age at last lambing, for
224 informative early life predictors, least-squares means (LSMs) and standard errors of the means (SEMs)
225 were estimated. For the age of the ewe's dam, no clear relationships were observed (Fig. 2). In the
226 Norwegian data, ewes with a dam aged 1 or 2 years old had a significantly lower age at last lambing
227 compared with those with a dam aged 3 years or older. However, in the UK data, ewes with a 3 year
228 old dam had a significantly lower age at last lambing ($P < 0.05$) when compared with those with a 2 or 4
229 year old dam, whilst other dam age groups did not differ significantly. There were no significant
230 differences between dam ages observed in the Irish results.

231 The effect of birth litter size (Fig. 2) was most evident in the Irish data, with significant ($P < 0.05$) LSM
232 differences observed between those born as singles, twins and triplets. Those born as a single had their
233 last lamb, on average, 6 months later than those reared as triplets (5.5 years for singles and 5.0 years for
234 triplets). The average age at last lambing for those reared as quadruplets was not significantly ($P > 0.05$)
235 different from those reared as twins or triplets. In the Norwegian data, there was a significant increase
236 in the average age at last lambing ($P < 0.05$) observed in ewes that were born as twins, compared with
237 those born as triplets, but neither of these litter sizes were significantly ($P > 0.05$) different to singles or
238 quadruplets. There was no significant difference between ewes born into different litter sizes ($P > 0.05$) in
239 the UK data. There was a tendency for triplet-born ewes to have their last lamb at an earlier age, but
240 low numbers in this category in the data set led to a high standard error associated with the mean.

241 Age at first lambing was only fitted in the Irish analyses. The LSM results from these analyses found
242 that the mean age at last lambing was significantly lower ($P < 0.05$) for ewes that had lambed for the first
243 time as a 1 year old compared with those that lambed for the first time at 2 years old. The mean age at
244 last lambing was 4.6 ± 0.19 and 5.3 ± 0.18 years, respectively.

245

Fig. 2 Here

237 **4. Discussion**

238 *4.1. Common reasons for ewe mortality/culling*

239 Breeding schemes across all three countries and sheep systems studied have seen improvements in traits associated with
240 growth, reproductive performance and carcass characteristics (Eikje et al., 2008; Lambe et al., 2014; Santos et al., 2015).
241 Improving ewe longevity remains another important goal, due to economic, environmental and welfare related benefits,
242 but progress has proven to be difficult due to the trait being lowly heritable, expressed later in life, and influenced by a
243 range of different factors (Lee et al., 2015). In an attempt to gather more detailed information as to why ewes are leaving
244 the flock at certain ages, both performance recording schemes in Norway and Ireland have, in recent years, provided
245 farmers the opportunity to submit animal fate information. Disposal reasons have also recently been introduced in the
246 UK national recording schemes, but it will take a few years for enough data to be collected before it can be analysed in
247 detail. In the meantime, data available from research flocks provides an indication as to why and when ewes are leaving
248 extensive-type hill farms similar to those being recorded in the national breeding scheme.

249 The most common reasons for mortality/culling, submitted by flock-masters, varied across all three countries. In
250 Norway, the most common reason was mastitis, closely followed by other udder associated problems, which included
251 issues such as poor conformation or damage to the udder or teats. Mastitis and udder problems also featured strongly
252 for ewes in the Irish data, but much less so for ewes in the UK data from the hill flocks (<5%). The high proportion of
253 Norwegian White Sheep in Norway culled for mastitis (plus other udder problems) may be influenced by the fact they
254 are a prolific breed (Jakobsen et al., 2017). Large litters, plus the emphasis on lamb growth and carcass characteristics,
255 may increase suckling frequency and increase the potential for damage to the udder. Indeed, Larsgard and Vaabenoe
256 (1993) and Waage and Vatn (2008) observed, when previously assessing mastitis in Norwegian sheep breeds, that the
257 incidence of mastitis increased as litter sizes increased. Previous research from Ireland also found that the incidence of
258 mastitis increased with ewe age (O'Brien et al., 2017). This, coupled with the large proportion of terminal sire bloodlines
259 represented in the Irish data (Texel, Suffolk and Charollais), which are also often associated with high lamb growth
260 rates, may explain why mastitis features highly as an end reason in the Irish data. In addition to the premature culling
261 of the ewe herself, the disease can also prove expensive in terms of treatment costs and poor lamb growth rates, therefore
262 having a considerable impact on affected flocks (Bergonier and Berthelot, 2003; Conington et al., 2008; Gelasakis et
263 al., 2015; McLaren et al., 2018).

264 Although Conington et al. (2008) reported mastitis to be the single biggest reason for premature culling in UK Texel
265 sheep, the data available from the UK in the current study was based on records from Scottish Blackface hill ewes, in
266 which there was a very low incidence of mastitis. This observation would be in line with findings from an earlier
267 Norwegian study comparing incidence in different breeds and pasture types, with ewes on lowland pasture having a
268 higher incidence when compared with those in hill or mountain environments (Larsgard and Vaabenoe, 1993). If ewes
269 spend the majority of their time in an extensive hill or mountain environment lower incidence of mastitis may be
270 observed, due to the reduced opportunity for causative bacteria to spread and infect other animals. Ewes that are housed
271 around lambing time may be at a higher risk of infection if there are high stocking densities and poorer hygiene, bedding
272 quality and ventilation levels, all of which provide favourable conditions for bacteria to develop and spread more easily
273 (Gelasakis et al., 2015). There are also lower litter sizes associated with ewes in extensive hill farm systems and Scottish
274 Blackface lambs are often associated with slower growth rates, when compared with terminal sire lambs (Lambe et al.,
275 2007). It could be hypothesised that cull data from terminal sire breeds in the UK may be more similar to the Norwegian
276 and Irish data presented here, although the data were not available to test this in the current study.

277 The data available from the UK hill flocks indicates that the most common reason for mortality/culling in hill ewes was
278 problems identified with their teeth, predominantly tooth loss, also known colloquially as “broken mouth”. Ewes were
279 culled if they had lost at least one of their four centre permanent incisors, usually identified in late autumn when farmers
280 typically sort through their flocks to remove ewes not fit for further breeding, or which are considered unlikely to survive
281 through the winter months. MacGregor (2011) also highlights a number of studies that show tooth loss can lead to
282 reductions in the live weight of the ewe (Dove and Milne, 1991; Williams 1993) and is associated with potential
283 reductions in feed intake. Annett et al. (2011) observed that the survival probability of hill ewes fell gradually as the
284 number of teeth missing increased. There have been a number of different reasons linked to the development of tooth
285 loss, including the effects of farm of origin, pasture type, breed, mineral nutrition and supplementary feeding to name
286 just a few (MacGregor, 2011). Less than 5% of the ewes recorded in Ireland were recorded as having been culled due
287 to their teeth whereas the option to identify ewes culled specifically for tooth loss was not available to farmers in
288 Norway. It is therefore possible that some ewes with missing teeth may have been identified in the Norwegian data set
289 as being culled for age, body condition, or another reason.

290 Ewes culled because of their age also featured relatively highly across all three countries. In the UK, it is common
291 practise for hill ewes to leave the flock at 5 years old, having had 4 lamb crops. These ewes would then be sold either
292 for slaughter or onto lowland flocks for further breeding. However, depending on the number of younger replacement

293 ewes available each year, hill farmers may decide to keep some ewes beyond this age, to maintain their overall flock
294 numbers, providing their body condition is good and they have retained all of their teeth. As the UK data in this study
295 came from research flocks where a strict culling policy was adhered to, animals were only recorded as culled for age if
296 there were no other reasons for culling them. This might not necessarily be the case in the Norwegian or Irish data as,
297 mentioned earlier, some of these animals recorded as being culled for age could have been culled for other reasons such
298 as tooth loss.

299 *4.2 Early-life predictors of age at last lambing*

300 In addition to identifying the main reasons for mortality/culling in flocks across Norway, Ireland and the UK, this study
301 also investigated potential early-life predictors of ewe longevity across all three countries. The longevity trait selected
302 was age at last lambing, as this was easily defined in the data collected from different performance recording schemes
303 and could be compared across all countries. The data available for age at last lambing, from each country, revealed that
304 on average, ewes in Norway left the flock at 3.38 years old, whereas those in Ireland and the UK left at 4.22 and 4.35
305 years old, respectively, although no statistical test could be performed to formally assess differences between countries.
306 It should also be noted that all of the Norwegian ewes lambed for the first time at 1 year old, whereas ewes in the UK
307 data did not have their first lamb until they were a year older. The Irish data contained ewes that lambed for the first
308 time as both 1 and 2 year olds, although an average age at first lambing of 1.88 would indicate a higher proportion of
309 ewes were lambing for the first time as a 2 year old.

310 The age at last lambing records differed across all three countries, likely influenced by the main mortality/culling reasons
311 highlighted above and the rules imposed for retaining data in the different analyses. In Norway and Ireland,
312 approximately the same percentage of ewe records came from each age group (1-6 years old). This would match with
313 the fact that a high percentage of ewes are culled for mastitis and other udder associated problems, which can occur at
314 any point throughout the ewe's lifetime. In contrast, the percentage of age at last lambing records available for ewes
315 culled in the UK hill flocks increased steadily up to 5 years old before tailing off in later age groups. This pattern
316 reflecting the fact that the main culling reasons identified were tooth loss or age, both of which are traits associated with
317 later life. Results observed by Mekki et al. (2009), found that younger crossbred Mule ewes were most commonly
318 culled for udder problems, such as mastitis, whereas the number of ewes culled for teeth associated issues increased
319 substantially after 4 years old.

320 In terms of possible early-life factors that may predict how long the ewe will remain in the flock, the results were not
321 consistent, with very few factors being significant across all three countries. Age at last lambing was significantly
322 affected by birth year of the ewe across all analyses, for the years that were studied here, but predicted means for these
323 years could not be used to predict future ewe longevity. Other significant effects included the ewe's dam's age. This
324 was most evident in the Norwegian analyses, where ewes born to younger dams (i.e. 1 and 2 year olds) left the flock
325 earlier than ewes born to older dams. Recent results from New Zealand have found that the survival rates of ewes born
326 (as a twin) to 1 year old mothers were shorter when compared with those born to older ewes, although it should be noted
327 that a relatively small number of animals were used in this study (Pettigrew et al., 2019). The relationship in the UK
328 analyses, although significant was less clear. Interestingly, age at first lambing (only fitted in the Irish analyses) proved
329 to be a significant effect and suggested ewes lambing for the first time as a 1 year old left the flock earlier than those
330 who lambed for the first time at 2 years old. Positive genetic and phenotypic correlations between age at first lambing
331 and age at last lambing have previously been observed by McLaren et al. (2017) in Lleyn and Dorset sheep in the UK
332 and therefore in agreement with the Irish results in this study. However, in contrast, Kenyon et al. (2011) found breeding
333 Romney ewes at 1 year old did not significantly reduce their longevity. This variation in results suggests there could be
334 breed and environment differences, and further investigation may prove useful.

335 *4.2 Limitations and recommendations*

336 Over the course of this study, several limitations were identified, particularly in terms of accurately assessing differences
337 between countries. Having identified the main reasons across each country (mastitis, tooth loss and old age) there
338 remains a wide range of additional reasons provided by each recording scheme that farmers can allocate to each ewe.
339 The question therefore arises – in how much detail should cull reasons be recorded? Is it more valuable to capture
340 detailed information on why the ewes are leaving the flock, or should the number of recorded culling reasons be
341 minimised to identify the main reason types and simplify recording? The reasons identified as responsible for over 5%
342 of culls/deaths in each country did not include any lambing-related issues or physical/structural reasons, such as those
343 associated with the legs or feet, which could potentially be under genetic control. Providing culling reasons such as old
344 age or slaughter on a list of options, may mean that breeders record these reasons as a default, rather than the root causes,
345 which may include poor body condition, tooth loss, lambing or reproductive issues, for example. Clear instruction should
346 therefore perhaps be provided to breeders to ensure that culling or death reasons that could have a genetic basis are
347 identified as the primary cause for ewe removal from the flock. Involuntary reasons, such as ewes lost due to
348 accident/injury, predation, or sold for management reasons such as food shortages (the latter two particularly relevant

349 to Norway), were also all below the 5% threshold. These involuntary culls or deaths are useful to know for management
350 or compensatory reasons, but could be combined, and records potentially removed, for consideration of longevity within
351 genetic analyses.

352 With mastitis and tooth loss being the main drivers of ewes leaving the flock in these data sets, genetic analyses of these
353 direct traits could be a useful option to explore in order to try to reduce the impact these traits have in terms of longevity.
354 These traits would be more easily defined and may be affected less by management policies across different farms,
355 reducing residual variation compared with the more general trait of age at last lambing. This could potentially increase
356 the heritability of the trait under selection, leading to faster genetic gain. However, to include reduced mastitis or tooth
357 loss in future breeding goals, accurate recording of these traits by the breeders would be required. The two traits are
358 very different in terms of their effect, mastitis affecting ewes of all ages while tooth loss predominantly affected older
359 ewes. They are also very different due to the nature of the different breeds and systems studied in the present analyses -
360 mastitis affecting predominantly terminal sire type breeds (i.e. high growth rates, high litter sizes) with lower incidence
361 in a more hill type breed; whilst tooth loss tends to be more prevalent in ewes grazing poorer quality hill swards.

362 Due to the different management and breed structures of the data available from each country, only observational
363 comparisons between countries could be made. Although the final models used to assess age at last lambing were as
364 similar as possible, the different analyses could not be compared using any statistical analyses. The main differences
365 across the three countries, which could not be ignored in the models used, were the breeds involved (pure breeds in
366 Norway and UK compared to a mix of pure and crossbred animals in Ireland) and the age of the ewes at their first
367 lambing (one year old in Norway, two years old in UK and a combination of both in Ireland). Nonetheless, these analyses
368 help towards improving our knowledge and understanding of factors influencing ewe longevity across each country.
369 The results from this study, both in terms of the main reasons identified across each country for mortality/culling and
370 the analyses of age at last lambing to assess ewe longevity, suggest further work would be beneficial before attempting
371 to combine data across countries or systems, or infer longevity implications in one country based on results from another.

372 **5. Conclusion**

373 This study has helped to further our understanding of the main causes of ewe mortality or culling (voluntary or
374 involuntary) across 3 different European sheep production systems. Differences were observed in the retention rate of
375 breeding ewes in the flocks across ages, as well as in the main reasons for ewes leaving the flock, which suggest that
376 longevity is influenced by different factors across countries and systems. The early life predictors investigated could not

377 be used to provide consistent recommendations across countries and production systems that would influence ewe
378 longevity. Therefore, one common definition or solution to select ewes for longer productive life in divergent sheep
379 flocks may not be appropriate.

380 **Acknowledgment and funding**

381 This work was funded by the Irish Department of Agriculture, Food and the Marine (DAFM), the Research Council of
382 Norway (Project number 272338), the Norwegian Association of Sheep and Goat Breeders (NSG) and the UK
383 Department for Environment, Food and Rural Affairs (DEFRA) through the ERA-NET SUSAN project SusSheP.

384 Thanks also to the Norwegian Meat and Poultry Research Centre and Sheep Ireland for access to commercial flock data.
385 SRUC receives financial support from the Scottish Government's Rural and Environment Science and Analytical
386 Services Division to collect research flock data.

387 Finally, many thanks to all farmers and research technicians involved in the collection and submission of data.

388

389 **References**

- 390 Animalia, 2018. Arsmelding 2018. [https://www.animalia.no/globalassets/saukontrollen---](https://www.animalia.no/globalassets/saukontrollen---dokumenter/arsmelding-sau-2018.pdf)
391 [dokumenter/arsmelding-sau-2018.pdf](https://www.animalia.no/globalassets/saukontrollen---dokumenter/arsmelding-sau-2018.pdf) (accessed 7th October 2019).
- 392 Annett, R.W., Carson, A.E., Dawson, L.E.R., Irwin, D., Gordon, A.W., Kilpatrick, D.J., 2011. Comparison
393 of the longevity and lifetime performance of Scottish Blackface ewes and their crosses within hill
394 sheep flocks. *Animal*. 5:3, 347-355.
- 395 Bergonier, D., Berthelot, X., 2003. New advances in epizootiology and control of ewe mastitis. *Liv. Prod.*
396 *Sci.* 79, 1-16.
- 397 Borg, R.C., Notter, D.R., Kott, R.W., 2009. Genetic analysis of ewe stayability and its association with lamb
398 growth and adult production. *J. Anim. Sci.* 87, 3515-3524
- 399 Brash, L., Fogarty, N., Gilmour, A., 1994. Reproductive performance and genetic parameters for Australian
400 Dorset sheep. *Aust. J. Agri. Res.* 45, 427-441.
- 401 Conington, J., Bishop, S.C., Grundy, B., Waterhouse, A., Simm, G., 2001. Multi-trait selection indexes for
402 sustainable UK hill sheep production. *Anim. Sci.* 73, 413-423.
- 403 Conington, J., Cao, G., Stott, A., Bunger, L., 2008. Breeding for resistance to mastitis in United Kingdom
404 sheep, a review and economic appraisal. *Vet. Rec.* 162, 369-376.
- 405 Dove, H., Milne, J.A., 1991. An evaluation of the effects of incisor dentition and of age on the performance
406 of lactating ewe and their lambs. *Anim. Prod.* 53, 183-190.
- 407 Eikje, L.S., Adnoy, T., Klemetsdal, G., 2008. The Norwegian sheep breeding scheme: description, genetic
408 and phenotypic change. *Animal*. 2, 167-176.
- 409 Gelasakis, A.I., Mavrogianni, V.S., Petridis, I.G., Vasileiou, N.G.C., Fthenakis, G.C., 2015. Mastitis in
410 sheep – The last 10 years and the future of research. *Vet. Microbiol.* 81, 136-146.
- 411 Jakobsen, J.H., Holme, I.-J., Boman, I.A., Blichfeldt, T., 2017. Curve linear weighting of litter size in the
412 Norwegian Total Merit Index. Poster at Sheep Breeders Round Table, Nottingham, UK.
413 <https://www.nationalsheep.org.uk/workspace/pdfs/norway-2.pdf> (accessed 7th September 2019).
- 414 Jones, O.R., Jones, D.L., Edwards-Jones, G., Cross, P., 2013. Informing decision making in agricultural
415 greenhouse gas mitigation policy: A Best–Worst Scaling survey of expert and farmer opinion in the
416 sheep industry. *Env. Sci. & Pol.* 29: 46-56.

417

418 Kenyon, P.R., van der Linden, D.S., West, D.M., Morris, S.T., 2011. The effect of breeding hoggets on
419 lifetime performance. *N. Z. J. Agri. Res.*, 54:4, 321-330.

420 Lambe, N.R., Navajas, E.A., McLean, K.A., Simm, G., Bunger, L., 2007. Changes in carcass traits during
421 growth in lambs of two contrasting breeds, measured using computer tomography. *Liv. Sci.* 107, 37-
422 52.

423 Lambe, N.R., Conington, J., Bishop, S.C., McLean, K.A., Bunger, L., McLaren, A., Simm, G., 2008.
424 Relationships between lambs carcass quality traits measured by X-ray computed tomography and
425 current UK hill sheep breeding goals. *Animal* 2, 36-43.

426 Lambe, N.R., Wall, E., Ludemann, C.I., Bunger, L., Conington, J., 2014. Genetic improvement of hill sheep
427 – Impacts on profitability and greenhouse gas emissions. *Small Rumin. Res.* 120, 27-34.

428 Larsgard, A.G., Vaabenoe, A., 1993. Genetic and environmental causes of variation in mastitis in sheep.
429 *Small Rumin. Res.* 12, 339-347.

430 Lee, M.A., Cullen, N.G., Mewman, S.A.N., Dodds, K.G., McEwan, J.C., Shackell, G.H., 2015. Genetic
431 analysis and genomic selection of stayability and productive life in New Zealand ewes. *J. Anim. Sci.*
432 93, 3268-3277.

433 McGregor, B.A., 2011. Incisor development, wear and loss in sheep and their impact on ewe production,
434 longevity and economics: A review. *Small Rumin. Res.* 95, 79-87.

435 McLaren, A., Kaseja, K., Moore, K., Mucha, S., Boon, S., Conington, J., 2017. Genetic aspects of ewe
436 longevity and fertility traits in Lleyn and Dorset sheep. In *Proceedings of the British Society of
437 Animal Science Conference, Advances in Animal Biosciences. The Future of Animal Science*, 26-27
438 April 2017, Chester, UK, pp. 101.

439 McLaren, A., Kaseja, K., Yates, J., Mucha, S., Lambe, N.R., Conington, J., 2018. New mastitis phenotypes
440 suitable for genomic selection in meat sheep and their genetic relationships with udder conformation
441 and lamb live weights. *Animal*, 12, 2470-2479.

442 Mekki, W., Roehe, R., Lewis, R., Davies, M.H., Bunger, L., Simm, G., Haresign, W., 2009. Genetic
443 relationship between longevity and objectively or subjectively assessed performance traits in sheep
444 using linear censored models. *J. Anim. Sci.*, 87, 3482-3489.

- 445 O'Brien, A.C., McHugh, N., Wall, E., Pabiou, T., McDermott, K., Randles, S., Fair, S., Berry, D.P., 2017.
446 Genetic parameters for lameness, mastitis and dagginess in a multi-breed sheep population. *Animal*,
447 11:911-919. doi: 10.1017/S1751731116002445.
- 448 Pabiou, T., Byrne, T., Wall, E., McHugh, N., 2014. Genetic improvement of sheep in Ireland. Proceedings
449 of the 10th World Congress of Genetics Applied to Livestock Production, Vancouver, Canada.
450 <http://www.wcgalp.org/system/files/proceedings/2014/genetic-improvement-sheep-ireland.pdf>
451 (accessed 7th October 2019)
- 452 Pettigrew, E.J., Hickson, R.E., Morris, S.T., Lopez-Villalobos, N., Pain, S.J., Kenyon, P.R., Blair, H.T.,
453 2019. The effects of birth rank (single or twin) and dam age on the lifetime productive performance
454 of female dual purpose sheep (*Ovis aries*) offspring in New Zealand. *PLoS ONE* 14(3): e0214021.
455 <https://doi.org/10.1371/journal.pone.0214021>
- 456 SAS (2013) SAS Institute Inc., 2013. SAS/STAT® 9.4 User's Guide. Cary, NC: SAS Institute Inc.
- 457 Santos, B.F.S., McHugh, N., Byrne, T.J., Berry, D.P., Amer, P.R., 2015. Comparison of breeding objectives
458 across countries with application to sheep indexes in New Zealand and Ireland. *J. Anim. Breed.*
459 *Genet.* 132; 144–154. doi:10.1111/jbg.12146.
- 460 VSN International (2013) *Gentstat for Windows 16th Edition*. VSN International, Hemel Hempstead, UK
- 461 Waage, S., Vatn, S., 2008. Individual animal risk factors for clinical mastitis in meat sheep in Norway. *Prev.*
462 *Vet. Med.* 87, pp 229-243. <https://doi.org/10.1016/j.prevetmed.2008.04.0024.002>
- 463 Williams, A., 1993. Evaluation of tooth grinding as a method for improving economic performance in flocks
464 with premature incisor tooth loss ("broken mouth"). Final Report Project DAV 5 Wool Research and
465 Development Corporation. Victorian Department of Agriculture, Werribee.
- 466 Zishiri, O.T., Cloete, S.W.P., Olivier, J.J., Dzama, K., 2013. Genetic parameters for growth, reproduction
467 and fitness traits in the South African Dorper sheep breed. *Small Rumin. Res.* 112, 39-48.

Table 1. Summary statistics for the fixed effects fitted in the linear models

Effects fitted	Norway				Ireland				UK			
	Av.	S.D.	Min	Max	Av.	S.D.	Min	Max	Av.	S.D.	Min	Max
Ewe birth weight (kg)	4.93	0.94	1.5	9.5	4.55	1.06	2	10	3.77	0.66	1.6	6.5
Ewe weight – approx. 6-8 weeks old (kg)*	14.56	2.72	5	30	19.30	3.81	12	32	18.15	1.04	14.2	22.17
Ewe weight – approx. 14-20 weeks old (kg)**	39.95	4.64	30	50	35.62	6.91	20	55	27.41	1.05	24.1	31.98
Ewe age at first lambing (years)	-	-	-	-	1.88	0.33	1	2	-	-	-	-
Dam of ewe age (years)	2.60	1.19	1	6	2.76	1.54	1	7	3.51	1.15	2	7
Birth litter size of ewe	2.41	0.75	1	4	1.87	0.68	1	4	1.59	0.53	1	3
Breed proportion of Texel (%)	-	-	-	-	45.91	48.69	0	100	-	-	-	-
Breed proportion of Suffolk (%)	-	-	-	-	16.89	35.16	0	100	-	-	-	-
Breed proportion of Belclare (%)	-	-	-	-	18.48	35.16	0	100	-	-	-	-
Breed proportion of Charollais (%)	-	-	-	-	5.20	21.59	0	100	-	-	-	-
Breed proportion of Vendeen (%)	-	-	-	-	4.70	20.21	0	100	-	-	-	-

*Ewe live weight: live weight corrected to 56 and 40 days old for UK and Irish data, respectively. Live weight gain from birth corrected to 42 days old for Norwegian data

**Ewe live weight: live weight corrected to 111 and 100 days old for UK and Irish data, respectively. Live weight gain from birth corrected to 140 days old for Norwegian data

221 **Table 2.** Most numerous cull and death reasons provided by commercial/research farms in Norway,
 222 Ireland and the UK (as a percentage of total ewe deaths or culls recorded in each country)

Norway	Ireland	UK
Mastitis (19.9%)	Age (20.9%)	Teeth (38.9%)
Udder problems (16.9%)	Died (reason unknown) (19.9%)	Age (23.5%)
Age (12.4%)	Slaughtered (15.9%)	Body condition (6.8%)
Non-breeder (6.3%)	Mastitis (13.5%)	Missing – presumed dead (6.8%)
	Unknown reason (6.4%)	Reproductive disorder (5.4%)
	Body condition (5.7%)	
	Udder problems (5.7%)	

223

224 **Table 3.** Summary statistics for age at last lambing data available from Norway, Ireland and the UK

Trait	Norway				Ireland				UK			
	Av.	S.D.	Min	Max	Av.	S.D.	Min	Max	Av.	S.D.	Min	Max
Ewe age at last recorded lambing event (years)	3.38	1.79	1	6	4.22	1.68	1	7	4.35	1.12	2	7

225

226

227


Table 4. Details of the effects fitted in each model, and their significance, across each country.

Effects fitted in the models	Norway	Ireland	UK
Ewe birth year	✓***	✓***	✓***
Flock of birth	✓***	✓***	✓*
Dam age	✓***	✓ns	✓*
Ewe birth weight	✓ns	✓***	✓ns
Birth litter size of ewe	✓**	✓***	✓ns
Ewe weight – approx. 6-8 weeks old	✓ns	✓**	✓ns
Ewe weight – approx. 14-20 weeks old	✓ns	✓ns	✓ns
Ewe weight – approx. 6-8 weeks old (squared)	✓ns		✓ns
Ewe weight – approx. 14-20 weeks old (squared)	✓ns		✓*
Ewe age at first lambing		✓***	
Breed proportion of Texel (%)		✓***	
Breed proportion of Suffolk (%)		✓***	
Breed proportion of Belclare (%)		✓***	
Breed proportion of Charollais (%)		✓ns	
Breed proportion of Vendeen (%)		✓*	
Ewe birth year x Flock of birth	✓***		✓***
Variance accounted for by the models (R ²)	10.8%	31.0%	3.5%

Residual means standard error (RMSE)	1.74	1.60	1.14
--------------------------------------	------	------	------

228
229
230

✓ Fitted in model; *** P<0.001; ** P<0.01; * P<0.05; ^{ns} Not Significant




231

232

Fig 1. Frequency of age at last lambing across all three counties

233



234

235 Fig 2. Predicted least squared means for the effects of a) the age of the ewe's dam and b) birth litter
 236 size of the ewe across all countries. (Means sharing the same superscript, within country, are not
 237 significantly different from each other ($P > 0.05$)).

238