

Scotland's Rural College

The global tendency in the research of biological activity in endophytic fungi

Dos Santos, I. R.; Mohamed, T. A.; Borges, L. L.; Nahas, HH Abo; Abdel-Azeem, M. A.; Carvalho, CDC; Bezerra, J. P.; Gupta, V. K.; Xavier-Santos, S.; Mossa, M. I.; Abdel-Azeem, A. M.

Published in:
Current Research in Environmental and Applied Mycology

DOI:
[10.5943/cream/12/1/1](https://doi.org/10.5943/cream/12/1/1)

First published: 12/01/2022

Document Version
Publisher's PDF, also known as Version of record

[Link to publication](#)

Citation for published version (APA):
Dos Santos, I. R., Mohamed, T. A., Borges, L. L., Nahas, HH. A., Abdel-Azeem, M. A., Carvalho, CDC., Bezerra, J. P., Gupta, V. K., Xavier-Santos, S., Mossa, M. I., & Abdel-Azeem, A. M. (2022). The global tendency in the research of biological activity in endophytic fungi: a scientometric analysis. *Current Research in Environmental and Applied Mycology*, 12, 1-14. Advance online publication. <https://doi.org/10.5943/cream/12/1/1>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



The global tendency in the research of biological activity in endophytic fungi: a scientometric analysis

dos Santos IR¹, Mohamed TA², Borges LL¹, Abo Nahas HH³, Abdel-Azeem MA⁴, Carvalho DDC⁵, Bezerra JP⁶, Gupta VK⁷, Xavier-Santos S¹, Mossa MI⁸ and Abdel-Azeem AM^{9*}

¹Universidade Estadual de Goiás, Campus Central, Laboratório de Micologia Básica, Aplicada e Divulgação Científica – FungiLab, Anápolis, Goiás, Brazil

²Dental Basic Sciences Department, College of Dentistry, University of Duhok 42001, Iraq

³Suez Canal University, Faculty of Science, Zoology Department, Ismailia 41522, Egypt

⁴Sinai University, Faculty of Pharmacy and Pharmaceutical Industries, Arish, North Sinai, Egypt

⁵Universidade Estadual de Goiás, Campus Sudeste, Laboratório de Fitopatologia, Ipameri, Goiás, Brazil

⁶Universidade Federal de Goiás, Instituto de Patologia Tropical e Saúde Pública, Departamento de Biociências e Tecnologia, Goiânia, Goiás, Brazil

⁷Biorefining and Advanced Materials Research Center, Scotland's Rural College (SRUC), Kings Buildings, West Mains Road, Edinburgh, EH9 3JG, United Kingdom; Center for Safe and Improved Food, Scotland's Rural College (SRUC), Kings Buildings, West Mains Road, Edinburgh, EH9 3JG, United Kingdom

⁸Al Arish University, Faculty of Science, Botany and Microbiology Department, North Sinai 45516, Egypt

⁹Suez Canal University, Faculty of Science, Botany and Microbiology Department, Ismailia 41522, Egypt

dos Santos IR, Mohamed TA, Borges LL, Abo Nahas HH, Abdel-Azeem MA, Carvalho DDC, Bezerra JP, Gupta VK, Xavier-Santos S, Abdel-Azeem AM 2022 – The global tendency in the research of biological activity in endophytic fungi: a scientometric analysis. *Current Research in Environmental & Applied Mycology (Journal of Fungal Biology)* 12(1), 1–14, Doi 10.5943/cream/12/1/1

Abstract

Endophytic fungi colonize plants' interior tissues and organs, establishing a mutualistic relation without causing apparent harm. The biotechnological importance of these fungi has inspired research worldwide. Through scientometrics methods, it is possible to verify tendencies and gaps in different research area, collaborating with academic society to foster new studies and correlations. To identify the production of scientific knowledge relating to biological activity within endophytic fungi, we performed a survey on the platform Web of Science from studies published between 1997 and 2019 that have the following terms in the title, summary, or keywords: (Fung* AND Endophytic) AND (Bioprospecting OR Bio-prospecting OR Prospecting OR Bioactivit* OR “Biological Activit*”). A total of 458 studies were obtained, of which the following information was verified: publication year; average citations per year; journal; h-index of the journal; area of concentration; keyword and co-occurrence between them; network of collaboration between countries, institutions, and journals; bioassays realized and biological activity verified among endophytic fungi. China represents the nation-state that most frequently hosts research, followed by Brazil and India. According to our findings, the period between 2016 and 2019 showed an increase in the number of bioassays, nearly half of which (43%) proved to have antimicrobial activity, followed by anti-tumor activity (29%). These properties show that endophytic fungi can produce bioactive molecules of pharmaceutical and agronomic interest.

Keywords – Anti-tumor – Bio-activity – China – H-index – Natural products – Quantitative research – Scientometric indicators

Introduction

Endophytic microbiota, including fungi, live asymptotically in the living tissue of healthy plants, interacting with the host plant by aiding growth and resistance to herbivory, pathogens, and abiotic stresses (Rana et al. 2019a). In return, the endophytes receive protection and nutrients from the host (Rana et al. 2019b). However, at some point, this may turn pathogenic due to biotic or abiotic interferences (Rodriguez & Redman 2008), suggesting that the interaction between the fungus and the host plant should be characterized by a balance of potential of virulence of the fungus and sustained defense of the plant (Schulz et al. 2002, Saikkonen et al. 2004). In addition, it is known that every plant host can harbor an endophytic community. However, the frequency and composition of the endophytes depend on the environmental characteristics (mainly the climate), the geographic distribution of the plants, their age, and the specific tissue involved (Strobel 2003, Saikkonen et al. 2003).

Enzymes and secondary metabolites from endophytic fungi have shown tremendous biotechnological importance, principally in the composition of novel pharmaceuticals and the biological control of pests within agriculture (Gangadevi & Muthumary 2009a). Assays of bio-prospecting among these organisms revealed natural composites with anti-bacterial, anti-fungal, anti-viral, anti-tumor, anti-inflammatory, anti-oxidant, and anti-protozoal activities with prospective for use in the treatment of disease and notable industrial applications (Strobel & Daisy 2003, Chapla et al. 2013). In addition, the manipulation of endophytic microbes enables more efficient methods of pharmaceutical production, minimizing the risk of extinction to plant species, some of which throughout the years have been suffering indiscriminate pressure from over harvest or malfeasance related to the extraction of their medicinal compounds (Peixoto et al. 2002).

The bibliographic survey realized by Jia et al. (2016) pointed out that 96 medicinal species of plants harbor endophytic fungi that produce bioactive compounds. The same hosts are commonly used as medication using direct consumption or extraction of active components. Species of the family Taxaceae are related to endophyte producers of taxol, a diterpenoid with anti-tumor activity mainly used to treat ovarian, uterine, and breast cancer (Gangadevi & Muthumary 2009b).

The endophytes possess a capacity to promote the accumulation of secondary metabolites within their host plants. Thus, researchers now isolate these microorganisms in plants with known therapeutic properties (Chen et al. 2016a). This is justified by the accelerated loss of medicinal species worldwide, compromising more than 50,000 species due to natural habitat destruction (Venieraki et al. 2017). To respond to the medical and agro-industrial demand, it is important to find alternative sources of bioactive compounds, especially for endangered and difficult to cultivate hosts (Atanasov et al. 2015).

Research involving the benefits of endophytic fungi has gained prominence throughout the world (Jalgaonwala et al. 2017), as presented by the review of Rashmi et al. (2019), which relates a list of more than 800 genera that present bio-active compounds, predominantly *Alternaria*, *Aspergillus*, *Colletotrichum*, *Fusarium*, *Penicillium* and *Phoma*. Intending to make a survey of the tendencies of biological activities of these microorganisms, scientometrics is an important ally in the evaluation of the quantitative aspects of research, including scientific journals, authors, and communication between the scientist (Macias-Chapula 1998, McGrath 1989). Considering the advancement of research verifying the biotechnological potential of endophytic fungi, in this review, we intend to contribute to the field with a scientometric analysis of global scientific literature on the biological activity of endophytic fungi.

Materials & Methods

From the publications available on the platform Web of Science during the period of 1945 to 2019, a study was performed with the following search terms and respective Boolean operators' included in the title, summary, and/or keywords: (Fung* AND Endophytic) AND (Bioprospecting

OR Bio-prospecting OR Prospecting OR Bioactivit* OR “Biological Activit*”). The retrieved publications were individually evaluated for compliance with the theme and categorized into original correction and review articles.

For every paper recovered from the platform, the following information was registered: a) year of publication, b) average of citations of articles per year, c) journal in which the study was published, d) h-index of the journal, e) area of concentration, f) keywords and co-occurrence between them, g) authors and co-authors, h) most cited articles, i) h-index of the ten most published authors, j) national affiliation of the authors and co-authors, k) collaboration between countries, institutions, and journals, l) bio-assays realized per year and m) verified biological activities of endophytic fungi. The analysis was made in the software (R Core Team 2020) with the help of the package bibliometrix (Aria & Cuccurullo 2017). For the association between keywords, a co-occurrence network was produced, stemming from the walk trap clustering algorithm, which captures the structure of the community simulating random walks in the networks (Pons & Latapy 2006). A flow chart (Sankey diagram) presented the collaboration between countries, institutions, and journals.

Results

The initial search resulted in 675 scientific publications, of which 458 correlated with the theme of the study, with 392 original articles, 3 corrections articles, and 63 review articles. The recovered publications are distributed among 197 journals. In addition, 1329 keywords and an average of 18.87 citations per article were found. Although the search began in 1945, the first studies found approaching this theme date to 1997 (n = 2). There was a marked increase in studies from the year 2005. The highest number of studies was in 2018 (n = 70) (Fig. 1a). The largest average of citations per article was in 2006 (n = 159) (Fig. 1b). The significant number of citations in this year is due to the article “Natural products from plant-associated microorganisms: distribution, structural diversity, bioactivity, and implications of their occurrence” published in the *Journal of Natural Products* (Gunatilaka 2006), that received 528 citations, followed by the articles “Molecular characterization and anti-microbial activity of endophytic fungi from coffee plants” published in the *World Journal of Microbiology and Biotechnology* (Sette et al. 2006), with 62 citations, and “Marinamide, a novel alkaloid and its methyl ester produced by the application of mixed fermentation technique to two mangrove endophytic fungi from the South China Sea” published in the *Chinese Science Bulletin* (Zhu & Lin 2006), with 45 citations. This time series shows the visibility of articles in 2006, with experimental and review articles, bringing an ample variety and characterization of bioactive molecules. On the other hand, the articles published from 2011 do not achieve scientific repercussion sufficient to be cited.

The journal *Marine Drugs* presented the highest number of publications (n = 16), followed by the *World Journal of Microbiology & Biotechnology* (n = 15) and the *Journal of Natural Products* (n = 15) (Fig. 2a). The journals with a higher h-index were the *European Journal of Organic Chemistry* (n = 13), followed by *Marine Drug* (n = 10), *Planta Medica* (n = 8), and *Journal of Natural Products* (n = 8) (Fig. 2b). Concerning the area of concentration, 52 studies are in the area of “Microbiology”, followed by “Chemistry” (n = 45), and “Applied Biotechnology and Microbiology” (n = 43) (Fig. 2c). In addition, other areas of concentration were found (e.g. “Pharmacology and Pharmacy” “Bio-chemistry and Molecular Biology”, and “Science and Technology”), noting that 229 articles fit into two or more areas of concentration.

The search for the most used keywords revealed that “natural-products” was included 95 times; followed by “metabolites” and “secondary metabolites”, with 84 and 56 occurrences, respectively (Fig. 3a); both terms also appear prominently in the co-occurrence network (Fig. 3b). The keyword “natural-products” is used with the keyword’s “metabolites,” “identification”, “diversity,” “microorganisms”, among others, as the keyword “secondary metabolites” is used with the keywords “biological-activity”, “biological-activities”, “endophytic fungus”, “in-vitro” and “fungi”, among others.

The authors and co-authors with the highest number of publications are Barbara Schulz (SCHULZ, B) and Siegfried Draeger (DRAEGER, S) (each with 16 articles), followed by Karsten

Krohn (KROHN, K) (n = 15) (Fig. 4a). The article with the most citations is “Natural products from plant-associated microorganisms: distribution, structural diversity, bioactivity, and implications of their occurrence” (Gunatilaka 2006) with 513 citations, the equivalent of 37 citations per year (Fig. 4b), which presents a review of microorganisms associated with plants as an enormous and unexplored resource of natural products, with chemical structures of biological relevance and a particular focus on endophytic fungi.

China represented the country with the highest number of publications (412 articles), followed by Brazil (n = 189) and India (n = 160) (Fig. 5a). The United States presents the largest collaboration network between countries, linking researchers in China, India, Brazil, Egypt, and the United Kingdom. The institutions most published between different journals are the University of São Paulo, Federal University of Paraná, and the Federal University of Minas Gerais (Brazil), and Sun Yat-sen University, China Agricultural University, and Zhejiang University (China). It was also verified that the University of Mississippi (USA) obtained the highest number of publications in different journals (n = 7). Sun Yat-sen University (China) was the only institution to publish Marine Drugs (Fig. 5b).




Fig. 1 – Temporal distribution. (a) scientific production. (b) the average number of citations of articles per year. The exponential crescent curve (a) denotes a general tendency (p-value: 2.85-13).






Fig. 2 – Distribution of the (a) articles per journal, (b) journals per h-index, (c) articles per area of concentration.




Fig. 4 – Distribution of (a) authors following the total of publications, (b) articles following the total citations.





Fig. 5 – Principal countries (a) to host research about the biological activity of endophytic fungi, (b) institutions and journals concerning collaboration of research.

The analyses related to biological activities addressed in the 395 experimental articles reveal 484 bio-assays (some studies verified only one activity and others more than one activity). A marked increase of bio-assays began in 2014, with the most in 2018 (n = 81), followed by 2019 (n = 79), 2016 (n = 51), 2017 (n = 48) e 2014 (n = 42) (Fig. 6).

The categorization of these biological activities resulted in 12 groups (e.g. antimicrobial, anti-tumor, enzymatic, anti-oxidant, insecticidal, anti-inflammatory, anti-diabetic, immunomodulatory, seed germination stimulant, anti-obesity, anti-algal, anti-hemolysis). The anti-microbial activity (anti-bacterial, anti-fungal, anti-parasitical, anti-viral) corresponded to almost half of the bio-assays (43%, n = 208), followed by anti-tumor (29%, n = 140) and enzymatic activity (9%, n = 44). Other types of activities (e.g. anti-oxidant, insecticidal, anti-inflammatory, anti-diabetic, immunomodulatory, seed germination stimulant, anti-obesity, anti-algal, anti-hemolysis) had less than 9% (Fig. 7).




Fig. 6 – Distribution of the bio-assays described in the scientific literature catalogued on the platform Web of Science, from 1997 to 2019.




Fig. 7 – Distribution of the articles following the type of biological activity.

Discussion

Throughout the years, the evolution of publications about biological activity among endophytic fungi has increased researcher's interest in discovering novel bioactive compounds, considering that the number of scientific papers is the most efficient way to measure this growth (Verbeek et al. 2002). The first publications in this area began in 1997, thus corroborating with the survey of (Gupta et al. 2020), which mentions that the first experimental studies involving this theme, although few, date to the 1990's, therein marking the start and establishing appreciation amongst researchers for the bioactive compounds of these microorganisms. Although the first studies of the biotechnological importance of endophytic fungi were published just before the turn of the 21st century, these endophytes already had been related in previous studies (e.g. from 1866) and described in the fossilized tissue of plants, indicating that this mutualistic symbiotic relation is ancient (Redecker et al. 2000, Krings et al. 2007).

Since the discovery of the biotechnological potential of endophytes, a significant increase in scientific production in the area has been observed, beginning around the turn of the 2000s, which may be due to better financing amongst universities and research institutes throughout the world, offering grants for academic training and investment in science, in addition to the increased attention given for the newfound applicability in the treatment of illnesses (Frank & Meyer 2007). The increase of scientific research is due to the constant search for novel bioactive compounds with opportunities for application in medicine. Pandey et al. (2014) pointed out that these investigations will continue to advance, bringing better and more accurate perceptions of such compounds and their biological activities, a fact observed in this study that demonstrates a significant increase of scientific publications in 2018 and 2019.

The research involved in the biological activity of endophytic fungi is aligned in three large areas of concentration (microbiology [including applied microbiology], chemistry and biotechnology), because the metabolites and extracellular enzymes produced by these microorganisms can be applied in various industrial and agricultural processes (Yadav 2018). This encourages the isolation of endophytic fungi from hosts not yet studied in the search for new natural products and justifies the use of the keywords like “natural products”, “metabolites,” and “secondary metabolites.” Strobel (2003) shows that the biological association between endophyte and host plant produces a diversity of secondary metabolites. This association may have developed a genetic system that permits transference between them. Meanwhile, because the publications were realized in a short and determined time (Fig. 1a) – less than two decades – the interaction between researchers can explain the repetition of methodologies, which facilitates the discussion of the data, and with this, the repetition of keywords (Chen et al. 2016b).

Between the ten authors with the most publications, four belong to institutions in Germany (Barbara Schulz, Siegfried Draeger, Karsten Krohn, and Peter Proksch). The German government has been establishing research and innovation in line with its growth policies, ensuring basic and essential conditions conducive to Education (Neves & Neves 2011). Within these policies, the 2005 program High tech Strategies stands out as the largest ongoing program of research and innovation; aside from this example, other policies can also be mentioned to foster the progress of research at universities (e.g. Initiative of Excellence, the Pact of Superior Education and the Pact for Research and Innovation) (Neves & Neves 2011).

However, in an evaluation between authors and co-authors, China holds the highest number of publications, pointing to investments in infrastructure and research financing, by public agencies and private companies. The National Science Foundation (2018), showed that since 2000 China increased its investments in research and development by an annual average of 18% compared to an average of 4% in the same period in the United States. This investment has already brought significant changes since 2018 when China surpassed the United States in the number of publications (Tollefson 2018), a result also observed in our study.

The data obtained in our study agree with those obtained by Gonçalves et al. (2017) demonstrated that China has the highest number of patents involving endophytic fungi applicable in pharmaceutical and cosmetic areas (31 patents, 86% of the documents found). Moreover, the Chinese

territory is rich in biodiversity, with already more than 30,000 documented species of microorganisms, constituting a great natural potential and reservoir for scientific research in strategic sectors that may be of significant interest to the pharmaceutical and biotechnological industries. The goal of these industries is the development and commercialization of innovative medicines, principally for the treatment of chronic sicknesses and infections (Li 2010). Other developing countries, such as Brazil and India, appear in the ranking as those that most published findings verify biological activity among endophytic fungi and therefore also show interest in new research in this strategic area of development. Although Brazil holds second place amongst countries with the highest number of publications, Gonçalves et al. (2017) note that larger investments in the universities are still necessary for the furtherance of new technologies, with few registered patents yet related to endophytic fungi and their applications in the database of the National Institute of Industrial Property (INPI).

The journals that most published articles proving biological activity in endophytic fungi, *Marine Drugs* (3,5%), the *World Journal of Microbiology & Biotechnology* (3,3%) and the *Journal of Natural Products* (3,3%), are responsible for a little more than 10% of the publications that contribute to the biochemistry of endophytes, whether by marine or terrestrial hosts. With respect to the journal of the highest h-index, it was verified that the *European Journal of Organic Chemistry*, between 2005 and 2006 was already in 4th place in the ranking of this index (Bornmann et al. 2009), and thus presented an increase in the number of publications, becoming one of the 20 most relevant journals in the area of organic chemistry.

The United States predominantly orchestrates networks of research collaboration, which deserves to be highlighted, as it presents a disparity of technical and scientific knowledge concerning the bio-prospecting of fungi (Lima et al. 2018), which leads to an increase of cooperation and knowledge exchange with developing countries, notably China, India, and Brazil. The highest number of bio-assays occurred in 2018 (17%), 2019 (16%), 2016 (11%), and 2017 (10%), and were responsible for almost 55% of the total research, demonstrating a tendency of growth.

In almost half of the bio-assays, the active ingredients studied have shown more frequent antimicrobial properties; the clinical importance of pathogenic microorganisms, especially drug-resistant bacteria, justifies the search for substances with novel antimicrobial activities (Schmitt et al. 2011). Moreover, almost 30% of the bio-assays proved to have anti-tumor properties, bringing new perspectives for exploring drugs that fight cancer. Due to the great economic value of these therapies, Chen et al. (2014) relate that between 2010 and 2013, about 100 studies of secondary metabolites of endophytic fungi with these properties were in process, principally in China and India, and at least 30 novel metabolites were isolated.

Conclusions

The analysis of the publications involving biological activity among endophytic fungi revealed a growing trend in the number of studies since the 2000s, with China, Brazil, and India representing the most research countries. In addition, the journals that most published scientific articles with results about bio-activity of endophytic fungi were *Marine Drugs*, the *World Journal of Microbiology & Biotechnology*, the *Journal of Natural Products*, and the *European Journal of Organic Chemistry*, the latter having the highest h-index. The rapid and important increase in the number of bio-assays between 2016 and 2019 points to the evolution of scientific production worldwide and reveals that developing countries have been concerned with seeking new sources of bioactive compounds within endophytic microorganisms. Studies that have proven anti-microbial and anti-tumor properties are highlighted, with the activities most frequently mentioned regarding the possibility of future applications.

Author contribution

IRS, LLB, AMA, and SXS conceived of the presented idea, designed, directed the project wrote the manuscript with support from DDC and JPB. VKG, TAM, HHA, MAA, MMI, and AMA helped

supervise the project. TAM, HHA, MMI and MAA lead the manuscript format, and all authors discussed the results and contributed to the final manuscript format.

Funding

This research was funded by the State University of Goiás, Brazil.

Consent for publication

All the authors have approved the manuscript for publication.

Competing interests

The authors declare no competing interests.

References

- Aria M, Cuccurullo C. 2017 – Bibliometrix: An R-tool for comprehensive science mapping analysis. *J Informetr* 11: 959–975. Doi 10.1016/J.JOI.2017.08.007
- Atanasov AG, Waltenberger B, Pferschy-Wenzig EM, Thomas L et al. 2015 – Discovery and resupply of pharmacologically active plant-derived natural products: A review. *Biotechnol Adv* 33: 1582–1614. Doi 10.1016/J.BIOTECHADV.2015.08.001
- Bornmann L, Marx W, Schier H. 2009 – Hirsch-Type Index Values for Organic Chemistry Journals: A Comparison of New Metrics with the Journal Impact Factor. Doi 10.1002/ejoc.200801243
- Chapla VM, Biasetto CR, Araujo AR. 2013 – Fungos endofíticos: Uma fonte inexplorada e sustentável de novos e bioativos produtos naturais. *Scopus* 5: 421–437
- Chen X, Chen J, Wu D, Xie Y et al 2016b – Mapping the Research Trends by Co-word Analysis Based on Keywords from Funded Project. *Procedia Comput Sci* 91: 547–555. Doi 10.1016/J.PROCS.2016.07.140
- Chen L, Zhang QY, Jia M, Ming QL et al. 2014 – Endophytic fungi with anti-tumor activities: Their occurrence and anticancer compounds. *Critical Reviews in Microbiology* 42(3): 454–473. Doi 10.3109/1040841X.2014.959892
- Chen L, Zhang QY, Jia M, Ming QL et al 2016a – Endophytic fungi with anti-tumor activities: Their occurrence and anticancer compounds. *Crit Rev Microbiol* 42: 454–473. Doi 10.3109/1040841X.2014.959892
- Frank DJ, Meyer JW. 2007 – University expansion and the knowledge society. *Theory Soc* 36: 287–311. Doi 10.1007/S11186-007-9035-Z
- Gangadevi V, Muthumary J. 2009a – A Novel Endophytic Taxol-Producing Fungus *Chaetomella raphigera* Isolated from a Medicinal Plant, *Terminalia arjuna*. *Appl Biochem Biotechnol* 2009 1583 158: 675–684. Doi 10.1007/S12010-009-8532-0
- Gangadevi V, Muthumary J. 2009b – Taxol production by *Pestalotiopsis terminaliae*, an endophytic fungus of *Terminalia arjuna* (arjun tree). *Biotechnol Appl Biochem* 52: 9–15. Doi 10.1042/BA20070243
- Gunatilaka AAL. 2006 – Natural Products from Plant-Associated Microorganisms: Distribution, Structural Diversity, Bioactivity, and Implications of Their Occurrence. *J Nat Prod* 69: 509–526. Doi 10.1021/NP058128N
- Gupta S, Chaturvedi P, Kulkarni MG, Van Staden J. 2020 – A critical review on exploiting the pharmaceutical potential of plant endophytic fungi. *Biotechnol Adv* 39: 107462. Doi 10.1016/J.BIOTECHADV.2019.107462
- Gonçalves B, Bastos E, Hanna S. 2017 – Prospecção tecnológica de fungos endofíticos e aplicações na indústria farmacêutica. *Cadernos de Prospecção* 10(1): 56–67. Doi 10.9771/cp.v10i1.20114
- Jalgaonwala R, Mohite BV, Mahajan R. 2017 – A review: Natural products from plant associated endophytic fungi. undefined

- Jia M, Chen L, Xin HL, Zheng CJ et al. 2016 – A friendly relationship between endophytic fungi and medicinal plants: a systematic review. *Frontiers in Microbiology* 7: 906. Doi 10.3389/fmicb.2016.00906
- Li Y. 2010 – Imitation to innovation in China: the role of patents in biotechnology and pharmaceutical industries. 189.
- Lima RA de, Velho LMLS, Faria LIL. 2018 – Delimitation of a multidisciplinary area for bibliometric analysis of scientific production: study case in the area of Bioprospecting. *Transinformação* 24: 168–153. Doi 10.1590/S0103-37862007000200006
- McGrath W. 1989 – What bibliometricians, scientometricians and informetricians study – a typology for definition and classification, and topics for discussion. In *International Conference on Bibliometrics, Scientometrics and Informetrics, 2. Proceedings...* Ontario: The University of Western Ontario.
- Krings M, Taylor TN, Hass H, Kerp H et al. 2007 – Fungal endophytes in a 400-million-yr-old land plant: infection pathways, spatial distribution, and host responses. *New Phytol* 174: 648–657. Doi 10.1111/J.1469-8137.2007.02008.X
- Macias-Chapula CA. 1998 – O papel da informetria e da cienciometria e sua perspectiva nacional e internacional. *Ciência da Informação* 27: nd-nd. Doi 10.1590/S0100-19651998000200005
- National Science Foundation. 2018 – State of US science enterprise report shows US leads in S&E as China rapidly advances. https://www.nsf.gov/news/news_summ.jsp?cntn_id=244271&org=NSF&from=news (Accessed on August 14, 2021).
- Neves CEB, Neves FM. 2011 – Pesquisa e inovação: novos desafios para an educação superior no Brasil e na Alemanha. *Cad CRH* 24: 481–502. Doi 10.1590/S0103-49792011000300003
- Pandey PK, Singh S, Yadav RNS, Singh AK, Singh MCK. 2014 – Fungal endophytes: Promising tools for pharmaceutical science. *International Journal of Pharmaceutical Sciences Review and Research*. 52(2): 128–138.
- Peixoto Neto PAS, Azevedo JL, Araújo WL. 2002 – Microrganismos endofíticos: interação com plantas e potencial biotecnológico. *Revista Biotecnologia Ciência & Desenvolvimento* 29: 62–76.
- Pons P, Latapy M. 2006 – Journal of Graph Algorithms and Applications Computing Communities in Large Networks Using Random Walks. 10: 191–218
- Rashmi M, Kushveer JS, Sarma VV. 2019 – A worldwide list of endophytic fungi with notes on ecology and diversity. *Mycosphere*.10(1): 798–1079. Doi 10.5943/mycosphere/10/1/19
- Rana KL, Kour D, Sheikh I, Yadav N et al 2019a – Endophytic Fungi: Biodiversity, Ecological Significance, and Potential Industrial Applications. In: Yadav AN, Mishra S, Singh S, Gupta A (eds) *Recent advancement in white biotechnology through fungi, vol 1. Diversity and enzymes perspectives*. Springer, Switzerland, Pp. 1–62. Doi 10.1007/978-3-030-10480-1
- Rana KL, Kour D, Sheikh I, Dhiman A et al 2019b – Biodiversity of Endophytic Fungi from Diverse Niches and Their Biotechnological Applications. In: Singh BP (ed) *Advances in endophytic fungal research: present status and future challenges*. Springer International Publishing, Cham, Pp. 105–144. Doi 10.1007/978-3-030-03589-1_6
- R Core Team. 2020 – A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria <https://www.R-project.org/>
- Redecker D, Kodner R, Graham LE. 2000 – Glomalean fungi from the Ordovician. *Science* 289: 1920–1921. Doi 10.1126/SCIENCE.289.5486.1920
- Rodriguez R, Redman R. 2008 – More than 400 million years of evolution and some plants still can't make it on their own: plant stress tolerance via fungal symbiosis. *J Exp Bot* 59: 1109–1114. Doi 10.1093/JXB/ERM342
- Saikkonen K, Faeth SH, Helander M, Sullivan TJ. 2003 – Fungal endophytes: A Continuum of Interactions with Host Plants. Pp. 319–343. Doi 10.1146/ANNUREV.ECOLSYS.29.1.319
- Saikkonen K, Wäli P, Helander M, Faeth SH. 2004 – Evolution of endophyte–plant symbioses. *Trends Plant Sci* 9: 275–280. Doi 10.1016/J.TPLANTS.2004.04.005

- Schmitt EK, Moore CM, Krastel P, Petersen F. 2011 – Natural products as catalysts for innovation: a pharmaceutical industry perspective. *Curr Opin Chem Biol* 15: 497–504. Doi 10.1016/J.CBPA.2011.05.018
- Schulz B, Boyle C, Draeger S et al 2002 – Endophytic fungi: a source of novel biologically active secondary metabolites. *Mycol Res* 106: 996–1004. Doi 10.1017/S0953756202006342
- Sette LD, Passarini MRZ, Delarmelina C et al 2006 – Molecular characterization and anti-microbial activity of endophytic fungi from coffee plants. *World J Microbiol Biotechnol* 2006 2211 22: 1185–1195. Doi 10.1007/S11274-006-9160-2
- Strobel G, Daisy B. 2003 – Bioprospecting for Microbial Endophytes and Their Natural Products. *Microbiol Mol Biol Rev* 67: 491–502. Doi 10.1128/MMBR.67.4.491-502.2003
- Strobel GA. 2003 – Endophytes as sources of bioactive products. *Microbes Infect* 5: 535–544. Doi 10.1016/S1286-4579(03)00073-X
- Tollefson J. 2018 – China declared world’s largest producer of scientific articles. *Nature* 553: 390. Doi 10.1038/D41586-018-00927-4
- Venieraki A, Dimou M, Katinakis P. 2017 – Endophytic fungi residing in medicinal plants have the ability to produce the same or similar pharmacologically active secondary metabolites as their hosts. *Hell, Plant Prot J* 10: 51–66. Doi 10.1515/HPPJ-2017-0006
- Verbeek A, Debackere K, Luwel M, Zimmermann E. 2002 – Measuring progress and evolution in science and technology – I: The multiple uses of bibliometric indicators. *Int J Manag Rev* 4: 179–211. Doi 10.1111/1468-2370.00083
- Yadav AN. 2018 – Biodiversity and biotechnological applications of host-specific endophytic fungi for sustainable agriculture and allied sectors. *Acta Scientific Microbiology* 1(5): 1–5.
- Zhu F, Lin Y. 2006 – Marinamide, a novel alkaloid and its methyl ester produced by the application of mixed fermentation technique to two mangrove endophytic fungi from the South China Sea. *Chinese Sci Bull* 2006 5112 51: 1426–1430. Doi 10.1007/S11434-006-1426-4